

# Computational Solid State Physics, part II

- **Instructor:** Lilia Boeri
- **Email:** [lilia.boeri@uniroma1.it](mailto:lilia.boeri@uniroma1.it)
- **Webpage:** <https://lboeri.wordpress.com>



**SAPIENZA**  
UNIVERSITÀ DI ROMA

# Outline

## ▶ **Single Particle Approximation:**

- Why is it useful?
- Wavefunction-based Methods: Hartree and Hartree-Fock Approximations

## ▶ **Basics of Density Functional Theory:**

- Hohenberg-Kohn Theorems: Density is the basic Variable.
- Kohn Sham Equations: Practical recipe to derive single-particle equations.
- Approximations for the Exchange and Correlation Energy, Homogeneous
- Electron Gas and beyond

# Density Functional Theory

**Density Functional Theory (DFT)** is considered the **workhorse method** to compute the electronic and vibrational properties of solids, and their response to external perturbations.

## ► Based on two fundamental papers:

- P. Hohenberg and W. Kohn, *Inhomogeneous electron gas*, Phys. Rev. 136, B864 (1964).

*“Density is the Basic Variable”*

- W. Kohn and L. J. Sham, *Self-Consistent Equations Including Exchange and Correlation Effects*, Phys. Rev. 140 A1133 (1965).

*Practical recipe to write down (and solve) effective single-particle equations.*

# Main Concepts Behind DFT:

The **full quantum-many body** problem (electrons + nuclei) in solids and large molecules is **untractable** (too many degrees of freedom)...

1. **Born-Oppenheimer Approximation:** Decouple ions & Electrons.
2. **Self-Consistent Field** (Hartree 1926): Single Particle Approximation  
(Effective single-particle equations).



How to determine the «best» single-particle equations?

**Wave-function-based methods:**

Hartree, Hartree-Fock, QMC

30's

**Density Functional Theory:**

(Ground-state) Density is the  
main variable.

60's  
80's

Si

$$Z = 14$$

$$Z_{val} = 4$$


in the diamond lattice:

FCC lattice,

$$a = 5.43 \text{ \AA}$$

$$\Omega = \frac{a^3}{4}$$

$$4(\{2\bar{2}3\}, \{2\bar{2}3\})$$



$$N_{dim} = N_{at} + N_{el} = 2 + 8 = 10$$

$$\Delta x = 0.1 \text{ \AA}$$

$$\underbrace{\left[ \frac{1}{4} \left( \frac{a}{\Delta x} \right)^3 \right]}_{N_{dim}} = \underline{\underline{10^{46}}} \text{ numbers} \rightarrow \text{EXPERIMENTAL WALL}$$

$\rightarrow N_{points} = N_p$

Independent-particle approximation:

$$\left(N_{\text{points}}\right) \cdot \left(N_{\text{nuclei}} + N_{e^{-}}\right) = \underline{\underline{10^5}} \rightarrow \text{Manageable}$$

Fully interacting system:

$$\left(N_{\text{points}}\right)^{\left(N_{\text{nuclei}} + N_e\right)} = 10^{46} \rightarrow \underline{\underline{\text{Unmanageable!}}}$$



# The Quantum-Many Body Problem:

All properties of a solid can be computed solving the **Quantum Many-Body Problem**, i.e. finding the eigenvalues and eigenstates of the Hamiltonian of interacting Nuclei and Electrons:

$$\hat{H}_{tot} = \hat{T}_N + \hat{T}_e + \hat{V}_{eN} + \hat{V}_{NN} + \hat{V}_{ee}$$

$$\hat{H}_{tot}\Phi(\mathbf{R}, \mathbf{r}) = E\Phi(\mathbf{R}, \mathbf{r}), \quad \mathbf{R} = \{\mathbf{R}_\alpha\}, \quad \mathbf{r} = \{\mathbf{r}_i\}$$

The size of the Hilbert space is **huge**!



# Born-Oppenheimer Approximation:

The **Quantum Many-Body Problem** of interacting nuclei and electrons:

$$\hat{H}_{tot} = \hat{T}_N + \hat{T}_e + \hat{V}_{eN} + \hat{V}_{NN} + \hat{V}_{ee}$$

$$\hat{H}_{tot}\Phi(\mathbf{R}, \mathbf{r}) = E\Phi(\mathbf{R}, \mathbf{r}), \quad \mathbf{R} = \{\mathbf{R}_\alpha\}, \quad \mathbf{r} = \{\mathbf{r}_i\}$$

$10^{23} - 10^4$

Can be **decoupled** due to the different characteristic scales of nuclear and electronic motion ( $\mathbf{M}_N \gg \mathbf{m}_{el}$ ):

► **Factorization of the wavefunction:**

$$\Phi(\mathbf{R}, \mathbf{r}) \simeq \Psi_{el}(\mathbf{r}; \mathbf{R}) X_N(\mathbf{R})$$

Diagram illustrating the factorization of the wavefunction  $\Phi(\mathbf{R}, \mathbf{r}) \simeq \Psi_{el}(\mathbf{r}; \mathbf{R}) X_N(\mathbf{R})$ . The term  $\Psi_{el}(\mathbf{r}; \mathbf{R})$  is associated with **Electrons** (indicated by a blue arrow), and the term  $X_N(\mathbf{R})$  is associated with **Nuclei** (indicated by a red arrow).

# Born-Oppenheimer Approximation:

Factorization of the wavefunction:

$$\Phi(\mathbf{R}, \mathbf{r}) \simeq \underbrace{\Psi_{el}(\mathbf{r}; \mathbf{R})}_{\text{Electrons}} \underbrace{X_N(\mathbf{R})}_{\text{Nuclei}}$$

Effective Equations (Electrons + Nuclei)

*clamped nuclei / ions*

$$\left[ -\frac{1}{2m_e} \nabla_{\mathbf{r}}^2 + V_{eN}(\mathbf{r}; \mathbf{R}) + V_{ee}(\mathbf{r}_{ij}) + V_{NN}(\mathbf{R}_{\alpha\beta}) \right] \Psi_{el}(\mathbf{r}; \mathbf{R}) = E_{El}(\mathbf{R}) \Psi_{el}(\mathbf{r}; \mathbf{R})$$

$$\left[ -\frac{1}{2M_N} \nabla_{\mathbf{R}}^2 + E_{el}(\mathbf{R}) \right] X_N(\mathbf{R}) = E_N X_N(\mathbf{R})$$

*Structural Relaxation*



The electronic Schrodinger's equation is **still too complicated** to solve analytically...



## Self – Consistent Field Approximation (Hartree, 1926)

Replace actual interacting electrons with effective non-interacting quasi-particles, **moving in an effective potential which incorporates the effect of all other (N-1) quasi-particles.**

$$H_{el} \simeq \sum_i h_i^{\text{eff}}$$

$$h_i^{\text{eff}} \phi_i = \varepsilon_i \phi_i$$

The original problem of solving a differential equation of 3N variables is recast into the manageable problem of solving a system of N equations of 3 variables, coupled self-consistently.



# How can we find the self-consistent field?

---

1) Wavefunction based methods



# Wavefunction-based methods:

**Trial wavefunction** + Minimization of the variational energy:

$$E_V[\Psi_T] = \frac{\langle \Psi_T | \hat{H}_{el} | \Psi_T \rangle}{\langle \Psi_T | \Psi_T \rangle}$$

**Hartree:** Trial wavefunction is a simple product of single-particle orbitals:

$$\Psi_{el}(\mathbf{r}; \mathbf{R}) \simeq \phi_1(\mathbf{r}_1; \mathbf{R}) \phi_2(\mathbf{r}_2; \mathbf{R}) \dots \phi_N(\mathbf{r}_N; \mathbf{R})$$

**Hartree-Fock:** Trial wavefunction is a properly antisymmetrized linear combination of Hartree's WFs:

$$\Psi_{HF}(x_1, \dots, x_N) = \frac{1}{\sqrt{N!}} \sum_{P=1}^{N!} (-1)^P \hat{P} [\psi_\alpha(x_1) \dots \psi_\nu(x_N)] = \frac{1}{\sqrt{N!}} \sum_{P=1}^{N!} (-1)^P \hat{P} [\Psi_H^{\alpha, \dots, \nu}]$$

# Hartree's Approximation:

Ansatz for the wavefunction:

$$\Psi_{el}(\mathbf{r}; \mathbf{R}) \simeq \phi_1(\mathbf{r}_1; \mathbf{R}) \phi_2(\mathbf{r}_2; \mathbf{R}) \dots \phi_N(\mathbf{r}_N; \mathbf{R})$$

quantum state

\* electron

PAULI  
exclusion

Variational minimization of the electronic energy yields a set of **N self-consistent equations**:

$$\left[ -\frac{\nabla_i^2}{2m_e} + V_{eN}(\mathbf{r}_i) + V_{coul}(\mathbf{r}_i) \right] \phi_i(\mathbf{r}_i) = \varepsilon_i \phi_i(\mathbf{r}_i)$$

Self-consistent  
potential


$$V_{coul}(\mathbf{r}_i) = \sum_{j_{occ}} \int d^3r' \frac{\phi_j^*(\mathbf{r}') \phi_j(\mathbf{r}')}{|\mathbf{r}' - \mathbf{r}_i|} - \int d^3r' \frac{\phi_i^*(\mathbf{r}') \phi_i(\mathbf{r}')}{|\mathbf{r}' - \mathbf{r}_i|}$$

Average Electrostatic  
Repulsion

Self-interaction

# Self-Consistent Field:

Start



# Hartree's Approximation:

## Ansatz for the wavefunction:

$$\Psi_{el}(\mathbf{r}; \mathbf{R}) \simeq \phi_1(\mathbf{r}_1; \mathbf{R})\phi_2(\mathbf{r}_2; \mathbf{R})\dots\phi_N(\mathbf{r}_N; \mathbf{R})$$

## Properties of the Hartree wavefunction:

- The Pauli exclusion principle is enforced «by hand» (one particle in one orbital).
- Accounts for average (electrostatic) effect of the Coulomb interaction (screening).
- No **correlation** in the position of the effective (quasi)-particles.

## (Serious) Drawback:

- Enforcing the exclusion principle by hand does not respect symmetry properties of electronic (fermionic wavefunction).



# Hartree-Fock Approximation:

## Fermionic Statistics (Exchange):

Improve the quality of the trial wavefunction by accounting correctly for fermionic statistics :

$$\Psi(x_1, \dots, x_j, \dots, x_k, \dots, x_N) = -\Psi(x_1, \dots, x_k, \dots, x_j, \dots, x_N)$$

The HF **trial wavefunction** is a properly **antisymmetrized** and renormalized linear combination of Hartree's wavefunctions:

$$\Psi_{HF}(x_1, \dots, x_N) = \frac{1}{\sqrt{N!}} \sum_{P=1}^{N!} (-1)^P \hat{P} [\psi_\alpha(x_1) \dots \psi_\nu(x_N)] = \frac{1}{\sqrt{N!}} \sum_{P=1}^{N!} (-1)^P \hat{P} [\Psi_H^{\alpha, \dots, \nu}]$$

$$\Psi_H(x_1, \dots, x_N) = \psi_\alpha(x_1) \dots \psi_\nu(x_N)$$

# Hartree-Fock Approximation:

**Slater determinant** permits to write the trial wavefunction in compact and transparent form:

$$\Psi_{HF}(x_1, \dots, x_N) = \frac{1}{\sqrt{N!}} \begin{vmatrix} \psi_\alpha(x_1) & \psi_\alpha(x_2) & \dots & \psi_\alpha(x_N) \\ \psi_\beta(x_1) & \psi_\beta(x_2) & \dots & \psi_\beta(x_N) \\ \vdots & \vdots & \ddots & \vdots \\ \psi_\nu(x_1) & \psi_\nu(x_2) & \dots & \psi_\nu(x_N) \end{vmatrix}$$

→ orbitals

electron

$$\psi_\alpha(x_i) = \psi_\alpha(\mathbf{r}_i, \sigma_i) = \phi_\alpha(\mathbf{r}_i)\chi_\alpha(\sigma_i)$$

- Putting two particles in the same spin-orbital is not allowed (determinant = 0).
- Exchanging two particles the sign of the N-body wavefunction changes.
- There is a **correlation** between electrons with the **same spin**.

# Hartree-Fock Approximation:

Variational minimization with respect to a single orbital ( $\gamma$ ) leads to the **effective single-particle** HF equations:

Direct (Hartree) Term

$$\left[ -\frac{\hbar^2 \nabla_1^2}{2m} + v_{ei}(\mathbf{r}_1) \right] \psi_\gamma(x_1) + \left[ \sum_\mu \int dx_2 \psi_\mu^*(x_2) \frac{1}{r_{12}} \psi_\mu(x_2) \right] \psi_\gamma(x_1) +$$

$$- \sum_\mu \left[ \int dx_2 \psi_\mu^*(x_2) \frac{1}{r_{12}} \psi_\gamma(x_2) \right] \delta_{\sigma_\gamma \sigma_\mu} \psi_\mu(x_1) = \varepsilon_\gamma \psi_\gamma(x_1)$$

Exchange (Fock) Term

(couples e- with the same spin) – favours FM

$$\hat{H}_{HF} |\psi_\gamma\rangle = \varepsilon_\gamma |\psi_\gamma\rangle$$

$$\hat{H}_{HF} = \hat{T}_e + \hat{v}_{ei} + \hat{v}_H [\{\psi_\gamma\}] + \hat{v}_x [\{\psi_\gamma\}]$$

# Hartree-Fock Approximation:

## Ansatz for the wavefunction:

$$\Psi_{HF}(x_1, \dots, x_N) = \frac{1}{\sqrt{N!}} \sum_{P=1}^{N!} (-1)^P \hat{P} [\psi_\alpha(x_1) \dots \psi_\nu(x_N)] = \frac{1}{\sqrt{N!}} \sum_{P=1}^{N!} (-1)^P \hat{P} [\Psi_H^{\alpha, \dots, \nu}]$$

## Advantages of the Hartree-Fock Approximation:

- Correctly accounts for fermionic anisymmetry of the wavefunction.
- Accounts for average (electrostatic) effect of the Coulomb interaction (screening) + effect of fermionic statistics (exchange).
- Koopman's theorem (HF orbital energies approximate ionization potentials).

## Drawbacks:

- Scales very badly with system size ( $N^4$ ).
- Does not include correlation effects beyond exchange (dynamic e-e avoidance, strong correlation requires multiple determinants)

# Effects of Electron-Electron Interaction:

## 1) Average Electrostatic Repulsion

Hartree

Classical

Screening of ionic potentials

## 2) Effect of Fermionic Statistics (Exchange Interaction)

Hartree-Fock

Quantum

Magnetism

## 3) Dynamic Interactions between all e- (Correlation)

Beyond H-F

Quantum

Binding energy, cohesion of solids

CONFIGURATION INTERACTION

Imp. v. Function  $\rightarrow$  QMC

$\rightarrow$  DENSITY FUNCTIONAL THEORY

# How can we find the self-consistent field?



2) Density Functional Theory

# Density Functional Theory:

1) **Density is the main variable:** the ground-state properties of a system of interacting electrons are uniquely determined by its **ground-state charge Density**.

$$n^0(\mathbf{r}) = N_e \int d^3r_2 \dots d^3r_N |\Psi_{el}^0(\mathbf{r}, \mathbf{r}_2, \dots, \mathbf{r}_N)|^2$$

P. Hohenberg and W. Kohn, *Inhomogeneous electron gas*, Phys. Rev. **136**, B864 (1964).

2) The **ground-state charge density** can be obtained solving a system of **self-consistent equations** for an **auxiliary system** of non-interacting quasi-particles (**Kohn-Sham electrons**):


$$n^0(\mathbf{r}) = \sum_i |\phi_i^{KS}(\mathbf{r})|^2$$

$$h^{KS} [n^0(\mathbf{r})] \phi_i^{KS}(\mathbf{r}) = \varepsilon_i \phi_i^{KS}(\mathbf{r})$$


W. Kohn and L. J. Sham, *Self-Consistent Equations Including Exchange and Correlation Effects*, Phys. Rev. **140** A1133 (1965).

# Kohn-Sham Equations:


- Single-Particle Equations (Kohn-Sham):


$$\left( -\frac{1}{2}\nabla^2 + v_{ext}(\mathbf{r}) + \int \frac{n(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}' + v_{xc}[n(\mathbf{r})] \right) \phi_i(\mathbf{r}) = \epsilon_i \phi_i(\mathbf{r})$$

- Self-Consistency Condition:



$$n_0(\mathbf{r}) = \sum_{i \in \text{occ}} |\phi_i(\mathbf{r})|^2$$

EXCHANGE AND correlation  
Potential


$$v_{\text{eff}}(\mathbf{r}) = v_{\text{ext}}(\mathbf{r}) + v_{\text{H}}[n(\mathbf{r})] + v_{\text{xc}}[n(\mathbf{r})]$$



# Self-Consistent Solution of Kohn-Sham Equations:



# Kohn-Sham Equations:

- Effective Potential (Kohn-Sham)

$$v_{\text{eff}}(\mathbf{r}) = v_{\text{ext}}(\mathbf{r}) + v_{\text{H}}[n(\mathbf{r})] + v_{\text{xc}}[n(\mathbf{r})]$$

$$v_{\text{ext}}(\mathbf{r})$$

**External Potential** (Nuclei + other external fields)

$$v_{\text{H}}[n(\mathbf{r})]$$

**Hartree Potential** (Direct Electrostatic Repulsion)

$$v_{\text{xc}}[n(\mathbf{r})]$$

**Exchange and Correlation Potential:** Contains all effects of e-e interaction beyond Hartree (unknown)

*It exists!*

# Exchange and Correlation Potential

$$v_{xc}[n(\mathbf{r})]$$

**Exchange and Correlation Potential:** Contains all effects of e-e interaction beyond Hartree (unknown)

## Local Density Approximation:

Assume that the exchange and correlation energy of the real system is **locally** the same as that of the **homogeneous electron gas**:

$$v_{xc}(\mathbf{r}) = \frac{\delta E_{xc}[n]}{\delta n(\mathbf{r})}$$

$$E_{xc}[n(\mathbf{r})] = \int n(\mathbf{r}) \epsilon_{xc}(n(\mathbf{r})) d\mathbf{r}$$

$$\epsilon_{xc}(n(\mathbf{r})) = \epsilon_x(n(\mathbf{r})) + \epsilon_c(n(\mathbf{r}))$$

ANALYTICALLY  
NUMERICALLY