

KET QUẢ SINH HOA MÁU



Patient:	Lu	Species:	Canine	Patient ID:	
Client:	Thy	Gender:	Female	Sample No.:	66
Doctor:		Age:	Adult	Time of analysis:	2025/12/15 08:53

Item		Current result		Ref. Ranges	
Protein	TP	↑	88.3	g/L	53.1-79.2
Protein	ALB	↓	22.8	g/L	23.4-40.0
Protein	GLOB	↑	65.6	g/L	25.4-52.0
Protein	A/G		0.3		
Liver and gallbladder	ALT	↑	>2000.0	U/L	10.1-100.3
Liver and gallbladder	AST	↑	436.3	U/L	0.0-51.7
Liver and gallbladder	AST/ALT		****		
Liver and gallbladder	ALP	↑	1758.9	U/L	15.5-212.0
Liver and gallbladder	GGT	↑	82.4	U/L	0.0-15.9
Liver and gallbladder	TBIL	↑	113.86	μmol/L	0.00-15.00
Liver and gallbladder	TBA	↑	>110.0	μmol/L	0.0-30.0
Pancreas	AMY		512.0	U/L	397.7-1285.1
Kidneys	BUN		9.62	mmol/L	2.50-9.77
Kidneys	CREA		80.10	μmol/L	20.00-123.70
Kidneys	BUN/CREA		29.7		
Cardiovasc./Muscle	CK	↑	276.5	U/L	66.4-257.5
Cardiovasc./Muscle	LDH	↑	210.6	U/L	0.0-143.6
Energy metabolism	GLU	↑	10.84	mmol/L	3.80-7.50
Energy metabolism	TC	↑	13.79	mmol/L	2.67-8.38
Energy metabolism	TG	↑	1.32	mmol/L	0.10-1.30
Minerals	Ca		2.21	mmol/L	2.10-2.97
Minerals	PHOS	↓	0.71	mmol/L	0.80-2.20
Minerals	CaxP		1.57	mmol/L^2	
Minerals	Mg		0.90	mmol/L	0.53-1.06
Electrolytes	Na+		141.7	mmol/L	138.0-160.0
Electrolytes	K+	↓	3.4	mmol/L	3.5-5.9
Electrolytes	Na/K		41.9		
Electrolytes	Cl-		120.1	mmol/L	102.7-125.0

Operator:

Comprehensive Diagnosis Panel			QC QC OK	
HEM(Hemolysis degree):	0	LIP(Lipemia degree):	1+	ICT(Jaundice degree): 2+

Report Explen.

TP



Increase is commonly associated with dehydration and increased globulin. Reduction is commonly associated with blood loss, protein-losing enteropathy, and decreased albumin.

The results only applies to this test sample. Time of Printing:2025-12-14 18:55:12

KET QUA SINH HOA MAU



Patient:	Lu	Species:	Canine	Patient ID:	
Client:	Thy	Gender:	Female	Sample No.:	66
Doctor:		Age:	Adult	Time of analysis:	2025/12/15 08:53

Report Explan.		
ALB	↓	Increase is commonly associated with dehydration and corticosteroid administration, etc. Reduction is commonly associated with excessive infusion, malnutrition, hepatic insufficiency or failure, nephropathy, and protein-losing enteropathy.
GLOB	↑	Increase is commonly associated with chronic inflammation and infection, and hyperimmunity, etc. Reduction is commonly associated with insufficient protein intake, anemia, and immunodeficiency.
ALT	↑	Increase is commonly associated with liver injury and muscle injury, etc.
AST	↑	Increase is commonly associated with liver injury and muscle injury, etc.
ALP	↑	Increase is commonly associated with fracture healing period, hepatobiliary diseases, hyperthyroidism, and osteosarcoma, etc.
GGT	↑	Elevated is commonly associated with bile duct injury or cholestasis, etc.
TBIL	↑	Increase is commonly associated with hemolysis and hepatobiliary dysfunction. Reduction is commonly associated with decreased erythropoiesis, etc.
TBA	↑	Increase is commonly associated with hepatic insufficiency or failure, portal vein shunt, and cholestasis, etc. Reduction is commonly associated with long-term fasting and intestinal malabsorption, etc.
CK	↑	Increase is commonly associated with trauma, increased muscle activity (such as tetanus and convulsion), myocarditis, and myocardial infarction, etc.
LDH	↑	Increase is commonly associated with hemolysis (especially in canine), post-exercise, liver injury, exertional rhabdomyolysis, white muscle disease, myocardial injury, tumors, etc.
GLU	↑	Increase is commonly associated with diabetes and hypercorticism, etc. Reduction is commonly associated with insulin administration, malnutrition, and insulinoma, etc.
TC	↑	Increase is commonly associated with biliary obstruction, hypothyroidism, hypercorticism, nephropathy, diabetes, etc. Reduction is commonly associated with protein loss enteropathy, pancreatic exocrine insufficiency, and hypoadrenocorticism, etc.
TG	↑	Increase is commonly associated with postprandial, obesity, diabetes and hypercorticism, etc.
PHOS	↓	Increase is commonly associated with nephropathy, bone healing period, and hyperthyroidism. Decreased in hyperparathyroidism, tumor, etc.
K+	↓	Increase is commonly associated with high potassium fluid replacement, diabetes, adrenocortical hypofunction, and acute kidney injury, etc. Reduction is commonly associated with low potassium or potassium-free fluid replacement, vomiting, diarrhea, and hypercorticism, etc.

Note: Due to the complexity and individuality of disease diagnosis, the report interpretation is only for your reference. Please consult your doctors for clinical diagnosis results.
The results only applies to this test sample.

Time of Printing:2025-12-14 18:55:12



PHONG KHAM THU Y VETGO – GO DUA
57 GO DUA, P. TAM BINH, TP. HO CHI MINH
0903.389.624 – 0867.483.384

Global Pioneer of Comprehensive Animal Medical Solutions
Better healthcare for all - Since 1991

