

## Your grade: 80%

Your latest: 80% • Your highest: 80% • To pass you need at least 70%. We keep your highest score.

[Next item →](#)

1. Which one of the following is common to both machine learning and statistical inference?

1 / 1 point

- ☒ Using sample data to infer qualities of the underlying population distribution.
- ☐ Using sample data to make inferences about a hypothesis.
- ☐ Using population data to model a null hypothesis.
- ☐ Using population data to make inferences about a null sample.

 ✓ **Correct**

Correct. In both machine learning and statistical inference, we're using sample data to infer qualities of the underlying population distribution.

2. Which one of the following describes an approach to customer churn prediction stated in terms of probability?

1 / 1 point

- ☒ Predicting a score for individuals that estimates the probability the customer will leave.
- ☐ Data related to churn may include the target variable for whether a certain customer has left.
- ☐ Churn prediction is a data-generating process representing the actual joint distribution between our x and the y variable.
- ☐ Predicting a score for individuals that estimates the probability the customer will stay.

 ✓ **Correct**

Correct. Churn prediction is often approached by predicting a score for individuals that estimates the probability the customer will leave.

3. What is customer lifetime value?

1 / 1 point

- ☒ The total purchases over the time which the person is a customer.
- ☐ The total value that the customer receives during their life.
- ☐ The total churn generated by a customer over their lifetime.
- ☐ The total churn a customer generates in the population.

 ✓ **Correct**

Correct. Customer lifetime value consists of the purchase amounts over the entire time that a person has been a customer.

4. Which one the following statements about the normalized histogram of a variable is true?

1 / 1 point

- ☐ It is a parametric representation of the population distribution.
- ☒ It provides an estimate of the variable's probability distribution.
- ☐ It is a non-parametric representation of the population variance.
- ☐ It serves as a bar chart for the null hypothesis.

 ✓ **Correct**

Correct. The normalized histogram of a variable estimates the variable's probability distribution, and the estimate improves with the amount of data used.

5. The outcome of rolling a fair die can be modelled as a \_\_\_\_\_ distribution.

1 / 1 point

- ☐ log-normal
- ☐ Poisson
- ☒ uniform
- ☐ normal

 ✓ **Correct**

Correct. The chance of rolling any particular value for a fair die is equally likely, so the outcome is uniformly distributed.

6. Which one of the following features best distinguishes the Bayesian approach to statistics from the Frequentist approach?

1 / 1 point

- ☐ Bayesian statistics is better than Frequentist.
- ☒ Bayesian statistics incorporate the probability of the hypothesis being true.
- ☐ Frequentist statistics incorporates the probability of the hypothesis being true.
- ☐ Frequentist statistics requires construction of a prior distribution.



**Correct**

Correct. Bayesian statistics allows for experimenters to incorporate their prior beliefs of the [population] distribution [of a given variable]. For frequentists, it's solely based on the data available. [that is, there is no formal mechanism in frequentist statistics for incorporating prior knowledge, one 'lets the data do the talking']

7. Which of the following best describes what a hypothesis is?

1 point

- ☒ A hypothesis is a statement about a sample of the population.
- ☐ A hypothesis is a statement about a prior distribution.
- ☐ A hypothesis is a statement about a population.
- ☐ A hypothesis is a statement about a posterior distribution.



**Incorrect**

Incorrect. Review the Introduction to Hypothesis video.

8. A Type 2 error in hypothesis testing is \_\_\_\_\_:

1 / 1 point

- ☐ correctly rejecting the alternative hypothesis.
- ☐ incorrectly accepting the alternative hypothesis.
- ☒ incorrectly accepting the null hypothesis.
- ☐ correctly rejecting the null hypothesis.



**Correct**

Correct. A type 2 error is incorrectly accepting the null hypothesis.

9. Which statement best describes a consequence of a type II error in the context of a churn prediction example? Assume that the null hypothesis is that customer churn is due to chance, and that the alternative hypothesis is that customers enrolled for greater than two years will not churn over the next year.

1 point

- ☐ You correctly conclude that a customer will eventually churn
- ☐ You correctly conclude that customer churn is by chance
- ☒ You incorrectly conclude that there is no effect
- ☐ You incorrectly conclude that customer churn is by chance



**Incorrect**

Incorrect. Review the Type 1 vs Type 2: Example video.

10. Which of the following is a statistic used for hypothesis testing?

1 / 1 point

- ☐ The standard deviation.
- ☐ The acceptance region.
- ☒ The likelihood ratio.
- ☐ The rejection region.



**Correct**

Correct. The likelihood ratio can be used as a test statistic, to decide whether to accept or reject the null hypothesis.