

Math 135, Spring 2022

Lecture #22: PDEs and boundary value problems

Wednesday May 18th

Recap

- We finished the topic “Fourier Series”.

So far we have covered

- Existence and uniqueness theory for ODEs
- Laplace transform
- Fourier series

And we are now going to deal with

- Partial Differential Equations (PDEs)

Learning objectives

Today we will discuss:

- Fourier series in an arbitrary domain.
- Derivation of the 1d wave equation as a model for a vibrating string.
- Solving the initial boundary value problem for the 1d wave equation by separation of variables.

Last topic on Fourier series

Scaling

Suppose that $f(x)$ is defined on $[-L, L]$, where $L > 0$.
We want to construct the Fourier series for $f(x)$.

Let $g(t) = f(Lt/\pi)$ be defined on $[-\pi, \pi]$

The Fourier series for $g(t)$ is

$$\frac{1}{2}a_0 + \sum_{n=1}^{\infty} \{a_n \cos(nt) + b_n \sin(nt)\}$$

where

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} g(t) \cos(nt) dt \quad \text{and} \quad b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} g(t) \sin(nt) dt$$

As $x = Lt/\pi$, take our Fourier series for $f(x)$ to be

$$t = \frac{\pi}{L} x$$

$$\frac{1}{2} a_0 + \sum_{n=1}^{\infty} \left\{ a_n \cos\left(n \frac{\pi x}{L}\right) + b_n \sin\left(n \frac{\pi x}{L}\right) \right\}$$

where

$$t = \frac{\pi}{L} x \rightarrow dt = \frac{\pi}{L} dx$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f\left(\frac{Lt}{\pi}\right) \cos(nt) dt = \frac{1}{L} \int_{-L}^L f(x) \cos\left(n \frac{\pi x}{L}\right) dx$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f\left(\frac{Lt}{\pi}\right) \sin(nt) dt = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n \pi x}{L}\right) dx$$

Definition: Let $f(x)$ be an integrable function on the interval $[-L, L]$, where $L > 0$. We define the **Fourier series** for $f(x)$ on $[-L, L]$ to be


$$\longrightarrow \frac{1}{2}a_0 + \sum_{n=1}^{\infty} \left[a_n \cos\left(\frac{n\pi}{L}x\right) + b_n \sin\left(\frac{n\pi}{L}x\right) \right],$$

where

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi}{L}x\right) dx \quad \text{and} \quad b_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi}{L}x\right) dx.$$

odd

An example



Find the Fourier series for $f(x) = x$ on $[-1, 1]$.

Step 1: Use any available symmetries to simplify the computation.

$a_n = 0$ for all n as $f(x)$ is odd

Step 2: Compute

$$b_n = 2 \int_0^1 x \sin(n\pi x) dx$$

$$b_n = \int_{-1}^1 \underbrace{x \sin(n\pi x)}_{\text{even}} dx = 2 \int_0^1 x \sin(n\pi x) dx$$

As $n \geq 1$,


$$b_n = \left[-\frac{2}{n\pi} x \cos(n\pi x) \right]_{x=0}^{x=1} + \int_0^1 \frac{2}{n\pi} \cos(n\pi x) dx$$

$$= -\frac{2}{n\pi} (-1)^n$$

$$= -\frac{2}{n\pi} (-1)^{n+1}$$

So, the Fourier series for $f(x)=x$ on $[-1,1]$ is

$$\sum_{n=1}^{\infty} \frac{2}{n\pi} (-1)^{n+1} \sin(n\pi x)$$



$$f(x) = x$$

$$S_4(x) = \sum_{n=1}^4 (-1)^{n+1} \frac{2}{n\pi} \sin(nx)$$


PDEs and boundary value problems

The vibrating string

Let $y=y(x,t)$ be the vertical displacement of a tight string constraint at its ends

Assumptions:

- Motion is strictly vertical
- No resistance to bending
- Small vibrations
- Constant density ρ
- Constant tension T



The 1d wave equation

We are lead to consider the **1d wave equation**:

$$y = y(x, t)$$

$$\frac{\partial^2 y}{\partial t^2} = c^2 \frac{\partial^2 y}{\partial x^2}$$


$$c^2 = \frac{T}{\rho}$$

with the **boundary conditions**

$$y(t, 0) = 0 = y(t, \pi)$$

and the **initial conditions**

$$y(0, x) = f(x) \quad \text{and} \quad \frac{\partial y}{\partial t}(0, x) = g(x).$$



Linearity

Theorem: If $y_1(t, x)$ and $y_2(t, x)$ are solutions of the linear wave equation

$$\frac{\partial^2 y}{\partial t^2} = c^2 \frac{\partial^2 y}{\partial x^2}$$

with the **boundary conditions**

$$y(t, 0) = 0 = y(t, \pi)$$

and C_1, C_2 are constants, then so is

$$y(t, x) = C_1 y_1(t, x) + C_2 y_2(t, x).$$

Proof: Easy computation.

Separation of variables

$$\begin{cases} \partial_{tt}^2 y = c^2 \partial_{xx}^2 y \\ y(0,t) = y(\pi,t) = 0 \end{cases}$$

Try to find a solution of the form

$$y(x,t) = v(t)u(x) \quad \text{where } u(0) = u(\pi) = 0$$

(educated guess)

Plug this into: $v''(t)u(x) = c^2 v(t)u''(x)$

$$\frac{1}{c^2} \frac{v''(t)}{v(t)} = \frac{u''(x)}{u(x)} = -\lambda \quad \text{a constant!}$$

Spatial oscillations

$$\frac{u''(x)}{u(x)} = -\lambda$$

If $\lambda < 0$, what is the general solution of

boundary
value problem

$$u''(x) + \lambda u(x) = 0 \quad ?$$

$$u(0) = u(\pi) = 0$$

A) $C_1 e^{\sqrt{-\lambda}x} + C_2 e^{-\sqrt{-\lambda}x}$

B) $C_1 \cos(\sqrt{-\lambda}x) + C_2 \sin(\sqrt{-\lambda}x)$

C) $C_1 x + C_2$

D) None of the above

$$u(x) = e^{\mu x}$$

Charact. poly.: $\mu^2 + \lambda = 0 \Rightarrow \mu^2 = -\lambda > 0$

$$\Rightarrow \mu = \pm \sqrt{-\lambda}$$

$$u'' + \lambda u = 0$$

$$\text{If } \lambda < 0: \quad u(x) = c_1 e^{\sqrt{-\lambda} x} + c_2 e^{-\sqrt{-\lambda} x}$$

Boundary conditions:

$$0 = u(0) = c_1 + c_2 \rightarrow c_2 = -c_1$$

$$\begin{aligned} 0 = u(\pi) &= c_1 e^{\sqrt{-\lambda} \pi} + c_2 e^{-\sqrt{-\lambda} \pi} \\ &= c_1 \left(\underbrace{e^{\sqrt{-\lambda} \pi} - e^{-\sqrt{-\lambda} \pi}}_{\neq 0} \right) \rightarrow c_1 = 0 \\ &\quad \rightarrow c_2 = 0 \end{aligned}$$

No non-trivial solutions.

If $\lambda > 0$, what is the general solution of

$$u''(x) + \lambda u(x) = 0 \quad ?$$

- A) $C_1 e^{\sqrt{\lambda}x} + C_2 e^{-\sqrt{\lambda}x}$
- B) $C_1 \cos(\sqrt{\lambda}x) + C_2 \sin(\sqrt{\lambda}x)$
- C) $C_1 x + C_2$
- D) None of the above

Eigenfunctions

- We have shown that we only have non-trivial solutions of the **boundary value problem**

$$\begin{cases} u'' + \lambda u = 0 \\ u(0) = 0 = u(\pi) \end{cases}$$

when $\lambda = n^2$ for $n = 1, 2, \dots$

- We can think of $-\frac{d^2}{dx^2}$, together with the boundary conditions $u(0) = 0 = u(\pi)$ as a **linear operator**, i.e. $-\frac{d^2}{dx^2} : \mathcal{X} \rightarrow \mathcal{Y}$ where \mathcal{X}, \mathcal{Y} are certain space functions.
- We call the functions

$$u_n(x) = \sin(nx)$$

eigenfunctions of this operator and we call the values $\lambda_n = n^2$ the corresponding **eigenvalues**.

Time oscillation

What is the general solution of

$$v''(t) + c^2 n^2 v(t) = 0 \quad ?$$

- A) $C_1 e^{nct} + C_2 e^{-nct}$
- B) $C_1 \cos(nct) + C_2 \sin(nct)$
- C) $C_1 nct + C_2$
- D) None of the above

- Suppose we take $\frac{\partial y}{\partial t}(0, x) = g(x) = 0$.
- Consequently, for each $n \geq 1$ we have a corresponding solution of

$$\begin{cases} v'' + n^2 c^2 v = 0 \\ v'(0) = 0 \end{cases}$$

given by

$$v_n(t) = \cos(nct).$$

Solving our PDE

See you next time!