



Variance Moderation of Locally Efficient Estimators and Supervised Clustering with Applications in High-Dimensional Biology

NIMA S. HEJAZI, MARK J. VAN DER LAAN, & ALAN E. HUBBARD *Graduate Group in Biostatistics*



School of
Public Health

UNIVERSITY OF CALIFORNIA, BERKELEY

OVERVIEW & MOTIVATIONS

- A general approach for applying variance moderation to locally efficient estimators in nonparametric models is introduced.
- While relevant estimation procedures are usually data-intensive, variance moderation stabilizes their small-sample properties, unlocking their use for differential expression analyses.
- Focusing on a targeted maximum likelihood estimator (TMLE) of the average treatment effect (ATE), we illustrate how the approach generalizes to asymptotically linear estimator.
- Supervised clustering is discussed in the context of influence functions, ...
- The `biotmle` R package [1] implements the inference and clustering approaches while leveraging state-of-the-art machine learning in the procedure.

DATA: BENZENE BIOMARKERS

- With the growing number of methods for measuring biomarkers there arises a need for methodologies able to simultaneously analyze multiple kinds of exposome data.
- Data was generated by the **Illumina Human Ref-8 BeadChips** platform.
- There were 125 subjects, for which background characteristics and expression measures for $\sim 22,000$ genes were obtained.
- Covariates in W were age, sex, and smoking status; all were discretized.
- The treatment (A) is degree of Benzene exposure: none, <1ppm, and >5ppm.
- The outcome (Y) is a vector of gene expression measures, normalized by median.

METHODOLOGY II: SUPERVISED DISTANCE MATRICES

- Let $\phi : (W, A, Y) \mapsto D_b(P_0)(O)$ be the EIF transformation, where $D_b(P_0)(O_i)$ is the contribution of subject i to the estimate of the biomarker-specific target parameter $\Psi_{b,n}$.
- $Z = \phi(W, A, Y)$ is then a $B \times N$ matrix, where each entry (b, i) may be interpreted as the degree to which subject i deviates from the target parameter $\Psi_{b,n}$ [2], and is thus an *association profile*.
- A *supervised distance matrix* [3] may be constructed by applying a distance metric of choice (e.g., Euclidean, correlation) to the transformed values Z .
- $\tilde{T}(Z)$, the resultant $B \times B$ empirical distance matrix, encodes the dissimilarity between pairs of biomarker association profiles.
- When $\tilde{T}(b, b') = g(Z_b, Z_{b'})$ is small, biomarkers b and b' have similar associations between expression and the target parameter Ψ , across the n subjects.
- *Supervised clustering* may be performed by applying standard clustering algorithms to the matrix \tilde{T} , thereby finding groups of biomarkers that share an association profile w.r.t. Ψ .
- In the case of the average treatment effect, a supervised cluster in \tilde{T} of biomarkers is a group whose causal differential expression profiles varies similarly with the treatment $A \in \{0, 1\}$.

METHODOLOGY I: VARIANCE MODERATION & LOCAL EFFICIENCY

- Let observed data $O = (W, A, Y) \sim P_0 \in \mathcal{M}$, where W represents potential baseline confounders, A the exposure of interest, and $Y = (Y_b, b = 1, \dots, B)$ a vector of potential biomarkers.
- We consider, as an example, the *average treatment effect* (ATE), as the causal parameter of interest, which is identified by the observed data parameter:

$$\Psi_b(P_0) = \mathbb{E}_W[Q_0^b(A = 1, W) - Q_0^b(A = 0, W)], \quad (1)$$

where $Q_0^b(A, W) \equiv \mathbb{E}_{P_0}(Y_b | A, W)$ and may be estimated via *ensemble machine learning* [4, 5, 6].

- Like the estimator $\hat{\beta}$ in a linear model $m(A, W | \beta)$, $\Psi_b(P_n)$ is **asymptotically linear** (for Ψ_b) [7]:

$$\sqrt{n}(\Psi_b(P_n) - \Psi_b(P_0)) = \frac{1}{\sqrt{n}} \sum_{i=1}^n D_b(O_i) + o_p(1). \quad (2)$$

- Ψ_b has efficient influence function (EIF), relative to the nonparametric model \mathcal{M} :

$$D_b(P_0)(o) = \left(\frac{I(a=1)}{g(1|w)} - \frac{I(a=0)}{g(0|w)} \right) \cdot [y_b - Q_0^b(a, w)] + (Q_0^b(1, w) - Q_0^b(0, w) - \Psi_b(P_0)(o)). \quad (3)$$

- A moderated test statistic [8, 9, 2] may be constructed for use with asymptotically linear estimators:

$$\tilde{t}_b = \frac{\sqrt{n}(\Psi_b(P_n) - \psi_0)}{S_b(D_{b,n})} \quad \text{where} \quad \tilde{S}_{b,n}^2 = \frac{d_0 S_0^2 + d_b S_b^2(D_{b,n})}{d_0 + d_b}, \quad (4)$$

$\{S_b, d_b\}$: var. EIF and df for b^{th} biomarker; $\{S_0, d_0\}$: var. EIF and df for other $(B-1)$ biomarkers.

RESULTS & DISCUSSION




Figure 1: A supervised heatmap of biomarker association profiles with the ATE.




Figure 2: Improved error rate control of the variance moderated TMLE for the ATE.

- Concluding comment...
- Concluding comment...

REFERENCES

- [1] N. S. Hejazi, W. Cai, and A. E. Hubbard, "biotmle: Targeted learning for biomarker discovery," *The Journal of Open Source Software*, vol. 2, no. 15, July 2017.
- [2] N. S. Hejazi, S. Kherad-Pajouh, M. J. van der Laan, and A. E. Hubbard, "Supervised variance moderation of locally efficient estimators in high-dimensional biology," 2018+.
- [3] K. S. Pollard and M. J. van der Laan, "Supervised distance matrices," *Statistical applications in genetics and molecular biology*, vol. 7, no. 1, 2008.
- [4] M. J. van der Laan, E. C. Polley, and A. E. Hubbard, "Super Learner," *Statistical applications in genetics and molecular biology*, vol. 6, no. 1, 2007.
- [5] L. Breiman, "Stacked regressions," *Machine learning*, vol. 24, no. 1, pp. 49–64, 1996.
- [6] D. H. Wolpert, "Stacked generalization," *Neural networks*, vol. 5, no. 2, pp. 241–259, 1992.
- [7] M. J. van der Laan and S. Rose, *Targeted learning: causal inference for observational and experimental data*. Springer Science & Business Media, 2011.
- [8] G. K. Smyth, "Linear models and empirical bayes methods for assessing differential expression in microarray experiments," *Statistical Applications in Genetics and Molecular Biology*, vol. 3, no. 1, pp. 1–25, 2004.
- [9] —, "Limma: linear models for microarray data," in *Bioinformatics and computational biology solutions using R and Bioconductor*. Springer, 2005, pp. 397–420.

CONTACT INFORMATION

- **N.S. Hejazi**, Ph.D. candidate, Group in Biostatistics, NHEJAZI@BERKELEY.EDU
- **M.J. van der Laan**, Professor of Biostatistics & Statistics, LAAN@BERKELEY.EDU
- **A.E. Hubbard**: Professor of Biostatistics, HUBBARD@BERKELEY.EDU