

Leveraging the causal effects of stochastic interventions to evaluate vaccine efficacy in two-phase trials

“Methods for the thorny challenges of real studies”, SER 2020


Nima Hejazi

Tuesday 3rd March, 2020

Graduate Group in Biostatistics, and
Center for Computational Biology,
University of California, Berkeley

 nshejazi

 nhejazi

 nimahejazi.org

Joint work with David Benkeser, Mark van der Laan,
Holly Janes, Peter Gilbert



The burden of HIV-1

- The HIV-1 epidemic — the facts:
 - now in its fourth decade,
 - 2.5 million new infections occurring annually worldwide,
 - new infections outpace patients starting antiretroviral therapy.
- *Most efficacious* preventive vaccine: 31% reduction rate.
- **Question:** To what extent can HIV-1 vaccines be improved by modulating immunogenic CD4⁺/CD8⁺ response profiles?

HVTN 505 trial examined new antibody boost vaccines

- HIV Vaccine Trials Network's (HVTN) 505 vaccine efficacy; randomized controlled trial, $n = 2504$ (Hammer et al. 2013).
- **Question:** How would HIV-1 infection risk in week 28 have changed had immunogenic response (due to vaccine) differed?
- Immunogenic response profiles only available for second-stage sample of $n = 189$ (Janes et al. 2017) due to cost limitations.
- Two-phased sampling mechanism: 100% inclusion rate if HIV-1 positive in week 28; based on matching otherwise.

Two-phase sampling censors the complete data structure

- Complete (unobserved) data $X = (L, A, Y) \sim P_0^X \in \mathcal{M}^X$, as per the full HVTN 505 trial cohort (Hammer et al. 2013):
 - L (baseline covariates): sex, age, BMI, behavioral HIV risk,
 - A (exposure): immune response profile for CD4+ and CD8+,
 - Y (outcome of interest): HIV-1 infection status at week 28.
- Observed data $O = (C, CX) = (L, C, CA, Y)$; $C \in \{0, 1\}$ is an indicator for inclusion in the second-stage sample.

NPSEM with static interventions

- Use a nonparametric structural equation model (NPSEM) to describe the generation of X (Pearl 2009), specifically

$$L = f_L(U_L); A = f_A(L, U_A); Y = f_Y(A, L, U_Y)$$

- Implies a model for the distribution of counterfactual random variables generated by interventions on the process.
- A *static intervention* replaces f_A with a specific value a in its conditional support $A \mid L$.
- This requires specifying a particular value of the exposure under which to evaluate the outcome *a priori*.

NPSEM with stochastic interventions

- *Stochastic interventions* modify the value A would naturally assume by drawing from a modified exposure distribution.
- Consider the post-intervention value $A^* \sim G^*(\cdot \mid L)$; static interventions are a special case (degenerate distribution).
- Such an intervention generates a counterfactual random variable $Y_{G^*} := f_Y(A^*, L, U_Y)$, with distribution P_0^δ .
- We aim to estimate $\psi_{0,\delta} := \mathbb{E}_{P_0^\delta}\{Y_{G^*}\}$, the counterfactual mean under the post-intervention exposure distribution G^* .

Stochastic interventions for the causal effects of shifts

- Díaz and van der Laan (2012; 2018)'s *stochastic* interventions

$$d(a, l) = \begin{cases} a + \delta, & a + \delta < u(l) \quad (\text{if plausible}) \\ a, & a + \delta \geq u(l) \quad (\text{otherwise}) \end{cases}$$

- Our estimand is $\psi_{0,d} := \mathbb{E}_{P_0^d}\{Y_{d(A,L)}\}$, mean of $Y_{d(A,L)}$.
- Statistical target parameter is $\Psi(P_0^X) = \mathbb{E}_{P_0^X}\overline{Q}(d(A, L), L)$, counterfactual mean of the *shifted* outcome mechanism.
- For HVTN 505, $\psi_{0,d}$ is the counterfactual risk of HIV-1 infection, had the observed value of the immune response been altered under the rule $d(A, L)$ defining $G^*(\cdot \mid L)$.

Targeted minimum loss estimator

- TML estimation algorithm updates initial estimators g_n and \bar{Q}_n so as to satisfy an arbitrary set of estimating equations.
- TML estimator built by updating initial estimates of outcome mechanism \bar{Q}_n via a (logistic) tilting model, yielding

$$\psi_n := \psi(P_n^*) = \frac{1}{n} \sum_{i=1}^n \bar{Q}_n^*(d(A_i, L_i), L_i).$$

- In this case, the efficient influence function (EIF) is

$$D(P_0^X)(x) = H(a, l)(y - \bar{Q}(a, l)) + \bar{Q}(d(a, l), l) - \psi(P_0^X)$$

- The auxiliary covariate $H(a, l)$ may be expressed

$$H(a, l) = \mathbb{I}(a + \delta < u(l)) \frac{g_0(a - \delta | l)}{g_0(a | l)} + \mathbb{I}(a + \delta \geq u(l))$$

Augmented estimators for two-phase sampling designs

- Rose and van der Laan (2011) introduce the IPCW-TMLE, to be used when observed data is subject to two-phase sampling.
- *Initial proposal*: correct for two-phase sampling by using a loss function with inverse probability of censoring weights:

$$\mathcal{L}(P_0^X)(O) = \frac{C}{\pi_0(Y, L)} \mathcal{L}^F(P_0^X)(X)$$

- When the sampling mechanism $\pi_0(Y, L)$ can be estimated by a parametric form, this procedure yields an efficient estimator.
- However, when machine learning is used, this is insufficient.

Efficient estimation under two-phase sampling

- When the sampling mechanism is not *known by design*, it is best to employ a flexible (ML) estimator of $\pi_0(Y, L)$.
- Then, the IPCW augmentation must be applied to the EIF:

$$D(P_0^X)(o) = \frac{c}{\pi_0(y, l)} D^F(P_0^X)(x) - \left(1 - \frac{c}{\pi_0(y, l)}\right) \cdot \mathbb{E}(D^F(P_0^X)(x) \mid C = 1, Y = y, L = l),$$

- Expresses observed data EIF $D^F(P_0^X)(o)$ in terms of full data EIF $D^F(P_0^X)(x)$; inclusion of second term induces efficiency.
- The expectation of the full data EIF $D^F(P_0^X)(x)$, taken only over units selected by the sampling mechanism (i.e., $C = 1$).

Emergent property: multiple robustness

- A unique multiple robustness property — combinations of $(g_0(L), \overline{Q}_0(A, L)) \times (\pi_0(Y, L), \mathbb{E}(D^F(P_0^X)(x) \mid C = 1, Y, L))$.
- We now have a semiparametric-efficient and robust procedure for assessing the effect of the intervention $d(a, l) = a + \delta$.
- Due to the construction of the IPCW-TMLE, the resultant estimator is robust and efficient under two-phase sampling.
- This allows us to assess how posited shifts in the assayed immune responses would have affected HIV-1 infection risk.

Finding the best efficient estimator




Figure 1: Relative performance of reweighted and augmented estimators.

Helping to fight the HIV-1 epidemic





Figure 2: Analysis of HIV-1 risk as a function of CD8+ immunogenicity, using R package txshift (<https://github.com/nhejazi/txshift>.)

Big picture takeaways

- Vaccine efficacy evaluation helps to develop enhanced vaccines better informed by biological properties of the target disease.
- HIV-1 vaccines modulate immunogenic response profiles as part of their mechanism for lowering HIV-1 infection risk.
- *Stochastic* interventions constitute a flexible framework for considering **realistic** treatment/intervention policies.
- Large-scale (vaccine) trials often use two-phase designs — need to (carefully!) accommodate for sampling complications.
- We've developed robust, open source statistical software for assessing stochastic interventions in observational studies.

Thank you

 <https://nimahejazi.org>

 <https://github.com/nhejazi>

 <https://twitter.com/nshejazi>

Appendix

From the causal to the statistical target parameter

Assumption 1: *Consistency*

$Y_i^{d(a_i, l_i)} = Y_i$ in the event $A_i = d(a_i, l_i)$, for $i = 1, \dots, n$

Assumption 2: *SUTVA*

$Y_i^{d(a_i, l_i)}$ does not depend on $d(a_j, l_j)$ for $i = 1, \dots, n$ and $j \neq i$, or lack of interference (Rubin 1978; 1980)

Assumption 3: *Strong ignorability*

$A_i \perp\!\!\!\perp Y_i^{d(a_i, l_i)} \mid L_i$, for $i = 1, \dots, n$

From the causal to the statistical target parameter

Assumption 4: *Positivity (or overlap)*

$a_i \in \mathcal{A} \implies d(a_i, l_i) \in \mathcal{A}$ for all $l \in \mathcal{L}$, where \mathcal{A} denotes the support of A conditional on $L = l_i$ for all $i = 1, \dots, n$

- This positivity assumption is not quite the same as that required for categorical interventions.
- In particular, we do not require that the intervention density place mass across all strata defined by L .
- Rather, we merely require the post-intervention quantity be seen in the observed data for given $a_i \in \mathcal{A}$ and $l_i \in \mathcal{L}$.

Literature: Díaz and van der Laan (2012)

- *Proposal*: Evaluate outcome under an altered *intervention distribution* — e.g., $P_\delta(g_0)(A = a \mid L) = g_0(a - \delta(L) \mid L)$.
- Identification conditions for a statistical parameter of the counterfactual outcome $\psi_{0,d}$ under such an intervention.
- Show that the causal quantity of interest $\mathbb{E}_0\{Y_{d(A,L)}\}$ is identified by a functional of the distribution of X :

$$\psi_{0,d} = \int_{\mathcal{L}} \int_{\mathcal{A}} \mathbb{E}_{P_0^X}\{Y \mid A = d(a, l), L = l\} \cdot q_{0,A}^X(a \mid L = l) \cdot q_{0,L}^X(l) d\mu(a) d\nu(l).$$

- Provides a derivation based on the efficient influence function (EIF) with respect to the nonparametric model \mathcal{M} .

Literature: Haneuse and Rotnitzky (2013)

- *Proposal*: Characterization of stochastic interventions as *modified treatment policies* (MTPs).
- Assumption of *piecewise smooth invertibility* allows for the intervention distribution of any MTP to be recovered:

$$g_{0,\delta}(a \mid l) = \sum_{j=1}^{J(l)} l_{\delta,j} \{h_j(a, l), l\} g_0 \{h_j(a, l) \mid l\} h'_j(a, l)$$

- Such intervention policies account for the natural value of the intervention A directly yet are interpretable as the imposition of an altered intervention mechanism.
- Identification conditions for assessing the parameter of interest under such interventions appear technically complex (at first).

Literature: Young et al. (2014)

- Establishes equivalence between g-formula when proposed intervention depends on natural value and when it does not.
- This equivalence leads to a sufficient positivity condition for estimating the counterfactual mean under MTPs via the same statistical functional studied in Díaz and van der Laan (2012).
- Extends earlier identification results, providing a way to use the same statistical functional to assess $\mathbb{E}Y_{d(A,L)}$ or $\mathbb{E}Y_{d(L)}$.
- The authors also consider limits on implementing shifts $d(A, L)$, and address working in a longitudinal setting.

Literature: Díaz and van der Laan (2018)

- Builds on the original proposal, accomodating MTP-type shifts $d(A, L)$ proposed after their earlier work.
- To protect against positivity violations, considers a specific shifting mechanism:

$$d(a, l) = \begin{cases} a + \delta, & a + \delta < u(l) \\ a, & \text{otherwise} \end{cases}$$

- Proposes an improved “1-TMLE” algorithm, with a single auxiliary covariate for constructing the TML estimator.
- Our (first) contribution: implementation of this algorithm.

Nonparametric conditional density estimation

- To compute the auxiliary covariate $H(a, l)$, we need to estimate conditional densities $g(A \mid L)$ and $g(A - \delta \mid L)$.
- There is a rich literature on density estimation, we follow the approach proposed in Díaz and van der Laan (2011).

- To build a conditional density estimator, consider

$$g_{n,\alpha}(a \mid L) = \frac{\mathbb{P}(A \in [\alpha_{t-1}, \alpha_t) \mid L)}{\alpha_t - \alpha_{t-1}},$$

for $\alpha_{t-1} \leq a < \alpha_t$.

- This is a classification problem, where we estimate the probability that a value of A falls in a bin $[\alpha_{t-1}, \alpha_t)$.
- The choice of the tuning parameter t corresponds roughly to the choice of bandwidth in classical kernel density estimation.

Nonparametric conditional density estimation

- Díaz and van der Laan (2011) propose a re-formulation of this classification approach as a set of hazard regressions.
- To effectively employ this proposed re-formulation, consider

$$\mathbb{P}(A \in [\alpha_{t-1}, \alpha_t) \mid L) = \mathbb{P}(A \in [\alpha_{t-1}, \alpha_t) \mid A \geq \alpha_{t-1}, L) \times \prod_{j=1}^{t-1} \{1 - \mathbb{P}(A \in [\alpha_{j-1}, \alpha_j) \mid A \geq \alpha_{j-1}, L)\}$$

- The likelihood of this model may be expressed to correspond to the likelihood of a binary variable in a data set expressed via a long-form repeated measures structure.
- Specifically, the observation of X_i is repeated as many times as intervals $[\alpha_{t-1}, \alpha_t)$ are before the interval to which A_i belongs, and the binary variables indicating $A_i \in [\alpha_{t-1}, \alpha_t)$ are recorded.

Density estimation with the Super Learner algorithm

- To estimate $g(A | L)$ and $g(A - \delta | L)$, use a pooled hazard regression, spanning the support of A .
- We rely on the Super Learner algorithm of van der Laan et al. (2007) to build an ensemble learner that optimally weights each of the proposed regressions, using cross-validation (CV).
- The Super Learner algorithm uses V -fold CV to train each proposed regression model, weighting each by the inverse of its average risk across all V holdout sets.
- By using a library of regression estimators, we invoke the result of van der Laan et al. (2004), who prove this likelihood-based cross-validated estimator to be asymptotically optimal.

Key properties of TML estimators

- **Asymptotic linearity:**

$$\Psi(P_n^\star) - \Psi(P_0^X) = \frac{1}{n} \sum_{i=1}^n D(P_0^X)(X_i) + o_P\left(\frac{1}{\sqrt{n}}\right)$$

- **Gaussian limiting distribution:**

$$\sqrt{n}(\Psi(P_n^\star) - \Psi(P_0^X)) \rightarrow N(0, \text{Var}(D(P_0^X)(X)))$$

- **Statistical inference:**

$$\text{Wald-type confidence interval : } \Psi(P_n^\star) \pm z_{1-\frac{\alpha}{2}} \cdot \frac{\sigma_n}{\sqrt{n}},$$

where σ_n^2 is computed directly via $\sigma_n^2 = \frac{1}{n} \sum_{i=1}^n D^2(\cdot)(X_i)$.

Algorithm for TML estimation

1. Construct initial estimators g_n of $g_0(A, L)$ and Q_n of $\bar{Q}_0(A, L)$, perhaps using data-adaptive regression techniques.
2. For each observation i , compute an estimate $H_n(a_i, l_i)$ of the auxiliary covariate $H(a_i, l_i)$.

3. Estimate the parameter ϵ in the logistic regression model

$$\text{logit} \bar{Q}_{\epsilon, n}(a, l) = \text{logit} \bar{Q}_n(a, l) + \epsilon H_n(a, l),$$

or an alternative regression model incorporating weights.

4. Compute TML estimator Ψ_n of the target parameter, defining update \bar{Q}_n^* of the initial estimate \bar{Q}_{n, ϵ_n} :

$$\Psi_n = \Psi(P_n^*) = \frac{1}{n} \sum_{i=1}^n \bar{Q}_n^*(d(A_i, L_i), L_i).$$

Algorithm for IPCW-TML estimation

1. Using all observed units (X), estimate sampling mechanism $\pi(Y, L)$, perhaps using data-adaptive regression methods.
2. Using only observed units in the second-stage sample $\Delta = 1$, construct initial estimators $g_n(A, L)$ and $\bar{Q}_n(A, L)$, weighting by the sampling mechanism estimate $\pi_n(Y, L)$.
3. With the approach described for the full data case, compute $H_n(a_i, l_i)$, and fluctuate submodel via logistic regression.
4. Compute IPCW-TML estimator Ψ_n of the target parameter, by solving the IPCW-augmented EIF estimating equation.
5. Iteratively update estimated sampling weights $\pi_n(Y, L)$ and IPCW-augmented EIF, updating TML estimate in each iteration, until $\frac{1}{n} \sum_{i=1}^n \text{EIF}_i < \frac{1}{n}$.

A linear modeling perspective

- Briefly consider a simple data structure: $X = (Y, A)$; we seek to model the outcome Y as a function of A .
- To posit a linear model, consider $Y_i = \beta_0 + \beta_1 A_i + \epsilon_i$, with error $\epsilon_i \sim N(0, 1)$.
- Letting δ be a change in A , $Y_{A+\delta} - Y_A$ may be expressed

$$\begin{aligned}\mathbb{E}Y_{A+\delta} - \mathbb{E}Y_A &= [\beta_0 + \beta_1(\mathbb{E}A + \delta)] - [\beta_0 + \beta_1(\mathbb{E}A)] \\ &= \beta_0 - \beta_0 + \beta_1\mathbb{E}A - \beta_1\mathbb{E}A + \beta_1\delta \\ &= \beta_1\delta\end{aligned}$$

- Thus, a *unit shift* in A (i.e., $\delta = 1$) may be seen as inducing a change in the difference in outcomes of magnitude β_1 .

A causal inference perspective

- Consider a data structure: $(Y_a, a \in \mathcal{A})$.
- To posit a linear model, let $Y_a = \beta_0 + \beta_1 a + \epsilon_a$ for $a \in \mathcal{A}$, with error $\epsilon_a \sim N(0, \sigma_a^2) \forall a \in \mathcal{A}$.
- For the counterfactual outcomes $(Y_{a'+\delta}, Y_{a'})$, their difference, $Y_{a'+\delta} - Y_{a'}$, for some $a' \in \mathcal{A}$, may be expressed

$$\begin{aligned}\mathbb{E}Y_{a'+\delta} - \mathbb{E}Y_{a'} &= [\beta_0 + \beta_1(a' + \delta) + \mathbb{E}\epsilon_{a'+\delta}] - [\beta_0 + \beta_1 a' + \mathbb{E}\epsilon_{a'}] \\ &= \beta_1 \delta\end{aligned}$$

- Thus, a *unit shift* for $a' \in \mathcal{A}$ (i.e., $\delta = 1$) may be seen as inducing a change in the difference in the counterfactual outcomes of magnitude β_1 .

Slope in a semiparametric model

- Consider the stochastic intervention $g^*(\cdot | L)$:

$$\begin{aligned}\mathbb{E}Y_{g^*} &= \int_L \int_a \mathbb{E}(Y | A = a, L) g(a - \delta | L) \cdot da \cdot dP_0(L) \\ &= \int_L \int_z \mathbb{E}(Y | A = z + \delta, L) g(z | L) \cdot dz \cdot dP_0(L),\end{aligned}$$

defining the change of variable $z = a - \delta$.

- For a semiparametric model, $\mathbb{E}(Y | A = z, L) = \beta z + \theta(L)$:

$$\begin{aligned}\mathbb{E}Y_{g^*} - \mathbb{E}Y &= \int_L \int_z [\mathbb{E}(Y | A = z + \delta, L) - \mathbb{E}(Y | A = z, L)] \\ &\quad g(z | L) \cdot dz \cdot dP_0(L) \\ &= [\beta(z + \delta) + \theta(L)] - [\beta z + \theta(L)] \\ &= \beta \delta\end{aligned}$$

References

- Breiman, L. (1996). Stacked regressions. *Machine Learning*, 24(1):49–64.
- Díaz, I. and van der Laan, M. J. (2011). Super learner based conditional density estimation with application to marginal structural models. *The international journal of biostatistics*, 7(1):1–20.
- Díaz, I. and van der Laan, M. J. (2012). Population intervention causal effects based on stochastic interventions. *Biometrics*, 68(2):541–549.
- Díaz, I. and van der Laan, M. J. (2018). Stochastic treatment regimes. In *Targeted Learning in Data Science: Causal Inference for Complex Longitudinal Studies*, pages 167–180. Springer Science & Business Media.

- Dudoit, S. and van der Laan, M. J. (2005). Asymptotics of cross-validated risk estimation in estimator selection and performance assessment. *Statistical Methodology*, 2(2):131–154.
- Hammer, S. M., Sobieszczyk, M. E., Janes, H., Karuna, S. T., Mulligan, M. J., Grove, D., Koblin, B. A., Buchbinder, S. P., Keefer, M. C., Tomaras, G. D., et al. (2013). Efficacy trial of a DNA/rAd5 HIV-1 preventive vaccine. *New England Journal of Medicine*, 369(22):2083–2092.
- Haneuse, S. and Rotnitzky, A. (2013). Estimation of the effect of interventions that modify the received treatment. *Statistics in medicine*, 32(30):5260–5277.
- Holland, P. W. (1986). Statistics and causal inference. *Journal of the American statistical Association*, 81(396):945–960.

- Janes, H. E., Cohen, K. W., Frahm, N., De Rosa, S. C., Sanchez, B., Hural, J., Magaret, C. A., Karuna, S., Bentley, C., Gottardo, R., et al. (2017). Higher t-cell responses induced by dna/rad5 hiv-1 preventive vaccine are associated with lower hiv-1 infection risk in an efficacy trial. *The Journal of infectious diseases*, 215(9):1376–1385.
- Kennedy, E. H. (2018). Nonparametric causal effects based on incremental propensity score interventions. *Journal of the American Statistical Association*, (just-accepted).
- Kennedy, E. H., Ma, Z., McHugh, M. D., and Small, D. S. (2017). Non-parametric methods for doubly robust estimation of continuous treatment effects. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 79(4):1229–1245.
- Pearl, J. (2009). *Causality: Models, Reasoning, and Inference*. Cambridge University Press.

- Rose, S. and van der Laan, M. J. (2011). A targeted maximum likelihood estimator for two-stage designs. *The International Journal of Biostatistics*, 7(1):1–21.
- Rosenbaum, P. R. and Rubin, D. B. (1983). The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70(1):41–55.
- Rubin, D. B. (1978). Bayesian inference for causal effects: The role of randomization. *The Annals of statistics*, pages 34–58.
- Rubin, D. B. (1980). Randomization analysis of experimental data: The fisher randomization test comment. *Journal of the American Statistical Association*, 75(371):591–593.
- Rubin, D. B. (2005). Causal inference using potential outcomes: Design, modeling, decisions. *Journal of the American Statistical Association*, 100(469):322–331.

- van der Laan, M. J., Dudoit, S., and Keles, S. (2004). Asymptotic optimality of likelihood-based cross-validation. *Statistical Applications in Genetics and Molecular Biology*, 3(1):1–23.
- van der Laan, M. J., Polley, E. C., and Hubbard, A. E. (2007). Super Learner. *Statistical Applications in Genetics and Molecular Biology*, 6(1).
- van der Laan, M. J. and Rubin, D. (2006). Targeted maximum likelihood learning. *The International Journal of Biostatistics*, 2(1).
- Wolpert, D. H. (1992). Stacked generalization. *Neural networks*, 5(2):241–259.
- Young, J. G., Hernán, M. A., and Robins, J. M. (2014). Identification, estimation and approximation of risk under interventions that depend on the natural value of treatment using observational data. *Epidemiologic methods*, 3(1):1–19.