

The Role of Endodontics After Dental Traumatic Injuries

BILL KAHLER*

CHAPTER OUTLINE

Unique Aspects of Dental Trauma

Most Common Types of Dental Trauma

Crown Fractures

Crown-Root Fractures

Root Fractures

Luxation Injuries and Avulsion

Follow-Up After Dental Trauma

Radiographic Examinations

Cone-Beam Computed Tomography and Dentoalveolar Trauma

Root Resorption

Horizontal (Transverse) Root Fractures

Luxation Injuries

Crown Fractures

Crown Infraction

Biologic Consequences

Uncomplicated Crown Fracture

Incidence

Biologic Consequences

Treatment

Complicated Crown Fracture

Incidence

Biologic Consequences

Treatment

Stage of Development of the Tooth

Time Between Trauma and Treatment

Concomitant Attachment Damage

Restorative Treatment Plan

Vital Pulp Therapy: Requirements for Success

Treatment Methods

Pulp Capping

Partial Pulpotomy

Full Pulpotomy

Pulpectomy

Treatment of the Nonvital Pulp

Immature Tooth: Apexification

Mature Tooth

Crown-Root Fracture

Root Fracture

Diagnosis and Clinical Presentation

Treatment

Healing Patterns

Treatment of Complications

Coronal Root Fractures

Midroot and Apical Root Fractures

Follow-Up

Prognosis

Luxation Injuries

Definitions

Incidence

Treatment

Biologic Consequences

External Root Resorption

Internal Root Resorption

Diagnostic Features of External Versus Internal Root Resorption

Summary of Possible Diagnostic Features

Clinical Management of the Avulsed Tooth

Consequences of Tooth Avulsion

Treatment Objectives

Clinical Management

Emergency Treatment at the Accident Site

Management in the Dental Office

Diagnosis and Treatment Planning

Preparation of the Root

Extraoral Dry Time Less Than 60 Minutes

Closed Apex

Open Apex

Extraoral Dry Time More Than 60 Minutes

Closed Apex

Open Apex

Preparation of the Socket

Splinting

Management of the Soft Tissues

Adjunctive Therapy

Second Visit

Endodontic Treatment

Extraoral Time Less Than 60 Minutes

Closed Apex

Open Apex

*The author acknowledges the outstanding work of Drs. Trope, Barnett, Sigurdsson, and Chivian in previous editions of this text. The present chapter is built on their foundational work.

Extraoral Time More Than 60 Minutes

Closed Apex
Open Apex (If Replanted)

Temporary Restoration

Root Filling Visit

Permanent Restoration**Follow-Up Care**

Late Complications
Dentoalveolar Trauma to the Primary
Dentition

A traumatic injury to the tooth results in damage to many dental and periradicular structures, making the management and consequences of these injuries multifactorial. Knowledge of the interrelating healing patterns of these tissues is essential. This chapter concentrates on the role of the dentinopulpal complex in the pathogenesis of disease subsequent to dentinal trauma and how treatment of this complex can contribute to favorable healing after an injury.

Unique Aspects of Dental Trauma

Most dental trauma occurs in the 7- to 12-year-old age group and is mainly due to falls and accidents near home or school.^{28,165} It occurs primarily in the anterior region of the mouth, affecting the maxillary more than the mandibular jaw.³³ Serious accidents, such as automobile crashes, can affect any tooth and can occur in all age ranges. In many cases, after a traumatic dental injury, endodontic treatment is provided to caries-free, single-rooted, young permanent teeth. If quick and correct treatment for these teeth is provided after injury, the potential for a successful endodontic outcome is very good.

Most Common Types of Dental Trauma

CROWN FRACTURES

Most crown fractures occur in young, caries-free anterior teeth.^{109,128} This makes maintaining or regaining pulp vitality essential. Luckily, vital pulp therapy supports a good prognosis in these situations if correct treatment and follow-up procedures are carefully followed.

CROWN-ROOT FRACTURES

Crown-root fractures are first treated periodontally to ensure that there is a sufficient and good margin to allow restoration. If the tooth can be maintained from a periodontal point of view, then the pulp is treated as for a crown fracture.

ROOT FRACTURES

A surprisingly large number of pulps in root-fractured teeth will survive this rather dramatic injury. In almost every case, the apical segment stays vital, and in many cases, the coronal segment stays vital or regains vitality subsequent to the injury. If the coronal segment permanently loses vitality, it should be treated as an immature permanent tooth with nonvital pulp. *The apical segment rarely needs treatment.*

LUXATION INJURIES AND AVULSION

Luxation injuries and avulsion often result in pulp necrosis and damage to the cemental protective layer of the root. The

potential complication of pulp infection in a root that has lost its cemental protective layer makes these injuries potentially catastrophic. Correct emergency and follow-up evaluation, which may include timely endodontic treatment, is critical.

Follow-Up After Dental Trauma

The reader is referred to Chapter 1 for specific descriptions of pulp tests, but a few general statements about pulp tests on traumatized teeth may be helpful in trying to interpret the results.

For decades, controversy has surrounded the validity of thermal and electric tests on traumatized teeth. Only generalized impressions may be gained from these tests following a traumatic injury. They are in reality sensitivity tests for nerve function and do not indicate the presence or absence of blood circulation within the pulpal space. It is assumed that subsequent to traumatic injury, the conduction capability of the nerve endings and/or sensory receptors is sufficiently deranged to inhibit the nerve impulse from an electrical or thermal stimulus. This makes the traumatized tooth vulnerable to false-negative readings from such tests.¹⁴⁶

Teeth that give a response at the initial examination cannot be assumed to be healthy and to continue to give a response over time. Teeth that yield no response cannot be assumed to have necrotic pulps because they may give a response at later follow-up visits. It has been demonstrated that it may take as long as 9 months for normal blood flow to return to the coronal pulp of a traumatized fully formed tooth. As circulation is restored, the responsiveness to pulp tests returns.⁸¹

The transition from a negative response to a positive response at a subsequent test may be considered a sign of a healing pulp. The repetitious finding of responses may be taken as a sign of a healthy pulp. The transition from a response to no response may be taken as an indication that the pulp is probably undergoing degeneration, and therefore an intervention in the form of endodontic treatment might be indicated. The persistence of no response would suggest that the pulp has been irreversibly damaged, but even this is not absolute.³⁹

Thermal and electrical pulp tests of all anterior teeth (canine to canine) of the maxillary and mandibular jaws should be performed at the time of the initial examination and carefully recorded to establish a baseline for comparison with subsequent repeated tests in later months. These tests should be repeated at 2 to 4 and 6 to 8 weeks; at 3, 6, and 12 months; and at yearly intervals after the trauma (according to the guidelines of the American Association of Endodontists). The purpose of the tests is to establish a trend as to the physiologic status of the pulps of these teeth. Particularly in traumatized teeth, carbon dioxide snow (CO₂, -78°C) or dichlorodifluoromethane (-40°C) placed on the incisal third of the facial surface gives more accurate responses than a water-ice pencil (Fig. 21.1).^{79,80} The intense cold seems to penetrate the tooth and covering splints or restorations and



Fig. 21.1 Dichlorodifluoromethane (-40°F [-40°C]) gas is sprayed on a cotton pellet and then placed on the incisal edge of the maxillary incisor.

reach the deeper areas of the tooth. Neither the dry ice nor the dichlorodifluoromethane spray forms ice water, which could disperse over adjacent teeth or gingiva to give a false-positive response. In trauma evaluation, there is no question that using water-ice pencils should be avoided because of this. Dichlorodifluoromethane spray is a very inexpensive alternative to CO_2 snow; its coldness elicits much more reliable responses than water-ice. The electrical pulp test relies on electrical impulses directly stimulating the nerves of the pulp. These tests have limited value in young teeth but are useful when the dentinal tubules are closed and do not allow dentinal fluid to flow in them. This situation is typical of teeth in elderly patients or in traumatized teeth that are undergoing premature sclerosis. In these situations, the thermal tests that rely on fluid flow in the tubules cannot be used, and the electrical pulp test becomes important.

Laser Doppler flowmetry (LDF) was introduced in the early 1970s for measurement of blood flow in the retina.¹⁰² The technique has also been used to assess blood flow in other tissue systems, such as the skin and renal cortex. It uses a beam of infrared (780 to 820 nm) or near-infrared (632.8 nm) light that is directed into the tissue by optical fibers. As light enters the tissue, it is scattered by moving red blood cells and stationary tissue cells. Photons that interact with moving red blood cells are scattered, and the frequency shifts according to the Doppler principle. Photons that interact with stationary tissue cells are scattered but not Doppler shifted. A portion of the light is returned to a photodetector, and a signal is produced (Fig. 21.2).



Fig. 21.2 Laser Doppler machine. (Courtesy Moor Instruments, Devon, United Kingdom.)

Attempts have been made to use LDF technology for pulp vitality diagnosis in traumatized teeth because this would provide a more accurate reading of the vitality status of the pulp.^{136,196,197} Studies have shown promising results, indicating that laser Doppler can detect blood flow more consistently and earlier than the standard vitality tests would be expected to render a response. In a study on young dogs, laser Doppler was able to correctly detect blood flow as early as 2 to 3 weeks after avulsion of an immature tooth and at the same time indicate no flow in those that would remain necrotic.¹⁹⁷

Presently the cost of an LDF machine limits its use in private dental offices; these are used primarily in hospitals and teaching institutions.

Radiographic Examinations

Radiographic imaging is essential for thorough examination, diagnosis, and management of dentoalveolar trauma. Imaging may reveal root fractures, subgingival crown fractures, tooth displacements, bone fractures, root resorptions, and embedded foreign objects. A single radiograph, even a panoramic image, is insufficient to properly diagnose just about any dental trauma case. In its 2020 guidelines for the management of traumatic dental injuries, and in the current recommendations on its interactive website (www.dentaltraumaguide.org), the International Association of Dental Traumatology (IADT) has recommended taking at least four different radiographs for almost every injury^{44a,64,75a}: a direct 90-degree on the axis of the tooth, two with different vertical angulations, and one occlusal film (Fig. 21.3).

Multiple radiographs increase the likelihood of diagnosing root fractures, tooth displacement, and other possible injuries. However, two-dimensional (2D) imaging methods



Fig. 21.3 An occlusal film of a luxation injury of central incisors. Both were diagnosed to be laterally luxated with apical translocation. Note that the left central is completely obliterated and has a history of being luxated some years prior.

have well-known inherent limitations, and their lack of three-dimensional (3D) information may prevent proper diagnosis and adversely affect long-term treatment outcomes. The interpretation of an image can be confounded by a number of factors, including the regional anatomy and superimposition of both the teeth and surrounding dentoalveolar structures. As a result of superimposition, periapical radiographs reveal only limited aspects (i.e., a 2D view) of the true 3D anatomy.^{52,142} In addition, there is often geometric distortion of the anatomic structures imaged with conventional radiographic methods.⁴ These problems can be overcome by using cone-beam computed tomography (CBCT) imaging techniques, which produce accurate 3D images of the teeth and surrounding dentoalveolar structures and have shown promise in improving the clinician's ability to properly diagnose luxation injuries, alveolar fractures, root fractures, and root resorption (Fig. 21.4).^{50,51}

In instances of soft-tissue laceration, it is advisable to radiograph the injured area before suturing to be sure that no foreign objects have been embedded. A soft-tissue radiograph with a normal-sized film briefly exposed at reduced kilovoltage should reveal the presence of many foreign substances, including tooth fragments (Fig. 21.5).

CONE-BEAM COMPUTED TOMOGRAPHY AND DENTOALVEOLAR TRAUMA

CBCT is accomplished by using a rotating gantry to which an x-ray source and detector are fixed (see Chapter 2). A divergent pyramidal or cone-shaped source of ionizing radiation is directed through the middle of the area of interest onto an area x-ray detector on the opposite side of the patient. The x-ray source and detector rotate around a fixed fulcrum within the region of interest. During the exposure sequence, hundreds of planar projection images are acquired of the field of view (FOV) in an arc of at least 180 degrees. In this single rotation, CBCT provides precise, essentially immediate, and accurate 3D radiographic images. Because CBCT exposure incorporates the entire FOV, only one rotational sequence of the gantry is necessary to acquire enough data for image reconstruction.

CBCT has been suggested as an adjunct imaging tool when the true nature of the dentoalveolar root fracture and dental injuries cannot be confidently diagnosed from a conventional examination and radiographs.^{13,50,134,143} However, it has been repeatedly demonstrated that the use of CBCT provides improved diagnostic images in cases of




Fig. 21.4 **A**, Panoramic radiograph taken on a patient with past history of dentoalveolar trauma. Tooth #9 appeared to have arrested root development with a periapical radiolucency. **B**, Sagittal view of tooth #9 of same patient from CBCT imaging. This revealed that tooth #9 had arrested root development, a large periapical radiolucency, and extensive root resorption along the palatal surface that was *not* evident on the panoramic image. **C**, Periapical radiograph of tooth #9 with extensive complicated crown fracture. The clinical examination revealed the crown fracture extended subgingivally on the palatal aspect. **D**, Sagittal view of tooth #9 of same patient from CBCT imaging. This revealed that the apical extent of the fractured palatal portion extended just apical to the crest of bone. The correct diagnosis of this injury was complicated crown-root fracture. **E**, Periapical radiograph of teeth #8, #9 revealed crown fractures. **F**, Sagittal view of tooth #9 of same patient from CBCT imaging. This revealed that tooth #9 suffered a lateral luxation displacement injury with extensive concomitant alveolar fracture. The necessary treatment of repositioning and splinting was apparent.



Fig. 21.4, cont'd **G**, Periapical radiograph of tooth #9 following dentoalveolar trauma. Tooth #8 was avulsed and tooth #9 appeared to be severely intruded. **H**, Sagittal view of tooth #9 of same patient from CBCT imaging. This revealed the labial position of tooth #9. **I**, Periapical radiograph of tooth #8 revealed a lateral luxation injury. **J**, Sagittal view of tooth #8 from CBCT imaging. The extensive tooth displacement and concomitant alveolar are evident. **K**, Periapical radiograph of teeth #8 to #10, that had recently been splinted in an emergency room. This image revealed that these teeth were not properly repositioned before splinting, evidenced by the spaces along each root. **L**, Axial view of maxillary anterior region of same patient from CBCT imaging. This revealed that teeth #8 to #10 are labially displaced. **M**, Sagittal view of tooth #10 of same patient revealed how severe the displacement was. Following consent and local anesthesia, the splint was removed and teeth #8 to #10 were repositioned and resplinted.

dentoalveolar injury.^{37,38,50,51,52,66,71,123,134,142,143} Perhaps the use of CBCT imaging for dentoalveolar injuries could be considered “best practice” and is recommended for most serious injuries (Fig. 21.6).⁶⁹

ROOT RESORPTION

Because the development and progression of root resorption occur without clinical signs or symptoms, their early detection is challenging. Therefore the definitive diagnosis of root resorption depends on its radiographic demonstration, which in turn is limited by the diagnostic accuracy of the imaging device used to determine its presence (see Chapter 18).¹⁴³ The radiodensity of the root requires that a significant amount of root substance be removed to cause enough contrast on the radiograph to allow it to be

detected. Thus only resorptive defects on the mesial or distal aspects of the root can be predictably detected after some time; the facial and palatal or lingual aspects are much harder to see. To overcome these difficulties, it is essential to take as many different horizontal-angled radiographs as feasible in cases of suspected root resorption. Early detection of small resorption defects has been shown to be poor with conventional dental radiographs, and the extent of the resorptive defect is grossly underestimated compared with CBCT.^{38,66,71}

The available literature supports the use of CBCT as a diagnostic tool to assess the true nature of teeth diagnosed with root resorption to improve diagnosis and aid management. This should ultimately improve the prognosis of teeth with root resorption that require endodontic management (see Fig. 21.4, A, B).¹⁴³




Fig. 21.5 Patient sustained complicated crown fractures on the left lower incisors and later on the canine tooth. **A**, An emergency pulp capping; Band-Aid composite was placed on the teeth. Six days later the patient was referred to the clinic because of a poorly healed laceration on the lip. **B**, Radiograph of the lip revealed a portion of the lateral crown still in the lip. The patient was anesthetized and the crown removed. **C**, Healing was quick and uneventful.




Fig. 21.6 **A1**, A clinical photograph of a maxillary right central incisor that sustained a crown-root fracture. **A2**, A periapical radiograph that failed to show the extent of the crown-root fracture. **A3**, A CBCT image revealed the extent of the crown-root fracture. **B1**, A clinical photograph of a right maxillary central incisor that had sustained a crown fracture that was restored with composite resin by the referring dentist. **B2**, A periapical radiograph revealed a vertical fracture in the crown and the coronal third of the root. An area of bone loss just below the alveolar crest on the mesial aspect of the root suggested further injury. **B3**, A CBCT image revealed the further injury as a transverse root fracture. (Reproduced from Andreasen FM, Kahler B: Diagnosis of acute dental trauma: the importance of standardized documentation: a review, *Dent Traumatol* 31:340, 2015.)

HORIZONTAL (TRANSVERSE) ROOT FRACTURES

There is a significant risk of misdiagnosing the true location of a root fracture in anterior teeth when intraoral radiography is used because of the possibility of the oblique course of the fracture line in the sagittal plane. It has been shown that horizontal root fracture can be detected sooner using CBCT than with periapical views, and the fracture can be assessed in coronal, axial, and cross-sectional views (see Fig. 21.4, C, D).¹¹⁷ Compared with conventional radiographs, CBCT increased the accuracy of diagnosing the actual nature of horizontal root fractures.^{37,134}

LUXATION INJURIES

As previously mentioned, conventional intraoral radiography provides poor sensitivity in the detection of minimal tooth displacements and root and alveolar fractures.¹²³ CBCT has significantly improved the ability to accurately diagnose traumatic injuries and has the potential to overcome most of the technical limitations of the plain-film projection (see Fig. 21.4, E–M).^{37,38,50,51,66,71,123,134,143}

Crown Fractures

As already mentioned, the primary aim from an endodontic point of view is to maintain pulp vitality after crown fractures.

CROWN INFRACTION

Crown infraction can be defined as an incomplete fracture of or a crack in the enamel, without loss of tooth structure.²¹

Biologic Consequences

Crown infractions are injuries that carry little danger of resulting in pulp necrosis. Meticulous follow-up over a 5-year period is the most important endodontic preventive measure in these cases. If, at any follow-up examination, the reaction to sensitivity tests changes, or if, on radiographic assessment, signs of apical or periradicular periodontitis develop or the root appears to have stopped development or is obliterating, endodontic intervention should be considered.

UNCOMPLICATED CROWN FRACTURE

Uncomplicated crown fracture can be defined as fracture of the enamel only or the enamel and dentin without pulp exposure.²¹

Incidence

Uncomplicated crown fracture is likely to be the most commonly reported dental injury. It is estimated to account for at least one third to one half of all reported dental trauma.

Biologic Consequences

Uncomplicated crown fractures are also injuries that have little danger of resulting in pulp necrosis. In fact, the biggest danger to the health of the pulp is through iatrogenic causes during the esthetic restoration of these teeth.

TREATMENT

There are two key issues in the treatment of crown fractures. First, all exposed dental tubules need to be closed as soon as reasonably possible. If the broken-off piece is not available or if it is not possible to reattach it and there is no time to do a full composite restoration at the time of the emergency appointment, a composite Band-Aid or temporary coverage should be placed on all exposed dentin. This prevents any ingress of bacteria into the tubules and reduces the patient's discomfort. The second issue is the remaining dentin thickness. Several studies have confirmed that if the remaining dentin is more than 0.5 mm thick, the tooth can be restored with the restoration of choice, including etching and bonding, and no special attention needs to be given to the pulp.^{4,54,70,137,173} However, if the remaining dentin is less than this thickness, a protective layer of hard-setting calcium hydroxide in the deepest part of the dentin exposure does reduce, if not completely prevent, reactive inflammation of the underlying pulp, which is a significantly different reaction compared with the negative reaction to composite bonding systems.^{4,54,70,137,173}

COMPLICATED CROWN FRACTURE

A *complicated crown fracture* involves enamel, dentin, and pulp (Fig. 21.7).²¹



Fig. 21.7 Complicated crown fracture involving enamel, dentin, and pulp.

Incidence

Complicated crown fractures occur in 0.9% to 13% of all dental injuries.^{47,153,175}

Biologic Consequences

A crown fracture involving the pulp, if left untreated, always results in pulp necrosis.¹¹⁶ However, the manner and time sequence in which the pulp becomes necrotic allow a great deal of potential for successful intervention to maintain pulp vitality. The first reaction after the injury is hemorrhage and local inflammation (Fig. 21.8).

Subsequent inflammatory changes are usually proliferative but can be destructive over time. A proliferative reaction is favored in traumatic injuries because the fractured surface is usually flat, allowing salivary rinsing with little chance of impaction of contaminated debris. Unless impaction of contaminated debris is obvious, it is expected that in the first 24 hours after the injury, a proliferative response with inflammation extending not more than 2 mm into the pulp will be present (see Fig. 21.8).^{57,60,90} In time, the bacterial challenge results in local pulpal necrosis and a slow apical progression of the pulpal inflammation (Fig. 21.9).

TREATMENT

Treatment options for complicated crown fracture are (1) vital pulp therapy, comprising pulp capping, partial pulpotomy, or full pulpotomy; and (2) pulpectomy. The choice of treatment depends on the stage of development of the tooth, the time between trauma and treatment, concomitant periodontal injury, and the restorative treatment plan.

Stage of Development of the Tooth

Loss of vitality in an immature tooth can have catastrophic consequences. Root canal treatment on a tooth with a blunderbuss canal is time consuming and difficult. It is probably more important that necrosis of an immature tooth leaves it



Fig. 21.8 Histologic appearance of the pulp within 24 hours of a traumatic exposure. The pulp proliferated over the exposed dentinal tubules. There is approximately 1.5 mm of inflamed pulp below the surface of the fracture.




Fig. 21.9 Histologic appearance of the pulp days after a traumatic exposure. Superficial necrosis above a zone of inflamed pulp is seen.

with thin dentinal walls that are susceptible to fracture both during and after the apexification procedure.¹¹⁸ Every effort must be made to keep the tooth vital, at least until the apex and cervical root have completed their development.

Removal of the pulp in a mature tooth is not as significant as in an immature tooth because a pulpectomy in a mature tooth has an extremely high success rate.¹⁶⁴ However, it has been shown that under optimal conditions, vital pulp therapy (rather than removal) can be carried out successfully on a mature tooth.^{133,193} Therefore this form of therapy can be an option under certain circumstances, even though a pulpectomy is the treatment that affords the most predictable success.

In an immature tooth, vital pulp therapy should always be attempted if at all feasible because of the tremendous advantages of maintaining the vital pulp.

Time Between Trauma and Treatment

For 48 hours after a traumatic injury, the initial reaction of the pulp is proliferative, with no more than a 2-mm depth of pulpal inflammation (see Fig. 21.9). After 48 hours, chances of direct bacterial contamination of the pulp increase, with the zone of inflammation progressing apically⁶⁰; as time passes, the likelihood of successfully maintaining a healthy pulp decreases.

Concomitant Attachment Damage

A periodontal injury compromises the nutritional supply of the pulp. This fact is particularly important in mature teeth, in which the chance of pulp survival is not as good as for immature teeth.^{18,68}

Restorative Treatment Plan

Unlike in an immature tooth, in which the benefits of maintaining vitality of the pulp are so great, pulpectomy is

a viable treatment option in a mature tooth. As pointed out, if performed under optimal conditions, vital pulp therapy after traumatic exposures can be successful. If the restorative treatment plan is simple and a composite resin restoration will suffice as the permanent restoration, this treatment option should be given serious consideration. If a more complex restoration is to be placed (e.g., a crown or bridge abutment), pulpectomy may be the more predictable treatment method.

VITAL PULP THERAPY: REQUIREMENTS FOR SUCCESS

Vital pulp therapy has an extremely high success rate if the following requirements can be met (also see Chapter 24).

- *Treatment of a noninflamed pulp.* Treatment of a healthy pulp has been shown to be an important requirement for successful therapy.^{174,183} Vital pulp therapy of the inflamed pulp yields an inferior success rate,^{174,183} so the optimal time for treatment is in the first 24 hours when pulp inflammation is superficial. As time between the injury and therapy increases, pulp removal must be extended apically to ensure that noninflamed pulp has been reached.
- *Bacteria-tight seal.* This requirement can be considered the most critical factor for successful treatment.¹⁷⁴ Challenge by bacteria during the healing phase causes failure (see also Chapter 15).⁵³ If the exposed pulp is effectively sealed from bacterial leakage, successful healing of the pulp with a hard-tissue barrier will occur independent of the dressing placed on the pulp and after more extended time periods between accident and treatment.^{54,88}
- *Pulp dressing.* Calcium hydroxide has traditionally been used for vital pulp therapy. Its main advantage is that it is antibacterial^{46,163} and disinfects the superficial pulp. Pure calcium hydroxide causes necrosis of approximately 1.5 mm of pulp tissue, which removes superficial layers of inflamed pulp if present (Fig. 21.10).¹³⁵ The

high pH (12.5) of calcium hydroxide causes a liquefaction necrosis in the most superficial layers.¹⁵⁶ The toxicity of calcium hydroxide appears to be neutralized as the deeper layers of pulp are affected, causing a coagulative necrosis at the junction of the necrotic and vital pulp, resulting in only mild irritation. This mild irritation initiates an inflammatory response, and in the absence of bacteria,¹⁵⁶ the pulp will heal with a hard-tissue barrier (Fig. 21.11).^{155,156} Hard-setting calcium hydroxide does not cause necrosis of the superficial layers of pulp; it has also been shown to initiate healing with a hard-tissue barrier.^{172,180}

The major disadvantage of calcium hydroxide is that it does not seal the fractured surface. Therefore an additional material must be used to ensure that bacteria do not challenge the pulp, particularly during the critical healing phase.

Many materials, such as zinc oxide eugenol,^{30,183} tricalcium phosphate,⁹⁸ and composite resin,²⁹ have been proposed as medicaments for vital pulp therapy. None to this date have afforded the predictability of calcium hydroxide used in conjunction with a well-sealed coronal restoration.^{70,122,161,173} In a dog study, for example, pulps capped directly with various adhesive agents showed moderate to severe inflammatory reactions, with progressive extension of tissue necrosis with time and total absence of continuous hard-tissue bridge formation.¹²² Application of a calcium hydroxide-based material was characterized by inflammatory cell infiltration, limited tissue necrosis, and partial to complete hard-tissue bridging.¹²² In a similar study on human teeth in vivo, it was found that Scotchbond Multi-Purpose Plus (3M, St. Paul, MN) caused inflammatory changes when applied directly to exposed pulp tissue; however, direct capping with Dycal followed by sealing with Scotchbond Multi-Purpose Plus showed favorable results in pulp tissue.¹⁷³

Currently, bioceramic materials are considered the pulp capping agents of choice.⁵ Mineral trioxide aggregate (MTA), a first-generation bioceramic material, has been shown to be a good pulp capping agent.^{72,139,141,195} It has a




Fig. 21.10 Pulp necrosis of 1.5 mm as a result of the high pH of calcium hydroxide.




Fig. 21.11 Hard-tissue barrier after calcium hydroxide partial pulpotomy. **A**, Histologic appearance of replacement odontoblasts and a hard-tissue barrier. **B**, Clinical appearance of the barrier on removal of the coronal restoration 3 months after placement of the calcium hydroxide. **C**, Radiographic appearance of the hard-tissue barrier (arrow).

high pH, similar to that of calcium hydroxide, when unset¹⁷⁷ and after setting creates an excellent bacteria-tight seal.¹⁷⁸ It is also hard enough to act as a base for a final restoration.¹⁷⁷ Yet MTA does not enjoy the same popularity as calcium hydroxide as a pulp capping agent in the treatment of traumatic exposures due to a prolonged setting time and discoloration of the crown because of the bismuth oxide filler in the material.

Recently, newer-generation bioceramic materials (e.g., BioDentine) have come to the market that have similar positive properties as MTA but not the disadvantages described previously. They set quickly enough for a one-visit procedure and do not discolor the tooth.¹⁹⁸ These materials are now considered superior to calcium hydroxide as the capping agent for traumatic pulp exposures.

TREATMENT METHODS

Pulp Capping

Pulp capping implies placing the dressing directly on the pulp exposure without any removal of the soft tissue.²⁹

Indications. There are few indications for pulp capping when traumatic pulp exposures are treated. The success rate of this procedure (80%),^{77,148} compared with partial pulpotomy (95%),⁵⁷ indicates that a superficial pulp cap should not be considered after traumatic pulp exposures. The lower success rate is not difficult to understand because superficial inflammation develops soon after the traumatic exposure. If the treatment is at the superficial level, a number of inflamed (rather than healthy) pulps will be treated, lowering the potential for success. In addition, a bacteria-tight coronal seal is much more difficult to attain in superficial pulp capping because there is no depth to the cavity to aid in creating the seal, as there is with a partial pulpotomy.

It must be acknowledged that the studies suggesting that an inflamed pulp is contraindicated for an attempt at pulp capping were performed in the 1970s with amalgam as the standard coronal restoration. From a microleakage perspective, as amalgam leaks and calcium hydroxide washes out in the presence of moisture as the reasons for the poor results, then it may be poor coronal seal, rather than the inflamed pulp, that leads to failure in these cases. More recent studies suggest that with the bioceramic as the pulp cap, the inflamed pulp is not the impediment as previously thought, but rather that the seal seems to be the major factor for success. Thus this material used as a base in these situations may provide more leeway for capping the inflamed pulp.

However, in a situation of a traumatic pulp exposure of a tooth (injuries in which most patients are young [large pulps]), the patient typically presents for treatment within 48 hours because of sensitivity and esthetic concerns. Therefore the pulpal inflammation is usually only superficial. In these cases, it may be prudent to remove the superficial layer of inflammation and place a bioceramic agent for pulp capping.

Partial Pulpotomy

Partial pulpotomy implies the removal of coronal pulp tissue to the level of healthy pulp. Knowledge of the reaction of the pulp after a traumatic injury makes it possible to accurately determine this level. This procedure is commonly called the *Cvek pulpotomy*.

Indications. As for pulp capping.

Technique. Administration of an anesthetic (possibly without a vasoconstrictor), rubber dam placement, and superficial disinfection are performed. A 1- to 2-mm-deep cavity is prepared into the pulp, using a high-speed handpiece with a sterile diamond bur of appropriate size and copious water coolant (Fig. 21.12).⁸⁵ A slow-speed bur or




Fig. 21.12 Cvek partial pulpotomy. **A**, The fractured teeth are cleaned and disinfected; a rubber dam is placed. **B**, Cavities are prepared at high speed with a round diamond bur 1 to 2 mm into the pulpal tissue. **C**, Calcium hydroxide on a plugger (**D**) is placed on the soft tissue of the pulp. **E**, Care is taken to avoid smearing the walls of the preparation with the calcium hydroxide. **F**, Cavity preparations are filled with glass ionomer cement. The exposed dentin is etched (**G**) and then covered with composite resin (**H**). **I**, Radiograph 6 months later shows formation of hard-tissue barriers in both teeth (arrows). (Courtesy Dr. Alessandra Ritter, Chapel Hill, NC.)

spoon excavator should be avoided. If bleeding is excessive, the pulp is amputated deeper until only moderate hemorrhage is seen. Excess blood is carefully removed by rinsing with sterile saline, and the area is dried with a sterile cotton pellet. Use of 5% sodium hypochlorite (NaOCl; bleach) has been recommended to rinse the pulpal wound.⁵⁴ The bleach causes chemical amputation of the blood coagulum; removes damaged pulp cells, dentin chips, and other debris; and provides hemorrhage control with minimal damage to the “normal” pulp tissue underneath.

Care must be taken not to allow a blood clot to develop because this would compromise the prognosis.^{57,155} Calcium hydroxide is carefully placed on the pulp stump. If the pulp size does not permit additional loss of pulp tissue, a commercial hard-setting calcium hydroxide can be used.¹⁵⁹ The prepared cavity is filled with a material with the best chance of a bacteria-tight seal such as a glass ionomer cement to a level flush with the fractured surface. The material in the pulpal cavity and all exposed dentinal tubules are etched and restored with bonded composite resin. Alternatively,

after hemostasis has been obtained, the pulp can be capped with MTA, covered with a lining material, and then restored with composite restoration. However, this has been shown to cause discoloration of the tooth.³⁵

Follow-Up. The follow-up is the same as for pulp capping. Emphasis is placed on evidence of maintenance of responses to sensitivity tests and radiographic evidence of continued root development (Fig. 21.13).

Prognosis. This method affords many advantages over pulp capping. Superficial inflamed pulp is removed during preparation of the pulpal cavity. Calcium hydroxide disinfects dentin and pulp and removes additional pulpal inflammation. Most important, space is provided for a material that can achieve a bacteria-tight seal to allow pulpal healing with hard tissue under optimal conditions. In addition, coronal pulp remains, which allows sensitivity testing to be carried out at the follow-up visits. The prognosis is extremely good (94% to 96%).^{57,78}

Full Pulpotomy

Full pulpotomy involves removal of the entire coronal pulp to a level of the root orifices. This level of pulp amputation is chosen arbitrarily because of its anatomic convenience. Because inflamed pulp sometimes extends past the canal orifices into the root pulp, many mistakes are made that result in treatment of an inflamed rather than a noninflamed pulp.

Indications. Full pulpotomy is indicated when it is anticipated that the pulp is inflamed to the deeper levels of the coronal pulp. Traumatic exposures after more than 72 hours and carious exposure of a young tooth with a partially developed apex are two examples of cases in which this treatment may be indicated. Because of the likelihood that the dressing will be placed on an inflamed pulp, full pulpotomy is contraindicated in mature teeth. In the immature tooth, benefits

outweigh risks for this treatment form only with incompletely formed apices and thin dentinal walls.

Technique. The procedure begins with administration of an anesthetic, rubber dam placement, and superficial disinfection, as for pulp capping and partial pulpotomy. The coronal pulp is removed as in partial pulpotomy but to the level of the root orifices. A calcium hydroxide dressing, a bacteria-tight seal, and a coronal restoration are carried out as for partial pulpotomy.

Follow-Up. Follow-up is the same as for pulp capping and partial pulpotomy. A major disadvantage of this treatment method is that sensitivity testing is not possible, owing to the loss of coronal pulp, so radiographic follow-up is extremely important to assess for signs of apical periodontitis and to ensure the continuation of root formation.

Prognosis. Because cervical pulpotomy is performed on pulps that are expected to have deep inflammation and the site of pulp amputation is arbitrary, many more mistakes are made that lead to treatment of the inflamed pulp. Consequently the prognosis, which is in the range of 75%, is poorer than for partial pulpotomy.⁸² Because of the inability to evaluate pulp status after full pulpotomy, some authors have recommended pulpectomy routinely after the roots have fully formed (Fig. 21.14). This philosophy is based on the pulpectomy procedure having a success rate in the range of 95%, whereas if apical periodontitis develops, the prognosis of root canal treatment drops significantly to approximately 80%.^{158,164}

Pulpectomy

Pulpectomy implies removal of the entire pulp to the level of the apical foramen.

Indications. Pulpectomy is indicated in a complicated crown fracture of mature teeth if conditions are not ideal




Fig. 21.13 Continued root development after partial pulpotomy. **A**, Radiograph of immature tooth with a complicated crown fracture. **B**, At the time of placement of calcium hydroxide after partial pulpotomy (arrow). **C**, Follow-up radiograph confirming that the pulp maintained vitality and the root continued to develop.




Fig. 21.14 Successful pulpotomy followed by a pulpectomy at 18 months. (Courtesy Dr. Leif Tronstad, Oslo, Norway.)

for vital pulp therapy or if it is foreseeable that restoring the tooth would require placement of a post. This procedure is no different from root canal treatment of a vital nontraumatized tooth.

TREATMENT OF THE NONVITAL PULP

Immature Tooth: Apexification

Indications. Apexification should be performed in teeth with open apices and thin dentinal walls in which standard instrumentation techniques cannot create an apical stop to facilitate effective root canal filling.

Biologic Consequences. A nonvital immature tooth presents a number of difficulties for adequate endodontic treatment. The canal is often wider apically than coronally, necessitating the use of a filling technique with softened filling material to mold it to the shape of the apical part of the canal. Because the apex is extremely wide, no barrier exists to stop this softened material from moving into and traumatizing the apical periodontal tissues. In addition, the lack of apical stop and extrusion of material through the canal might result in a canal that is underfilled and susceptible to leakage. An additional problem in immature teeth with thin dentinal walls is their susceptibility to fracture, both during and after treatment.⁵⁸

These problems are overcome by stimulating the formation of a hard-tissue barrier to allow for optimal filling of the canal and by reinforcing the weakened root against fracture, both during and after apexification.^{120,176}

Technique

DISINFECTION OF THE CANAL. In most cases, nonvital teeth are infected,^{36,160} so the first phase of treatment is to disinfect the root canal system to ensure periapical healing.^{46,61} The canal length is estimated with a parallel preoperative radiograph, and after access to the canals has been prepared, a file is placed to this estimated length. When the

length has been confirmed radiographically, very light filing (because of the thin dentinal walls) is performed with copious irrigation with 0.5% NaOCl.^{62,169} A lower strength of NaOCl is used because of the increased danger of placing the agent through the apex in immature teeth. The increased volume of irrigant used compensates for this lower concentration of NaOCl. An irrigation needle that can passively reach close to the apical length is useful in disinfecting the canals of these immature teeth. The canal is dried with paper points and a creamy mix of calcium hydroxide (toothpaste thickness) spun into the canal with a Lentulo spiral instrument (Fig. 21.15). Additional disinfecting action of calcium hydroxide is effective after its application for at least 1 week,¹⁶³ so continuation of treatment can take place any time after 1 week. Further treatment should not be delayed more than 1 month because the calcium hydroxide could be washed out by tissue fluids through the open apex, leaving the canal susceptible to reinfection.

Hard-Tissue Apical Barrier

TRADITIONAL METHOD. Formation of the hard-tissue barrier at the apex requires a similar environment to that required



Fig. 21.15 Creamy mix of calcium hydroxide on a Lentulo spiral instrument ready for placement into the canal.

for hard-tissue formation in vital pulp therapy: a mild inflammatory stimulus to initiate healing and a bacteria-free environment to ensure that inflammation is not progressive.

As with vital pulp therapy, calcium hydroxide is used for this procedure.^{58,91,99} Calcium hydroxide is packed against the apical soft tissue with a plugger or thick point to initiate hard-tissue formation. This step is followed by backfilling with calcium hydroxide to completely fill the canal. The calcium hydroxide is meticulously removed from the access cavity to the level of the root orifices, and a well-sealing, temporary filling is placed in the access cavity. A radiograph is taken; the canal should appear to have become calcified, indicating that the entire canal has been filled with the calcium hydroxide (Fig. 21.16). Because calcium hydroxide washout is evaluated by its relative radiodensity in the canal, it is prudent to use a calcium hydroxide mixture without the addition of a radiopaquer, such as barium sulfate. These additives do not wash out as readily as calcium hydroxide, so if they are present in the canal, evaluation of washout is impossible.

At 3-month intervals, a radiograph is exposed to evaluate whether a hard-tissue barrier has formed and if the calcium hydroxide has washed out of the canal. This is assessed to have occurred if the canal can be seen again radiographically. If no washout is evident, the calcium hydroxide can be left intact for another 3 months. Excessive calcium hydroxide dressing changes should be avoided if at all possible because the initial toxicity of the material is thought to delay healing.¹²⁵

When completion of a hard-tissue barrier is suspected, the calcium hydroxide should be washed out of the canal with NaOCl. A file of a size that can easily reach the apex can be used to gently probe for a stop at the apex. When a hard-tissue barrier is indicated radiographically and can be probed with an instrument, the canal is ready for filling.

Bioceramic Barrier. The creation of a physiologic hard-tissue barrier, although quite predictable, takes anywhere from 3 to 18 months with calcium hydroxide. The disadvantages of this long time period are that the patient is

required to present for treatment multiple times and the tooth may fracture during treatment before the thin, weak roots can be strengthened. In addition, one study has indicated that long-term treatment with calcium hydroxide may weaken the roots and make them even more susceptible to fracture.²⁴ However, this is not conclusive because another more recent study showed that calcium hydroxide did not weaken teeth over time when compared with positive and negative controls and that fracture. Therefore fracture may be more related to the stage of root development rather than calcium hydroxide use.¹¹⁵

Bioceramic material has been used to create a hard-tissue barrier quickly after disinfection of the canal (see earlier in the section Traditional Method) (Fig. 21.17). Calcium sulfate is pushed through the apex to provide a resorbable extraradicular barrier against which to pack the bioceramic material. The material is mixed and packed into the apical 4 to 5 mm of the canal. Historically, a wet cotton pellet was placed against the bioceramic material and left for at least 6 hours and this is now not considered necessary as long as the bioceramic material does not dry out. The canal is filled with thermoplasticized gutta-percha. The cervical canal is then reinforced with composite resin to below the marginal bone level (described later; also see Fig. 21.17).

A number of case reports have been published using this apical barrier technique,^{83,101,131} and it has steadily gained popularity with clinicians. Numerous reports attest to the efficacy of this technique.^{48,114}

FILLING THE ROOT CANAL. Because the apical diameter is larger than the coronal diameter of most of these canals, a softened filling technique is indicated in these teeth (see Chapter 9). Care must be taken to avoid excessive lateral force during filling, owing to the thin walls of the root. If the hard-tissue barrier was produced by long-term calcium hydroxide therapy, it consists of irregularly arranged layers of coagulated soft tissue, calcified tissue, and cementum-like tissue (Fig. 21.18). Included also are islands of soft connective tissue, giving the barrier a “Swiss cheese” consistency.^{40,56} Because of the irregular nature of the barrier, it is not




Fig. 21.16 A and B, Root canal that “disappears” after placement of a thick mix of pure calcium hydroxide and then over time washes out again. (Courtesy Dr. Cecilia Bourguignon, Paris, France.)




Fig. 21.17 Apexification with mineral trioxide aggregate. **A**, The canal is disinfected with light instrumentation, copious irrigation, and a creamy mix of calcium hydroxide for 1 month. **B**, Calcium sulfate is placed through the apex as a barrier against which the mineral trioxide aggregate (MTA) is placed. **C**, A 4-mm MTA plug is placed at the apex. **D**, The body of the canal is filled with the Resilon obturation system. **E**, A bonded resin is placed below the cemento-enamel junction (CEJ) to strengthen the root. (Courtesy Dr. Marga Ree, Purmerend, Netherlands.)




Fig. 21.18 Histologic appearance of hard-tissue barrier after calcium hydroxide apexification. The barrier is composed of cementum and bone with soft-tissue inclusions.

unusual for cement or softened filling material to be pushed through it into the apical tissues during filling (Fig. 21.19). Formation of the hard-tissue barrier might be some distance short of the radiographic apex because the barrier forms wherever the calcium hydroxide contacts vital tissue. In teeth with wide, open apices, vital tissue can survive and proliferate from the periodontal ligament a few millimeters into the root canal. Filling should be completed to the level of the hard-tissue barrier and not forced toward the radiographic apex.

REINFORCEMENT OF THIN DENTINAL WALLS. The apexification procedure has become a predictably successful procedure (see the section Prognosis),^{76,91} but thin dentinal walls still present a clinical problem. Should secondary injuries occur, teeth with thin dentinal walls are more susceptible to fractures that render them unrestorable.^{65,179} It has been reported that approximately 30% of these teeth will fracture during or after endodontic treatment.¹²⁰ Consequently, some clinicians have questioned the advisability of the

apexification procedure and have opted for more radical treatment procedures, including extraction followed by extensive restorative procedures, such as dental implants. However, this approach cannot be considered in children or young adults because facial growth has not been completed. Studies have shown that intracoronally bonded restorations can internally strengthen endodontically treated teeth and increase their resistance to fracture.^{84,118} Thus, after root filling, the material should be removed to below the marginal bone level and a bonded resin filling placed (see Fig. 21.17).

FOLLOW-UP. Routine recall evaluation should be performed to determine success in the prevention or treatment of apical periodontitis. Restorative procedures should be assessed to ensure that they in no way promote root fractures.



Fig. 21.19 Root filling with a soft technique after calcium hydroxide apexification. Sealer and softened filling material are expressed through the "Swiss cheese" holes in the barrier.

PROGNOSIS. Periapical healing and the formation of a hard-tissue barrier occur predictably with long-term calcium hydroxide treatment (79% to 96%).^{58,120} However, long-term survival is jeopardized by the fracture potential of the thin dentinal walls of these teeth. It is expected that the newer techniques of internally strengthening the teeth described earlier will increase their long-term survivability.

PULP REVASCULARIZATION/REGENERATIVE ENDODONTIC TREATMENT. Regeneration of a necrotic pulp has been considered possible only after avulsion of an immature permanent tooth that was reimplanted within 40 minutes (discussed later). The advantages of pulp revascularization lie in the possibility of further root development and reinforcement of dentinal walls by deposition of hard tissue, thus strengthening the root against fracture (see also Chapter 12). After reimplantation of an avulsed immature tooth, a unique set of circumstances exists that allows regeneration to take place. The young tooth has an open apex and is short, which allows new tissue to grow into the pulp space relatively quickly. The pulp is necrotic but usually neither degenerated nor infected, so it acts as a matrix into which the new tissue can grow. It has been experimentally shown that the apical part of a pulp may remain vital and after reimplantation proliferate coronally, replacing the necrotized portion of the pulp.^{30,140,167} Because in most cases the crown of the tooth is intact and caries free, bacterial penetration into the pulp space through cracks¹²⁷ and defects will be a slow process. The race between new tissue growth and infection of the pulp space favors the new tissue.¹⁴

Traditionally, regeneration of pulp tissue in a necrotic infected tooth with apical periodontitis was thought to be impossible.¹³⁸ Regeneration/repair can occur when the canal is effectively disinfected, a matrix into which new tissue

can grow is provided, and coronal access is effectively sealed.^{30,105} There have been many studies in the past few years aimed at making the procedure more predictable both in healing of apical periodontitis and in promoting pulp and not nonspecific connective tissue (see Chapter 12). A clinical protocol can be found in the AAE "Clinical Considerations for a Regenerative Procedure." However, although there is the potential for further root maturation, a prospective study has shown that increases in root width and length are variable in a cohort of predominantly traumatized teeth (Fig. 21.20).¹¹²

In regenerative endodontic procedures there are three outcome measures for success. The primary goal is the elimination of symptoms and the evidence of bony healing. The secondary goal is increased root wall thickness and/or increased root length, which is considered desirable but perhaps not essential. The tertiary goal is a positive response to vitality testing (which if achieved, could indicate a more organized vital pulp tissue).¹ It is known that results are variable for these objectives and true regeneration of the pulp/dentin complex is not achieved. Repair derived primarily from the periodontal and osseous tissues has been shown histologically.^{34,132}

Disinfection of the canal involved no mechanical preparation, but copious irrigation with NaOCl was performed, followed by placement of an intracanal mixture of antibiotics consisting of equal amounts of ciprofloxacin, metronidazole, and minocycline, to a final concentration of 1 to 5 mg/mL.^{1,103} The minocycline in the triantibiotic paste causes unacceptable discoloration of the teeth¹¹³; this has led to many studies on eliminating or replacing this antibiotic in the paste or eliminating the tripaste. Alternatively, calcium hydroxide can be used as an intracanal




Fig. 21.20 **A**, Preoperative radiograph showing an immature incisor (tooth #9) that sustained an uncomplicated crown fracture where pulp necrosis developed as shown by cessation of root development when compared with tooth #8. **B**, A 6-year review radiograph showing the tooth after treatment with a regenerative endodontic protocol. There were no signs of infection (primary goal). Calcific material at the apex is present indicating apical closure. A quantitative analysis as described by Bose et al.⁴⁴ reports a 1.9% increase in root length and a 5% increase in canal wall width (secondary goal). The tooth was responsive to electric pulp testing in a comparable range to tooth #8 (tertiary goal).

medicament rather than an antibiotic paste. Furthermore, Biodentine when used as an intracanal barrier results in less crown discoloration than when MTA is used. Avoiding discoloration of the crown is a patient-centered outcome.

Studies are underway to find a synthetic matrix that can act as a more predictable scaffold for new ingrowth of tissue than the introduction of a blood clot into the canal.^{121,194} Studies are also underway on stem cells and growth factors aimed at a more predictable production of odontoblasts and thus pulp cells. It is hoped that with the concept of tissue engineering, namely stem cells, scaffolds, and signaling molecules, that true pulp regeneration is an achievable goal.¹²¹ The procedure described here can be attempted in most cases and has been recommended as treatment of choice for immature teeth with incomplete root formation, although teeth with near or complete root formation may be more suited for conventional endodontic therapy or MTA barrier techniques.¹²¹ If there are no signs of regeneration or continued signs of infection, the more traditional treatment methods for treatment of immature teeth can be initiated.

Mature Tooth

Conventional endodontic therapy is recommended for mature teeth with closed apex.

Crown-Root Fracture

Crown-root fracture is a periodontal rather than an endodontic challenge. The tooth must be treated periodontally to enable a well-sealed coronal restoration. This could be accomplished by simple gingivectomy if the extent of the root component of the fracture is large. Alternatively, the tooth could be orthodontically or surgically extruded such that the exposed surface of the root fracture is treatable. Once the feasibility of the coronal restoration is ensured, the particular crown fracture is treated as previously described.

Root Fracture

Root fracture implies fracture of the cementum, dentin, and pulp. These injuries are relatively infrequent, occurring in less than 3% of all dental injuries.¹⁹⁹ Immature teeth with vital pulps rarely sustain horizontal root fractures.¹⁰ When a root fractures horizontally, the coronal segment is displaced to a varying degree, but generally the apical segment is not displaced. Because the apical pulpal circulation is not disrupted, pulp necrosis in the apical segment is extremely rare. Permanent pulpal necrosis of the coronal segment, requiring endodontic treatment, occurs in approximately 25% of cases.^{11,12,107}

DIAGNOSIS AND CLINICAL PRESENTATION

The clinical presentation of a root fracture is similar to that of luxation injuries. The extent of displacement of the coronal segment is usually indicative of the location of the fracture and can vary from none, simulating a concussion injury (apical fracture), to severe, simulating an extrusive luxation (cervical fracture).

Radiographic examination for root fractures is extremely important. Because root fractures are usually oblique

(facial to palatal; Fig. 21.21), one periapical radiograph may easily miss it. It is imperative to take at least three angled radiographs (45 degrees, 90 degrees, and 110 degrees) so that at least at one angulation, the x-ray beam passes directly through the fracture line to make it visible on the radiograph (Fig. 21.22).

TREATMENT

Emergency treatment involves repositioning the segments into close proximity as much as possible. In the case of severe displacement of the coronal segment, its apical extension is frequently lodged in (if not perforating through) the cortical bone facial to the tooth. Forcing the crown facially is not possible, and the two segments will not be properly aligned. The only way to accomplish reapproximation of the two segments is to release the coronal segment from the bone by gently pulling it slightly downward with finger pressure or extraction forceps and, once it is loose, rotating it back to its original position (Fig. 21.23). The traditionally recommended¹⁴⁷ splinting protocol has been changed from 2 to 4 months with rigid splinting to a flexible splint to adjacent teeth for 2 to 4 weeks.^{23,69,110} If a long time has elapsed between the injury and treatment, repositioning of the segments close to their original position probably will not be possible; this compromises the long-term prognosis for the tooth.

HEALING PATTERNS

Investigators²⁶ have described four types of responses to root fractures.

1. *Healing with calcified tissue.* Radiographically, the fracture line is discernible, but the fragments are in close contact (Fig. 21.24, A).
2. *Healing with interproximal connective tissue.* Radiographically, the fragments appear separated by a narrow



Fig. 21.21 Extracted root after a root fracture. Note the oblique angle of the fracture.

radiolucent line, and the fractured edges appear rounded (see Fig. 21.24, B).

3. *Healing with interproximal bone and connective tissue.* Radiographically, the fragments are separated by a distinct bony ridge (see Fig. 21.24, C).
4. *Interproximal inflammatory tissue without healing.* Radiographically, a widening of the fracture line and/or a developing radiolucency corresponding to the fracture line becomes apparent (see Fig. 21.24, D).

The first three healing patterns are considered successful. The teeth are usually asymptomatic and respond positively

to sensitivity testing. Histologically, healing can be derived from pulpal and periodontal tissues (Fig. 21.25 for healing with calcified tissue; Fig. 21.26 for healing with interproximal connective tissue).⁹⁷

Coronal yellowing is possible because calcific metamorphosis of the coronal segment is not unusual (Fig. 21.27).^{108,199}

The fourth type of root fracture response is typical when the coronal segment loses its vitality. The infective products in the coronal pulp cause an inflammatory response and typically radiolucencies are at the fracture line (see Fig. 21.24, D).²⁶




Fig. 21.22 Radiographs showing the importance of different vertical angulations for diagnosis of root fracture. All three radiographs were taken within minutes of each other.



Fig. 21.23 Figure of facial root fracture manipulated back into place.

TREATMENT OF COMPLICATIONS

Coronal Root Fractures

Historically it had been thought that fractures in the cervical segment had a poor prognosis, and extraction of the coronal segment was recommended. Research does not

support this treatment; in fact, if these coronal segments are adequately splinted, chances of healing do not differ from those for midroot or apical fractures (Fig. 21.28).¹⁹⁹ However, if the fracture occurs at the level of or coronal to the crest of the alveolar bone, the prognosis is extremely poor.

If reapproximation of the fractured segments is not possible, extraction of the coronal segment is indicated. The level of fracture and length of the remaining root are evaluated for restorability. If the apical root segment is long enough, gentle orthodontic eruption of this segment can be carried out to enable fabrication of a restoration.

Midroot and Apical Root Fractures

Permanent pulpal necrosis occurs in 25% of root fractures.^{56,108} Initially it is likely that in many cases the pulp in the coronal segment will become necrotic after the injury; however, because of a very large apical opening in the coronal segment, revascularization is possible if the segments are well reapproximated. In most cases, permanent necrosis occurs in the coronal segment, with only the




Fig. 21.24 Healing patterns after horizontal root fractures. **A1**, A transverse root fracture in a 7.5-year-old boy. **A2**, Healing with calcified tissue with internal thickening of root dentin into the pulp chamber at a 3-year review. **B**, Healing with interproximal connective tissue. **C**, Healing with bone and connective tissue. Note extensive intracanal calcification of tooth #9. **D**, Interproximal connective tissue without healing. (Courtesy Professor Geoffrey Heithersay, Adelaide, Australia.)




Fig. 21.25 **A**, Sagittal histologic section in a labiopalatal plane showing pulp and hard-tissue deposition between the fractured segments on the labial and to a greater extent on the palatal aspect (Van Gieson stain, original magnification $\times 4$). **B**, Labial section showing hard-tissue deposition between approximately half of the fractured dentin surfaces and bulging into the root canal space (Van Gieson stain, original magnification $\times 10$). **C**, Higher-power view of 25A showing fibrillar dentin deposit (FD) with some small and larger cell/tissue inclusions and tubular dentin (TD) with a sunray orientation bulging into the root canal and lined by pulp (original magnification $\times 20$). **D**, Polarized light view of the pulpal extremity of dentin deposition showing the tubular orientation and deeper irregular hard-tissue formation with cell/tissue inclusions (original magnification $\times 25$). **E**, Palatal section showing hard-tissue deposition between the majority of the fractured dentin surfaces and budding into the root canal. The calcified tissue deposits consist of tubular dentin (TD) fibrillar dentin (FD) with cell/tissue inclusions, considered to be of pulpal origin (P), and sclerotic calcified tissue (SCT), considered to be of periodontal ligament origin (Van Gieson stain, original magnification $\times 10$). **F**, Higher-power view of the calcified tissue considered to have been pulpal deposited showing further details of the orientation of the dentinal tubules in the tubular dentin layers (TD) and the deeper fibrillar dentin layers (FD) (original magnification $\times 20$). (Reproduced from Heithersay GS, Kahler B: Healing responses following transverse root fracture: a historical review and case reports showing healing with (a) calcified tissue and (b) dense fibrous connective tissue, *Dent Traumatol* 29:253, 2013.)

apical segment remaining vital. Therefore endodontic treatment is indicated in the coronal root segment unless periapical pathology is seen in the apical segment. In most cases the pulpal lumen is wide at the apical extent of the coronal segment, and long-term calcium hydroxide treatment or an MTA apical plug is indicated (see the section Apexification). The coronal segment is filled after a hard-tissue barrier has formed apically in the coronal segment and periradicular healing has taken place.

In rare cases in which both the coronal and apical pulp are necrotic, treatment is more complicated. Endodontic treatment through the fracture is extremely difficult. Endodontic manipulations, medicaments, and filling materials all have a detrimental effect on healing of the fracture site. If healing of the fracture has been completed and is followed by necrosis of the apical segment, the prognosis is much improved.

In more apical root fractures, necrotic apical segments can be surgically removed. This is a viable treatment if the remaining coronal root segment is long enough to provide adequate periodontal support. Removal of the apical segment in midroot fractures leaves the coronal segment with a compromised attachment.

Follow-Up

After the splinting period is completed, follow-up is as for all dental traumatic injuries: at 3, 6, and 12 months and yearly thereafter for at least 5 years.

Prognosis

The following factors influence repair:

1. The degree of dislocation and mobility of the coronal fragment are extremely important in determining outcome.^{10,22,108,170,199} Increased dislocation and coronal fragment mobility result in a poorer prognosis.
2. Immature teeth are seldom involved in root fractures, but in the unlikely event they are, the prognosis is good.^{22,106}
3. The quality of treatment is vital to successful repair. The prognosis improves with quick treatment, close reduction of the root segments, and semirigid splinting for 2 to 4 weeks.²³

Complications include pulp necrosis and root canal obliteration. Pulp necrosis can be treated successfully^{56,108} by treating the coronal segment with calcium hydroxide to stimulate hard-tissue barrier formation. Root canal obliteration is common if the root segment (coronal or apical) remains vital.




Fig. 21.26 **A**, The intact central incisor after extraction and formalin fixation. **B**, The histologic appearance of the extracted tooth sectioned in a mesiodistal sagittal plane showing the interposition of fibrous connective tissue (*FCT*) calcification of the root canals but evidence of residual pulp tissue (*P*) (H&E, original magnification $\times 4$). **C**, View of the fracture site showing dentin resorption lines (*RL*) repaired with reparative hard tissue (*RHT*) and the interposed dense fibrous connective tissue (*DFCT*) (H&E, original magnification $\times 10$). **D**, Higher-power view showing dense connective tissue, generally with a parallel orientation to the fractured dentin surfaces that show reparative hard tissue laid down on previously resorbed dentin. There is minimal vascularity, but some extravasated erythrocytes are evident (Van Gieson stain, original magnification $\times 20$). **E**, Highlighted section from (**C**) using polarized light to demonstrate Sharpey fiber insertion into the reparative hard tissue (*arrows*) (original magnification $\times 40$). (Reproduced from Heithersay GS, Kahler B: Healing responses following transverse root fracture: a historical review and case reports showing healing with (a) calcified tissue and (b) dense fibrous connective tissue, *Dent Traumatol* 29:253, 2013.)

Luxation Injuries

DEFINITIONS

The types of luxation injury can be defined as follows:

1. *Concussion* implies no displacement, normal mobility, and sensitivity to percussion.
2. *Subluxation* implies sensitivity to percussion, increased mobility, and no displacement.
3. *Lateral luxation* implies displacement labially, lingually, distally, or incisally.
4. *Extrusive luxation* implies displacement in a coronal direction.
5. *Intrusive luxation* implies displacement in an apical direction into the alveolus.

Definitions 1 through 5 describe injuries of increasing magnitude in terms of intensity of the injury and subsequent sequelae.

INCIDENCE

Luxation injuries as a group are the most common of all dental injuries, with a reported incidence ranging from 30% to 44%.⁶³

TREATMENT

Teeth that are concussed or subluxated do not need any immediate treatment. Responses to vitality tests should be investigated and noted. Even after mild injury, such as subluxation, the pulp might be unresponsive to vitality tests for several weeks if not months.¹⁶⁶ When the pulp is unresponsive initially after the trauma, patients should be recalled on a regular basis and monitored for any additional signs of infection of the root canal (discussed later).

Teeth with lateral and extrusive luxation should be repositioned as soon as possible. In lateral luxation, the apex might be perforating the facial bone plate, and the tooth




Fig. 21.27 **A**, Pulp canal calcification after a luxation injury. Radiographic appearance of a calcified maxillary lateral incisor; note that despite calcification, it has now become necrotic and infected, evident by the periapical lesion. **B**, The typical yellow appearance of the tooth caused by thickened dentin in the pulp chamber.

must be slightly and gently pulled down to loosen the hold before it is repositioned in its original position. Current IADT guidelines call for 2 weeks of physiologic splinting in cases of extrusion luxation and 4 weeks for lateral luxation. A decision on root canal therapy follows the guidelines for

avulsion (discussed later). If the tooth has a fully formed apex and was diagnosed to have moved into (if not through) the cortical plate (apical translocation), there is a good likelihood of the pulp being devitalized; therefore endodontic treatment should be initiated as early as 2 weeks after the injury. If the apex is still not fully formed, waiting for signs of revascularization is strongly recommended.

Permanent teeth that are intruded are not likely to spontaneously reerupt, especially if the apex is fully formed.¹¹⁹ Alternative treatment, such as orthodontic extrusion or immediate surgical repositioning, should be considered. If orthodontic extrusion is planned for an intruded tooth, it should be initiated as soon as possible and should not be delayed longer than 2 to 3 weeks after the trauma. Only a few studies have evaluated the true efficacy of this approach. One study¹⁸⁹ using a dog model indicated that severely intruded teeth showed signs of ankylosis as early as 11 to 13 days after the trauma, despite initiation of orthodontic movement 5 to 7 days after the injury.

For the surgical approach, one study⁵⁵ (using a dog model) concluded that “a careful immediate surgical repositioning of a severely intruded permanent tooth with complete root formation has many advantages with few disadvantages.” Another investigation,⁶⁷ a retrospective study of 58 intruded human teeth, indicated that surgical repositioning resulted in a predictable outcome, with only five of the teeth lost over the observation period. However, it was also observed that less surgical manipulation positively influenced the healing. If an intruded tooth is immediately reimplanted, it should be splinted for at least 4 weeks, but in most cases the splint needs to be left on the tooth longer.

BIOLOGIC CONSEQUENCES

Luxation injuries result in damage to the attachment apparatus (periodontal ligament and cemental layer), the severity of which depends on the type of injury sustained




Fig. 21.28 Internal root resorption in the root fracture area; tooth was diagnosed with compound midroot fracture. **A**, Coronal portion was repositioned and then splinted to the lateral incisor with a composite splint. **B**, At 6-month recall, the pulp responded normally to cold, but an internal root-resorptive defect was noted on radiograph. No treatment was indicated, because the pulp did respond normally to vitality tests. **C**, At 42-month recall, the tooth was still asymptomatic, and the pulp still responded to vitality tests. The resorptive defect had healed, and signs of dystrophic calcifications were evident in the apical and coronal segments.

(concussion least, intrusion most). The apical neurovascular supply to the pulp is also affected to varying degrees, resulting in an altered or nonvital tooth.

Healing can be favorable or unfavorable. Favorable healing after a luxation injury occurs if the initial physical damage to the root surface and the resultant inflammatory response to the damaged external root surface are again covered with cementum. An unfavorable response occurs when there is direct attachment of bone to the root, with the root ultimately being replaced by bone.

There are two resorption responses in which the pulp plays an essential role:

1. In *external* inflammatory root resorption, the *necrotic infected pulp* provides the stimulus for periodontal inflammation in the ligament space. If the cementum has been damaged, the inflammatory stimulators in the pulp space are able to diffuse through the dentinal tubules and stimulate an inflammatory response over large areas of the periodontal ligament. Because of the lack of cemental protection, the periodontal inflammation includes root resorption in addition to the expected bone resorption.
2. With *internal* inflammatory root resorption, the inflamed pulp is the tissue involved in resorbing the root structure. The pathogenesis of internal root resorption is not completely understood. Here it is thought that coronal necrotic infected pulp provides a stimulus for a pulpal inflammation in the more apical parts of the remaining vital pulp.¹⁰ If the internal root surface has lost its precemental protection during an injury, internal root resorption occurs in the area adjacent to the inflamed pulp. Thus both the necrotic infected pulp and the inflamed pulp contribute to this type of root resorption.

External Root Resorption

Caused by an Injury (Alone) to the External Root Surface. If an injury harms the attachment, the by-products of this mechanical damage stimulate an inflammatory response. The healing response depends on the extent of the initial damage.

LOCALIZED INJURY: HEALING WITH CEMENTUM. When the injury is localized (e.g., after concussion or subluxation injury), mechanical damage to the cementum occurs, resulting in a local inflammatory response and a localized area of root resorption. If no further inflammatory stimulus is present, periodontal healing and root surface repair occur within 14 days (Fig. 21.29).⁸⁶ The resorption is localized to the area of mechanical damage, and treatment is not required because it is free of symptoms and not even visualized radiographically in most cases. However, in a minority of cases, small radiolucencies can be seen on the root surface if the radiograph is taken at a specific angle. It is important to avoid misinterpreting these cases as progressive. The pulp is not involved. If the pulp responds to sensitivity tests, this is a clue that no treatment should be performed. A wait-and-see attitude can allow spontaneous healing to take place. It is important to realize that in the initial stages, radiolucency could be followed by spontaneous repair, and endodontic treatment should *not* be initiated.




Fig. 21.29 Histologic section showing previous root resorptive defect healed with new cementum and periodontal ligament.

DIFFUSE INJURY: HEALING BY REPLACEMENT RESORPTION. When the traumatic injury is severe (e.g., intrusive luxation or avulsion with extended dry time), involving diffuse damage on more than 20% of the root surface, an abnormal attachment can occur after healing takes place.¹²⁶ The initial reaction, as always, is inflammation in response to the severe mechanical damage to the root surface. After the initial inflammatory response, a diffuse area of root surface devoid of cementum results. Cells in the vicinity of the denuded root now compete to repopulate it. Often cells that are precursors of bone, rather than the slower-moving periodontal ligament cells, move across from the socket wall and populate the damaged root. Bone comes into contact with the root without an intermediate attachment apparatus. This phenomenon is termed *dentoalveolar ankylosis*.¹²⁶ Ankylosis results in failure of the tooth to erupt, leading to infraposition in the developing dentition and a high-pitch sound to percussion. However, the ankylosis and replacement resorption that follows cannot be reversed and can be considered a physiologic process because bone resorbs and reforms throughout life. Thus root is resorbed by osteoclasts, but in the reforming stage, bone is laid down instead of dentin, slowly replacing the root with bone. This process is termed *replacement resorption* (Fig. 21.30).^{8,19} However, in these traumatic cases, the resorptive phase includes the root being “replaced” with bone (Fig. 21.31). An alternative term of “osseous replacement” has been used.¹⁸⁶

TREATMENT. Treatment strategies for limiting the effect of the traumatic injury to the periodontal structures are outside the scope of this chapter. However, in general they involve minimizing the initial inflammatory response to the injury. The initial inflammation is destructive in nature and increases the surface area of root to be covered in the healing phase.²⁵ The smaller the surface area to be covered by new cementum, the higher the chances of favorable repair.




Fig. 21.30 Histologic appearance of replacement resorption. There is direct contact between bone and root structure. Resorptive defects are seen in the bone and root tissue, which is the normal physiologic process of bone turnover. The resorptive defects are filled by new bone and additional areas resorbed. In this way the entire root is replaced by bone at a rate dependent on the metabolic rate of the patient.



Fig. 21.31 Radiographic appearance of replacement resorption. The root acquires the radiographic appearance of the surrounding bone (without a lamina dura). Note that radiolucencies typical of active inflammation are not present.

A study has indicated that if Ledermix, a drug combining corticosteroid and tetracycline, is placed in the root canal immediately after a severe trauma in which replacement resorption is expected, favorable healing occurs at a very high rate.⁴⁵ In a more recent study, it was shown that triamcinolone (the corticosteroid portion of the Ledermix paste) was as effective as Ledermix at inhibiting external root resorption.⁴⁹ Both of these studies were done on young dogs and need to be replicated in human studies.

Caused by an Injury to the External Root Surface and Inflammatory Stimulus in the Root Canal. Recognized inflammatory stimuli that cause root resorption are pressure,

pulp space infection, and sulcular infection. This chapter focuses on pulp space infection.

Consequences of Apical Neurovascular Supply Damage

PULP CANAL OBLITERATION (CALCIFICATION). *Pulp canal obliteration* is common after luxation injuries (see Fig. 21.27). The frequency of pulp canal obliteration appears inversely proportional to pulp necrosis. The exact mechanism of pulp canal obliteration is unknown. It has been theorized that the sympathetic/parasympathetic control of blood flow to the odontoblasts is altered, resulting in uncontrolled reparative dentin.^{10,20} Another theory is that hemorrhage and blood clot formation in the pulp after injury form a nidus for subsequent calcification if the pulp remains vital.^{10,20} One study¹⁶ found that pulp canal obliteration could usually be diagnosed within the first year after injury and was more frequent in teeth with open apices (>0.7 mm radiographically), those with extrusive and lateral luxation injuries, and those that had been rigidly splinted.¹⁶

PULP NECROSIS. The factors most important for the development of pulp necrosis are the type of injury (concussion least, intrusion most) and the stage of root development (mature apex more than an immature apex).¹⁵ Pulp necrosis will most likely to lead to infection of the root canal system, with problematic consequences.

PULP SPACE INFECTION. Pulp space infection in conjunction with damage to the external root surface results in periradicular root and bone resorption and continues in its active state as long as the pulpal stimulus (infection) remains. When the root loses its cemental protection, lateral periodontitis with root resorption can result (Fig. 21.32). CBCT imaging can show the extent of the resorption (Fig. 21.33).

To have pulp space infection, the pulp must first become necrotic. Necrosis occurs after a fairly serious injury in which displacement of the tooth results in severing of the apical blood vessels. In mature teeth, pulp regeneration cannot occur, and usually by 3 weeks, the necrotic pulp becomes infected. (For details of the typical bacterial contents of a traumatized necrotic pulp, the reader is referred to Chapter 15 or to Bergenholz.³⁶) Because a serious injury is required for pulp necrosis, areas of cemental covering of



Fig. 21.32 Inflammatory root resorption caused by a pulp space infection. Note the radiolucencies in the root and surrounding bone. (Courtesy Dr. Fred Barnett.)




Fig. 21.33 **A**, Periapical radiograph of tooth #9 of a 16-year-old boy showing a mature tooth 6 months after avulsion and replantation within 5 minutes. Inflammatory root resorption is occurring as endodontic treatment was not commenced. **B** and **C**, CBCT sagittal images taken same day as the periapical radiograph. **D**, CBCT axial images.

the root usually are also affected, resulting in loss of its protective (insulating) quality. Now bacterial toxins can pass through the dentinal tubules and stimulate an inflammatory response in the periodontal ligament. The result is resorption of the root and bone. The periodontal infiltrate consists of granulation tissue with lymphocytes, plasma cells, and polymorphonuclear leukocytes. Multinucleated giant cells resorb the denuded root surface, and this continues until the stimulus (pulp space bacteria) is removed (Fig. 21.34).¹⁸¹ Radiographically, the resorption is observed as progressive radiolucent areas of the root and adjacent bone (see Fig. 21.32).

TREATMENT. Assessing attachment damage caused by the traumatic injury and minimizing the subsequent inflammation should be the focus of the emergency visit. The clinician's attention to pulp space infection should ideally be 7 to 10 days after the injury.^{187,188} Root canal disinfection removes the stimulus to the periradicular inflammation, thus the resorption will stop.^{86,187,188} In most cases a new attachment will form, but if a large area of root is affected, replacement resorption can result by the mechanism already described. Again, treatment principles include prevention of pulp space infection or elimination of the bacteria if they are present in the pulp space.




Fig. 21.34 Histologic appearance of multinucleated osteoclasts (dentoclasts) resorbing the dentin of the root.

1. Prevention of pulp space infection

- a. *Reestablish the vitality of the pulp.* If the pulp stays vital, the canal will be free of bacteria, and external inflammatory root resorption will not occur. In severe injuries in which vitality has been lost, it is possible under some circumstances to promote revascularization of the pulp. Revascularization is possible in young teeth with incompletely formed apices if the teeth are replaced in their original position within 60 minutes of the injury (Fig. 21.35).⁵⁹ If the tooth has been avulsed, soaking it in doxycycline for 5 minutes or covering the root with minocycline powder before replantation has been shown to double or triple the revascularization rate.^{59,150} However, even under the best conditions, revascularization fails to occur on many occasions, and a diagnostic dilemma results. If the pulp revascularizes, external root resorption will not occur, and the




Fig. 21.35 Revascularization of immature root. A tooth with an open apex was replanted soon after the avulsion. The checkup radiograph 12 years later confirms that regrowth has taken place into the pulp chamber. It appears that a bone plug has grown in, and new periodontal ligament has formed, with a lamina dura within the pulp canal. (Courtesy Dr. Cecilia Bourguignon, Paris, France.)

root will continue to develop and strengthen. If the pulp becomes necrotic and infected, the subsequent external inflammatory root resorption that develops could result in loss of the tooth in a very short time. At present, the diagnostic tools available cannot detect a vital pulp in this situation before approximately 6 months after successful revascularization. This period of time is obviously unacceptable because by that time the teeth that have not revascularized could be lost to the resorption process. Recently the laser Doppler flowmeter or the pulse oximeter has been shown to have diagnostic potential for the detection of revascularization in immature teeth (see Fig. 21.2). These devices appear to detect the presence of vital tissue in the pulp space by 4 weeks after the traumatic injury.¹⁹⁷

- b. *Prevent root canal infection by initiating root canal treatment at 7 to 10 days.* In teeth with closed apices, revascularization cannot occur. These teeth should be endodontically treated within 7 to 10 days of the injury, before the ischemically necrosed pulp becomes infected.^{187,188} Endodontic treatment is extremely difficult so soon after a serious traumatic injury. It is beneficial to start the endodontic treatment with chemomechanical preparation, after which an intracanal dressing with a creamy mix of calcium hydroxide is placed (see Fig. 21.15).¹⁸⁷ Then, the clinician can fill the canal at his or her convenience after periodontal healing of the injury is complete, approximately 1 month after the instrumentation visit. There appears to be no necessity for long-term calcium hydroxide treatment in cases in which the endodontic treatment is started within 10 days of the injury.¹⁸⁷
2. *Eliminate pulp space infection.* When root canal treatment is initiated later than 10 days after the accident or if active external inflammatory resorption is observed, the preferred antibacterial protocol consists of microbial control followed by long-term dressing with calcium hydroxide.¹⁸⁸ Calcium hydroxide can effect an alkaline pH in the surrounding dentinal tubules (Fig. 21.36), kill bacteria, and neutralize endotoxin, a potent inflammatory stimulator.




Fig. 21.36 High pH of calcium hydroxide. The root was filled with calcium hydroxide and then cut in cross section. A pH indication shows the high pH in the canal and surrounding root, whereas the surrounding tissue is a neutral pH.

The first visit consists of the microbial control phase, with cleaning and shaping of the canal and the placement of either a corticosteroid/antibiotic material or a creamy mix of calcium hydroxide using a Lentulo spiral. The patient is seen in approximately 1 month, at which time the canal is filled with a dense mix of calcium hydroxide. Once filled, the canal should appear radiographically to be calcified because the radiodensity of calcium hydroxide in the canal is usually similar to that of the surrounding dentin (see Fig. 21.16). A radiograph is then exposed at 3-month intervals. At each visit the tooth is tested for symptoms of periodontitis. In addition to stopping the resorptive process, calcium hydroxide washout is assessed. Because the root surface is so radiodense as to make assessment of healing difficult, the adjacent bone healing is assessed. If adjacent bone has healed, it is assumed that the resorptive process has stopped in the root as well; then the canal can be obturated with the permanent root filling material (Fig. 21.37).

Internal Root Resorption

Internal root resorption is rare in permanent teeth. Internal resorption is characterized by an oval-shaped enlargement of the root canal space.²¹ External resorption, which is much more common, is often misdiagnosed as internal resorption.

Etiology. Internal root resorption is characterized by resorption of the internal aspect of the root by multinucleated giant cells adjacent to the granulation tissue in the pulp (Fig. 21.38). Chronic inflammatory tissue is common in the pulp, but only rarely does it result in resorption. There are different theories on the origin of the pulpal granulation tissue involved in internal resorption. The most logical explanation is that it is inflamed pulp tissue caused by an infected coronal pulp space. Communication between the coronal necrotic tissue and the vital pulp is through appropriately oriented dentinal tubules (see Fig. 21.38).¹⁹¹ One investigator¹⁹¹ reported that resorption of the dentin is frequently associated with deposition of hard tissue resembling bone or cementum and not dentin. He postulates that the resorbing tissue is not of pulpal origin but is “metaplastic” tissue derived from




Fig. 21.37 Healing of external inflammatory root resorption after calcium hydroxide treatment. The radiolucencies seen before treatment have disappeared with the reestablishment of the lamina dura. (Courtesy Dr. Fred Barnett.)




Fig. 21.38 Histologic appearance of internal root resorption. **A**, Section stained with Brown and Brenn. Bacteria are seen (in the dentinal tubules) communicating between the necrotic coronal segment and the apical granulomatous tissue and resorbing cells. **B**, An area of active internal root resorption. (Courtesy Dr. Leif Tronstad, Oslo, Norway.)

the pulpal invasion of macrophage-like cells. Others¹⁷¹ concluded that the pulp tissue was replaced by periodontium-like connective tissue when internal resorption was present. In addition to the requirement of the presence of granulation tissue, root resorption takes place only if the odontoblastic layer and predentin are lost or altered.¹⁸¹

Reasons for the loss of predentin adjacent to the granulation tissue are not obvious. Trauma frequently has been suggested as a cause.^{157,192} Another reason for the loss of predentin might be extreme heat produced when cutting on dentin without an adequate water spray. The heat presumably would destroy the predentin layer, and if later the coronal aspect of the pulp becomes infected, the bacterial products could initiate the typical inflammation in conjunction with resorbing giant cells in the vital pulp adjacent to the denuded root surface. Internal root resorption has been produced experimentally by the application of diathermy.¹⁹¹

Clinical Manifestations. Internal root resorption is usually asymptomatic and is first recognized clinically through

routine radiographs. For internal resorption to be active, at least part of the pulp must be vital. The coronal portion of the pulp is often necrotic, whereas the apical pulp that includes the internal resorptive defect can remain vital. Therefore a negative sensitivity test result does not rule out active internal resorption. It is also possible that the pulp becomes nonvital after a period of active resorption, giving a negative sensitivity test, radiographic signs of internal resorption, and radiographic signs of apical inflammation. Traditionally, the pink tooth has been thought to be pathognomonic of internal root resorption. The pink color is due to the granulation tissue in the coronal dentin undermining the crown enamel. The pink tooth can also be a feature of invasive cervical resorption, which must be excluded before a diagnosis of internal root resorption is made. Invasive cervical resorption has been described and classified by Heithersay^{92,93,94} by the extent of the resorptive lesion into the root, which initiates at the external cervical aspect of the tooth. Simple elimination of the resorptive tissue is ineffective in arresting its progress. Total removal or inactivation of the resorptive tissue is essential if recurrence (or concurrence) is to be avoided. Concurrence indicates the incomplete removal of the resorptive tissue at the time of treatment and recurrence is the reestablishment of the resorptive process. The reason for recurrence or concurrence is probably due to the invasive nature of the resorptive tissue whereby small infiltrative channels are created within the dentin, and these may interconnect with the periodontal ligament in positions more apical to the main resorptive defect. The use of trichloroacetic acid has been shown to be effective.⁹⁴

Radiographic Appearance. The usual radiographic presentation of internal root resorption is a fairly uniform radiolucent enlargement of the pulp canal (Figs. 21.39 and 21.40). Because the resorption is initiated in the root canal, the resorptive defect includes some part of the root canal space, so the original outline of the root canal is distorted.

Histologic Appearance. Like that of other inflammatory resorptive defects, the histologic picture of internal resorption



Fig. 21.39 Angled radiographs to show internal resorption. Radiographs from two different horizontal projections depict **(A)** the lesion within the confines of the canal on both views and **(B)** the adjacent bone intact on both views.



Fig. 21.40 A maxillary incisor with internal root resorption. Uniform enlargement of the pulp space is apparent. Outline of the canal cannot be seen in the resorptive defect.

is granulation tissue with multinucleated giant cells (see Fig. 21.37). An area of necrotic pulp is found coronal to the granulation tissue. Dentinal tubules containing microorganisms and communicating between the necrotic zone and the granulation tissue can sometimes be seen (see Fig. 21.38).^{162,181,188,191} Unlike external root resorption, the adjacent bone is not affected with internal root resorption.

Treatment. Treatment of internal root resorption is conceptually very easy. Because the resorptive defect is the result of the inflamed pulp and the blood supply to the tissue is through the apical foramina, endodontic treatment that effectively removes the blood supply to the resorbing cells is the treatment approach. After adequate anesthesia has been obtained, the canal apical to the internal defect is explored, and a working length short of the radiographic apex is used. The apical canal is thoroughly cleaned and shaped to ensure that the blood supply to the tissue resorbing the root is cut off.

By completion of the root canal instrumentation, it should be possible to obtain a blood-free and dry canal with paper points. Calcium hydroxide is spun into the canal to facilitate the removal of the tissue in the irregular defect at the next visit. At the second visit, the tooth and defect are filled with a warm gutta-percha technique (Fig. 21.41).




Fig. 21.41 **A**, Internal root resorption in a maxillary premolar with a history of trauma 7 years before the diagnosis (patient's head slammed against side window during an automobile accident). **B**, Three-year follow-up radiograph after endodontic treatment.

Diagnostic Features of External Versus Internal Root Resorption

It is often very difficult to distinguish external from internal root resorption, so misdiagnosis and incorrect treatment may result. The following sections present a list of typical diagnostic features of each resorptive type.

Radiographic Features. A change of angulation of x-rays should give a fairly good indication of whether a resorptive defect is internal or external. A lesion of internal origin appears close to the canal, whatever the angle of the x-ray (see Fig. 21.39). A defect on the external aspect of the root moves away from the canal as the angulation changes (Fig. 21.42). By using the buccal object rule, it is usually possible to distinguish whether the external root defect is buccal or lingual-palatal.

With internal resorption, the outline of the root canal is usually distorted, and the root canal and radiolucent resorptive defect appear contiguous (see Fig. 21.40). When the defect is external, the root canal outline appears normal




Fig. 21.42 External root resorption. Radiographs from two different horizontal projections depict movement of the lesion to outside the confines of the root canal.

and can usually be seen “running through” the radiolucent defect (Fig. 21.43).

Invasive cervical resorption may be accompanied by resorption of the bone in addition to the root (Fig. 21.44); radiolucencies can be apparent in the root and the adjacent bone. Internal root resorption does not involve the bone, and as a rule the radiolucency is confined to the root (see Fig. 21.39). On rare occasions, if the internal defect perforates the root, the bone adjacent to it is resorbed and appears radiolucent on the radiograph.

Vitality Testing. External inflammatory resorption in the apical and lateral aspects of the root involves an infected pulp space, so no response to sensitivity tests supports the diagnosis. However, because invasive cervical resorption does not involve the pulp, a normal response to sensitivity testing is usually associated with this type of resorption. Internal root resorption usually occurs in teeth with vital pulps and responds to sensitivity testing. In teeth that exhibit internal root resorption, it is common to register a nonresponse to sensitivity testing because often the coronal pulp has been removed or is necrotic, and the active resorbing cells are more apical in the canal. In addition, the pulp might have become necrotic after active resorption took place.

Pink Spot. With apical and lateral external root resorption, the pulp is nonvital, so the granulation tissue that



Fig. 21.44 Tooth #9 exhibiting invasive cervical resorption in a 31-year-old male. The tooth was injured 3 years prior after a fall from a scooter and was root filled and restored with a post crown. The resorption commences in the cervical external root and extends deeper into the root consistent with a Heithersay Grade III invasive cervical resorptive lesion. A resorptive defect is also present in the adjacent bone.




Fig. 21.43 **A**, Periapical radiograph of invasive cervical root resorption in tooth #8 and #9 in a 68-year-old female where the only known trauma was a fall onto bitumen aged 7 years when hit by a car. These teeth are diagnosed as having Heithersay Grade IV invasive cervical resorptive lesions which can be confirmed by CBCT imaging. **B** and **C**, CBCT coronal images. **D**, CBCT sagittal image of tooth #8. **E**, CBCT axial image. The outline of the canal is evident in Panels A, B, D, and E. The pulp does not appear exposed due to the anticlastic properties of predentin.

produces the pink spot is not present in these cases. For invasive cervical (Fig. 21.45) and internal root resorption, the pink spot due to the granulation tissue undermining the enamel is a possible sign.

Transient Apical Breakdown. A variation on internal resorption, noted especially after luxation injuries to more mature teeth, is transient apical internal resorption. Disruption of the neurovascular bundle at the apex may result in inflammatory mediators that allow osteoclastic changes at the apex to accommodate vascular derived repair followed by remodeling of the apex when the resorptive processes resolve. Teeth considered to show a combination of color changes, altered or V-notched apical radiographic changes, and altered pulp sensibility over time may have “transient apical breakdown” (TAB) for which no treatment other than observation is necessary.⁹

Summary of Possible Diagnostic Features

- External inflammatory root resorption due to pulp infection.
 - *Apical:* No response of the pulp to thermal or electric stimuli, with or without a history of trauma.
 - *Lateral:* History of trauma, no response of the pulp to thermal or electric stimuli, lesion moves on angled x-rays, root canal visualized radiographically overlying the defect, bony radiolucency also apparent.
- Invasive cervical resorption. Often a history of trauma (maybe forgotten, or its long-term risks not appreciated by the patient); positive pulp sensitivity test; lesion located at the attachment level of the tooth; lesion moves on angled x-rays; root canal outline is undistorted and can be visualized radiographically; crestal bony defect associated with the lesion; pink spot possible. CBCT imaging more reliably illustrates the extent of the resorption throughout the root (see Fig. 21.43).
- Internal root resorption. History of trauma, crown preparation, or pulpotomy; responsive pulp to thermal or electric stimuli likely, may occur at any location along the root canal (not only attachment level); lesion stays associated with the root canal on angled x-rays, radiolucency contained in the root without an adjacent bony defect; pink spot possible.




Fig. 21.45 Pink spot of invasive cervical root resorption. **A**, Radiographic appearance. **B**, Clinical appearance.

Most misdiagnoses of resorptive defects are made between invasive cervical and internal root resorptions. The diagnosis should always be confirmed as treatment proceeds. If root canal therapy is the treatment of choice for an apparent internal root resorption, the bleeding within the canal should cease quickly after pulp extirpation because the blood supply of the granulation tissue is the apical blood vessels. If bleeding continues during treatment, and particularly if it is still present at the second visit, the source of the blood supply is external, and treatment for perforating external resorption should be carried out. On obturation, it should be possible to fill the entire canal from within in internal resorption. Failure to achieve this should make the clinician suspicious of an external lesion that perforates the root. Finally, if the blood supply of an internal resorption defect is removed on pulp extirpation, any continuation of the resorptive process on recall radiographs should alert the clinician to the possibility that an external resorptive defect was misdiagnosed.

Clinical Management of the Avulsed Tooth

Favorable healing after an avulsion injury requires quick emergency intervention followed by evaluation and possible treatment at decisive times during the healing phase. The urgency of the emergency visit and the multidisciplinary nature of follow-up evaluations require that both the public and clinicians from many dental disciplines be knowledgeable about the treatment strategies involved.

CONSEQUENCES OF TOOTH AVULSION

Tooth avulsion results in attachment damage and pulp necrosis. The tooth is “separated” from the socket due mainly to tearing of the periodontal ligament that leaves viable periodontal ligament cells on most of the root surface. In addition, small and localized cemental damage occurs from the crushing of the tooth against the socket.

If the periodontal ligament left attached to the root surface does not dry out, the consequences of tooth avulsion are usually minimal.^{19,168} The hydrated periodontal ligament cells will maintain their viability and repair after replantation, with minimal destructive inflammation as a by-product. Because the areas of the crushing injury are localized, inflammation stimulated by the damaged tissues will be correspondingly limited, and favorable healing with new replacement cementum is likely to occur after the initial inflammation subsides (see Fig. 21.29).

If excessive drying occurs before replantation, the damaged periodontal ligament cells elicit a severe inflammatory response over a diffuse area on the root surface. Unlike the situation described earlier, in which the area to be repaired after the initial inflammatory response is small, here a large area of root surface is affected that must be repaired by new tissue. The slower-moving cementoblasts cannot cover the entire root surface in time, and it is likely that in certain areas, bone will attach directly onto the root surface. In time, through physiologic bone recontouring, the entire root will be replaced by bone. As earlier noted, this has been termed *replacement resorption* (see Figs. 21.30 and 21.31).^{25,184}

Pulpal necrosis always occurs after an avulsion injury. Although a necrotic pulp itself is not of consequence, the necrotic tissue is extremely susceptible to bacterial contamination. If revascularization does not occur or effective endodontic therapy is not carried out, the pulp space inevitably becomes infected. The combination of bacteria in the root canal and cemental damage on the external surface of the root results in an external inflammatory resorption that can be very serious and can lead to rapid loss of the tooth root (see Figs. 21.32 and 21.33).¹⁸¹

The consequences after tooth avulsion appear to be directly related to the severity and surface area of the inflammation on the root surface and resultant damaged root surface that must be repaired. Treatment strategies should always be considered in the context of limiting the extent of the periradicular inflammation, thus tipping the balance toward favorable responses (cemental) rather than unfavorable ones (replacement resorption or inflammatory resorption).

TREATMENT OBJECTIVES

Treatment is directed at avoiding or minimizing resultant inflammation due to the two main consequences of the avulsed tooth: attachment damage and pulpal infection.

Attachment damage as a direct result of the avulsion injury cannot be avoided. However, considerable additional damage can occur to the periodontal ligament in the time that the tooth is out of the mouth (primarily because of drying). Treatment is directed at minimizing this damage (and the resultant inflammation) so that the fewest possible complications result. When severe additional damage cannot be avoided and replacement resorption of the root is considered certain, steps are taken to slow the replacement of the root by bone to maintain the tooth in the mouth for as long as possible.

In the open apex tooth, all efforts are made to promote revascularization of the pulp, thus avoiding pulp space infection. When revascularization fails (in the open apex tooth) or is not possible (in the closed apex tooth), all treatment efforts are made to prevent or eliminate toxins from the root canal space.

CLINICAL MANAGEMENT

Emergency Treatment at the Accident Site

Replant if possible, or place in an appropriate storage medium. As mentioned, damage to the attachment apparatus that occurred during the initial injury is unavoidable but usually minimal. However, all efforts must be made to minimize necrosis of the remaining periodontal ligament while the tooth is out of the mouth. Pulpal sequelae are not a concern initially and are dealt with at a later stage of treatment.

The single most important factor to ensure a favorable outcome after replantation is the speed with which the tooth is replanted.^{21,27} Of utmost importance is the prevention of drying, which causes loss of normal physiologic metabolism and morphology of the periodontal ligament cells.^{27,168} Every effort should be made to replant the tooth within the first 15 to 20 minutes.³² This usually requires emergency personnel at the site of the injury with some knowledge of treatment protocol. The clinician should

communicate clearly with the person at the site of the accident. Ideally this information should have been given at an earlier time; for example, as an educational offering to school nurses or athletic trainers. Failing this, the information can be given over the phone. The aim is to replant a clean tooth with an undamaged root surface as gently as possible, after which the patient should be brought to the office immediately. If doubt exists that the tooth can be replanted adequately, the tooth should quickly be stored in an appropriate medium until the patient can get to the dental office for replantation. Suggested storage media, in order of preference, are milk, saliva (either in the vestibule of the mouth or in a container into which the patient expectorates), physiologic saline, and water.¹⁰⁰ Water is the least desirable storage medium because the hypotonic environment causes rapid cell lysis and increased inflammation on replantation.^{42,43}

Cell culture media in specialized transport containers, such as Hanks Balanced Salt Solution (HBSS), have shown superior ability to maintain the viability of the periodontal ligament fibers for extended periods.¹⁸⁶ Presently they are considered impractical because they need to be present at the accident site before the injury occurs. However, if we consider that more than 60% of avulsion injuries occur close to home or school,⁸⁹ it seems reasonable to assume that it would be beneficial to have these media available in emergency kits at these sites. It would also be advantageous to have them in ambulances and in the kits of emergency response personnel who are likely to treat the more serious injuries in which teeth might otherwise be sacrificed to a more serious life-threatening situation.

Management in the Dental Office

Emergency Visit. Prepare socket, prepare root, replant, construct a functional splint, and administer local and systemic antibiotics.

Recognizing that a dental injury might be secondary to a more serious injury is essential. The attending dental clinician is likely to be the first health care provider the patient sees after a head injury, so ruling out any injuries to the brain (e.g., concussion) and/or central nervous system (CNS) in general is paramount. If on examination a CNS injury is suspected, immediate referral to the appropriate expert is the first priority, above and beyond the dental injury. Once a CNS injury has been ruled out, the focus of the emergency visit is the attachment apparatus. The aim is to replant the tooth with a minimum of irreversibly damaged cells (that will cause inflammation) and the maximal number of periodontal ligament cells that have the potential to regenerate and repair the damaged root surface.

DIAGNOSIS AND TREATMENT PLANNING

If the tooth was replanted at the site of injury, a complete history is taken to assess the likelihood of a favorable outcome. The position of the replanted tooth is assessed and adjusted if necessary. On rare occasions, the tooth may be gently removed to prepare the root to increase the chances of a favorable outcome (discussed later).

If the patient presents with the tooth out of the mouth, the storage medium should be evaluated and the tooth placed in a more appropriate medium if required. HBSS is

presently considered the best medium for this purpose. Milk or physiologic saline is also appropriate for storage purposes.

The medical and accident histories are taken, and a clinical exam is carried out, with emphasis on questions about when, how, and where the injury occurred.

The clinical examination should include an examination of the socket to ascertain whether it is intact and suitable for replantation. The socket is gently rinsed with saline, and when it has been cleared of the clot and debris, its walls are examined directly for the presence, absence, or collapse of the socket wall. The socket and surrounding areas, including the soft tissues, should be radiographed. Three vertical angulations are required for diagnosis of the presence of a horizontal root fracture in adjacent teeth.²¹ The remaining teeth in both the upper and lower jaws should be examined for injuries, such as crown fractures. Any soft-tissue lacerations should be noted and, if tooth fragments are missing, explored. A photograph prior to any treatment is beneficial especially if legal concerns or insurance claims are involved.¹³

Preparation of the Root

Preparation of the root depends on the maturity of the tooth (open versus closed apex) and on the dry time of the tooth before it was placed in a storage medium. A dry time of 60 minutes is considered the point where survival of root periodontal ligament cells is unlikely.

EXTRAORAL DRY TIME LESS THAN 60 MINUTES

Closed Apex

The root should be rinsed of debris with water or saline and replanted in as gentle a fashion as possible.

If the tooth has a closed apex, revascularization is not possible,⁵⁹ but because the tooth was dry for less than 60 minutes (replanted or placed in appropriate medium), the chance for periodontal healing exists. Most importantly, the chance of a severe inflammatory response at the time of replantation is lessened. A dry time of less than 15 to 20 minutes is considered optimal, and periodontal healing would be expected.^{19,32,168}

A continuing challenge is the treatment of the tooth that has been dry for more than 20 minutes (periodontal cell survival is assured) but less than 60 minutes (periodontal cell survival unlikely). In these cases, logic suggests that the root surface consists of some cells with the potential to regenerate and some that will act as inflammatory stimulators.

Open Apex

Gently rinse off debris, if available, soak in doxycycline for 5 minutes or cover with minocycline, and replant.

In an open apex tooth, revascularization of the pulp and continued root development are possible (see Fig. 21.35). Investigators⁵⁹ found in monkeys that soaking the tooth in doxycycline (1 mg in approximately 20 mL of physiologic saline) for 5 minutes before replantation significantly enhanced complete revascularization (Fig. 21.46, A). This result was confirmed later in dogs by other investigators.^{150,197} A study found that covering the root with minocycline (Arestin, OraPharma, Warminster, PA), which attaches to the root




Fig. 21.46 **A**, Avulsed tooth soaking in doxycycline. **B**, Minocycline powder placed on the root surface before replantation.

for approximately 15 days, further increased the revascularization rate in dogs (see Fig. 21.46, B).¹⁵⁰ Although animal studies do not provide us with a prediction of the rate of revascularization in humans, it is reasonable to expect that the same enhancement of revascularization that occurred in two animal species also will occur in humans.³¹ As for a closed apex tooth, the open apex tooth is gently rinsed and replanted.

EXTRAORAL DRY TIME MORE THAN 60 MINUTES

Closed Apex

Remove the periodontal ligament by placing in acid for 5 minutes, soak in fluoride, and replant.

When the root has been dry for 60 minutes or more, survival of the periodontal ligament cells is not expected.^{27,168} In such cases the root should be prepared to be as resistant to resorption as possible (attempting to slow the osseous replacement process). These teeth should be soaked in acid for 5 minutes to remove all remaining periodontal ligament and thus remove the tissue that would initiate the inflammatory response on replantation. The tooth can then be soaked in 2% stannous fluoride for 5 minutes and replanted.^{41,159} A few years ago, studies were published that

indicated that an enamel matrix protein, Emdogain (Straumann USA, Andover, MA), could be beneficial in teeth with extended extraoral dry times, not only to make the root more resistant to resorption but also, possibly, to stimulate the formation of new periodontal ligament from the socket.^{74,104} Unfortunately, more recent studies have shown that the positive effect of Emdogain is only temporary, and most of these teeth start to resorb after a few years.¹⁵⁴

If the tooth has been dry for more than 60 minutes and no consideration is given to preserving the periodontal ligament, the endodontics may be performed extraorally. In the case of a tooth with a closed apex, no advantage exists to this additional step at the emergency visit. However, in a tooth with an open apex, endodontic treatment performed after replantation involves a long-term apexification procedure. In these cases, completing the root canal treatment extraorally, in which a seal in the blunderbuss apex is easier to achieve, may be advantageous. When endodontic treatment is performed extraorally, it must be performed aseptically with the utmost care to achieve a root canal system that is thoroughly disinfected.

Open Apex

Replant? If yes, treat as with closed apex tooth. Endodontic treatment may be performed out of the mouth.

Because these teeth are in young patients in whom facial development is usually incomplete, many pediatric clinicians consider the prognosis to be so poor and the potential complications of an ankylosed tooth so severe, they recommend that these teeth not be replanted. Considerable debate exists as to whether it would be beneficial to replant the root even though it will inevitably be lost due to osseous replacement. If the patients are followed carefully and the root submerged by decoronation procedure at the appropriate time,^{8,73,130} the height, and, more important, the width of the alveolar bone will be maintained, allowing for easier permanent restoration at the appropriate time when the child's facial development is complete.

Preparation of the Socket

The socket should be left undisturbed before replantation.²¹ Emphasis is placed on removal of obstacles in the socket to facilitate replacement of the tooth into the socket.⁸⁷ It should be lightly aspirated if a blood clot is present. If the alveolar bone has collapsed or may interfere with replantation, a blunt instrument should be inserted carefully into the socket in an attempt to reposition the wall.

Splinting

A splint that allows for physiologic movement of the tooth during healing and that is in place for a minimal period results in a decreased incidence of ankylosis.^{4,17,87} Flexible (physiologic) fixation for 1 to 2 weeks is recommended.^{4,6,7} Composite and wire splints where the wire diameter is no greater than 0.4 mm is considered to be a flexible splint.¹²⁴ The splint should allow movement of the tooth, should have no memory (so the tooth is not moved during healing), and should not impinge on the gingiva and/or prevent maintenance of oral



Fig. 21.47 Titanium trauma splint (TTS) in place.

hygiene in the area. Many splints satisfy the requirements of an acceptable device.¹¹¹ Fishing line has recently been recommended as a splinting material and, when bonded with resin modified glass ionomer cement, reduces the amount of iatrogenic damage to enamel when the splint is removed.^{75a,111} A new titanium trauma splint (TTS) has recently been shown to be particularly effective and easy to use (Fig. 21.47).¹⁹⁰ After the splint is in place, a radiograph should be exposed to verify the positioning of the tooth and as a preoperative reference for further treatment and follow-up. When the tooth is in the best possible position, adjusting the bite to ensure that it has not been splinted in a position causing traumatic occlusion is important. One week is sufficient to create periodontal support to maintain the avulsed tooth in position.¹⁸⁵ Therefore the splint should be removed after 1 to 2 weeks. The only exception is with avulsion in conjunction with alveolar fractures, for which 4 to 8 weeks is the suggested time of splinting.¹⁸⁵

Management of the Soft Tissues

Soft-tissue lacerations of the socket gingiva should be tightly sutured. Lacerations of the lip are fairly common with these types of injuries. The clinician should approach lip lacerations with some caution; a plastic surgery consult might be prudent. If these lacerations are sutured, care must be taken to clean the wound thoroughly beforehand because dirt or even minute tooth fragments left in the wound affect healing and the esthetic result.

Adjunctive Therapy

Systemic antibiotics given at the time of replantation and before endodontic treatment are effective in preventing bacterial invasion of the necrotic pulp and therefore subsequent inflammatory resorption.⁸⁷ Tetracycline has the additional benefit of decreasing root resorption by affecting the motility of the osteoclasts and reducing the effectiveness of collagenase.¹⁵¹ The administration of systemic antibiotics is recommended, beginning at the emergency visit and continuing until the splint is removed.⁸⁷ For patients not susceptible to tetracycline staining, the antibiotic of choice is doxycycline twice daily for 7 days at the appropriate dosage for patient age and weight.^{151,152} Penicillin V

1000 mg as a loading dose, followed by 500 mg 4 times daily for 7 days, has also been shown to be beneficial. The bacterial content of the sulcus also should be controlled during the healing phase. In addition to stressing to the patient the need for adequate oral hygiene, the use of chlorhexidine rinses for 7 to 10 days is helpful.

As stated previously, there is a benefit in removal of the pulp contents at the emergency visit and placing Ledermix or corticosteroid into the root canal.^{45,49} The medicament was able to shut down the inflammatory response after replantation and inhibit the spread of dentinoclasts to allow for more favorable healing compared with those teeth that did not have the medicament.^{144,145}

The need for analgesics should be assessed on an individual case basis. The use of pain medication stronger than nonprescription nonsteroidal antiinflammatory drugs (NSAIDs) is unusual. The patient should be sent to a physician for consultation regarding a tetanus booster within 48 hours of the initial visit.

SECOND VISIT

The second visit should take place 1 to 2 weeks after the trauma. At the emergency visit, emphasis was placed on the preservation and healing of the attachment apparatus. The focus of the second visit is the prevention or elimination of potential irritants from the root canal space. These irritants, if present, provide the stimulus for the progression of the inflammatory response, bone and root resorption. Also at this visit, the course of systemic antibiotics is completed; the chlorhexidine rinses can be stopped. At this appointment the splint is removed; the tooth might still have class I or class II mobility after splint removal, but all indications are that it will continue to heal better without the splint.⁴

Endodontic Treatment

EXTRAORAL TIME LESS THAN 60 MINUTES

Closed Apex

Initiate endodontic treatment after 1 to 2 weeks. When endodontic treatment is delayed or signs of resorption are present, provide long-term calcium hydroxide treatment before obturation.

No chance exists for revascularization of teeth with closed apices; therefore endodontic treatment should be initiated at the second visit 7 to 10 days later.^{18,59} If therapy is initiated at this optimum time, the pulp should be necrotic without (or with minimal) infection.^{129,187} Endodontic therapy with an effective interappointment antibacterial agent¹⁸⁷ over a relatively short period (1 to 2 weeks) is sufficient to ensure effective disinfection of the canal.¹⁶³ Long-term calcium hydroxide treatment should always be considered if the injury occurred more than 2 weeks before initiation of the endodontic treatment or especially if radiographic evidence of inflammatory resorption is present.¹⁸⁷

The root canal is thoroughly cleaned and shaped, irrigated, and then filled with calcium hydroxide. The canal is obturated when a radiographically intact periodontal membrane can be demonstrated around the root (see Fig. 21.37). Calcium hydroxide is an effective antibacterial agent^{46,163} and favorably

influences the local environment at the resorption site, theoretically promoting healing.¹⁸² It also changes the environment in the dentin to a more alkaline pH, which may slow the action of the resorptive cells and promote hard-tissue formation.¹⁸² However, changing of the calcium hydroxide should be kept to a minimum (not more than every 3 months) because it has a necrotizing effect on the cells attempting to repopulate the damaged root surface.¹²⁵

Calcium hydroxide is considered the material of choice in the prevention and treatment of inflammatory root resorption, but it is not the only medicament recommended in these cases. Some attempts have been made not only to remove the stimulus for the resorbing cells but also to affect them directly. The antibiotic-corticosteroid paste Ledermix is effective in treating inflammatory root resorption by inhibiting the spread of dentinoclasts^{144,145} without damaging the periodontal ligament; however, its ability to diffuse through the human tooth root has been demonstrated,² and its release and diffusion are enhanced when it is used in combination with calcium hydroxide paste.³

Open Apex

Avoid endodontic treatment and look for signs of revascularization. At the first sign of an infected pulp, initiate apexification procedure.

Teeth with open apices have the potential to revascularize and continue root development; initial treatment is directed toward reestablishing the blood supply.^{59,150,196} The initiation of endodontic treatment is avoided if at all possible, unless definite signs of pulp necrosis are present (e.g., periradicular inflammation). An accurate diagnosis of pulp vitality is extremely challenging in these cases. After trauma, a diagnosis of necrotic pulp is particularly undesirable because infection in these teeth is potentially more harmful due to cemental damage accompanying the traumatic injury. External inflammatory root resorption can be extremely rapid in these young teeth because the tubules are wide and allow irritants to move freely to the external surface of the root.^{59,196}

Patients should be recalled every 3 to 4 weeks for pulp vitality testing. Studies indicate that thermal tests with carbon dioxide snow (-78°C) or dichlorodifluoromethane (-40°C) placed at the incisal edge or pulp horn are the best methods of sensitivity testing, particularly in young permanent teeth.^{79,80,140} One of these two tests must be included in the pulp vitality testing. Reports confirm the superiority of the laser Doppler flowmeter in the diagnosis of revascularization of traumatized immature teeth¹⁹⁷; however, the cost of such an instrument precludes its use in the average dental office. Radiographic signs (apical breakdown and/or signs of lateral root resorption) and clinical signs (pain on percussion and palpation) of pulp pathosis are carefully assessed. At the first sign of pathosis, endodontic treatment should be initiated, and after disinfection of the root canal space, an apexification procedure should be carried out.

EXTRAORAL TIME MORE THAN 60 MINUTES

Closed Apex

Teeth with closed apices are treated endodontically in the same way as teeth that had an extraoral time of less than 60 minutes.

Open Apex (If Replanted)

If endodontic treatment was not performed out of the mouth, initiate apexification procedure.

The chance of revascularization in these teeth is extremely poor,^{184,188} so no attempt is made to revitalize them. An apexification procedure is initiated at the second visit if root canal treatment was not performed at the emergency visit. If endodontic treatment was performed at the emergency visit, the second visit is a recall visit to assess initial healing only.

Temporary Restoration

Effectively sealing the coronal access is essential to prevent infection of the canal between visits. Recommended temporary restorations are reinforced zinc oxide eugenol cement, acid etch composite resin, or glass ionomer cement. The depth of the temporary restoration is critical to its sealability. A depth of at least 4 mm is recommended, so a cotton pellet should not be placed; the temporary restoration is placed directly onto the calcium hydroxide in the access cavity. Calcium hydroxide should be removed from the walls of the access cavity before the temporary restoration is placed because calcium hydroxide is soluble and washes out when it comes into contact with saliva, leaving a defective temporary restoration.

After initiation of the root canal treatment, the splint is removed. If time does not permit complete removal of the splint at this visit, the resin tacks are smoothed so as not to irritate the soft tissues; the remaining resin is removed at a later appointment.

At this appointment, healing is usually sufficient to allow a detailed clinical examination of the teeth surrounding the avulsed tooth. Pulp vitality tests, reaction to percussion and palpation, and periodontal probing measurements should be carefully recorded for reference at follow-up visits.

ROOT FILLING VISIT

This visit should take place at the clinician's convenience or after long-term calcium hydroxide therapy, when an intact lamina dura is traced.

If the endodontic treatment was initiated 1 to 2 weeks after the avulsion and a thorough examination confirms normality, filling of the root canal at this visit is acceptable. Long-term use of calcium hydroxide is also a proven option for these cases. If endodontic treatment was initiated more than 2 weeks after the avulsion or active resorption is visible, the pulp space must first be disinfected before root filling. Traditionally, the reestablishment of a lamina dura (see Fig. 21.37) is a radiographic sign that the canal bacteria have been controlled. When an intact lamina dura can be traced throughout, root filling can take place.

The canal is cleaned, shaped, and irrigated under strict asepsis (i.e., a rubber dam). After completion of cleaning and shaping, the canal can be filled.

Permanent Restoration

Much evidence exists that coronal leakage caused by defective temporary and permanent restorations results in a

clinically relevant amount of bacterial contamination of the root canal after root filling.¹⁴⁹ Therefore the tooth should receive a permanent restoration as soon as possible. The depth of restoration is important for a tight seal, so the deepest restoration possible should be made. A post should be avoided if possible. Because most avulsions occur in the anterior region of the mouth where esthetics is important, composite resins combined with dentin bonding agents are recommended in these cases.

FOLLOW-UP CARE

Follow-up evaluations should take place at 3 months, 6 months, and yearly for at least 5 years. If replacement resorption is identified (see Fig. 21.31), a more closely monitored follow-up schedule is indicated. In the case of inflammatory root resorption (see Figs. 21.32 and 21.33), a new attempt at disinfection of the root canal space by standard retreatment might reverse the process. Teeth adjacent to and surrounding the avulsed tooth or teeth may show pathologic changes long after the initial accident, so these teeth, too, should be tested at follow-up visits and the results compared with those collected soon after the accident.

Late Complications

The IADT and AAE guidelines as stated previously recommend 5-year follow-up after the initial trauma. Heithersay^{95,96} wrote a series on the "life cycles of traumatized teeth" discussing follow-up of cases of up to 51 years post injury. Resorption is a common posttraumatic sequelae presenting many years after this initial trauma. Changes in the color of the crown also need to be monitored as either a need for bleaching of the tooth or a change in the condition of the tooth such as intracanal calcification. This chapter has presented many concepts and treatment approaches at the time of trauma. However, it is important to consider that there can be either unfortunate events over time that need secondary management or failure in diagnosis that is apparent years later when signs and symptoms of infection or resorption and inappropriate or poor treatment were performed at the time of injury (Figs. 21.48 and 21.49; see also Fig. 21.43).

Dentoalveolar Trauma to the Primary Dentition

The classification of injuries to the primary dentition is the same as the permanent dentition, although management may be different. Consideration must be given to the risk of damage to the developing permanent teeth. Furthermore, there may also be behavioral issues of the child that require different approaches in patient care. Instances where care is required to minimize risk to the permanent dentition include intrusive injuries when the intruded root is close to the developing permanent tooth. It is generally not advisable to replant an avulsed primary tooth. However, most luxation injuries heal spontaneously, so conservative treatment for preschool children is recommended. Immediate treatment is advised for complicated crown fractures, crown-root fractures, root fractures when the coronal fragment is displaced, extrusive luxations, and alveolar fractures.^{75,130}




Fig. 21.48 **A**, An occlusograph showing root fillings in teeth #9 and #10 in a 44-year-old male 25 years after the teeth had been traumatized. Symptoms and infection returned after endodontic treatment and surgical correction was required. **B**, CBCT imaging revealed internal resorption within the canal.




Fig. 21.49 **A**, An occlusograph showing extensive periapical radiolucencies associated with the maxillary incisor teeth in a 23-year-old male. The patient sustained a traumatic injury when he was 8 years of age. Tooth #8 has extruded gutta-percha through the open apex indicating either a failed apexification procedure or poor obturation of the canal due to the open apex. Tooth #9 has an open apex indication pulp necrosis likely occurred at or soon after the traumatic injury but no treatment was undertaken. **B**, CBCT imaging revealing the immature root formation of tooth #9 and periapical lesion. Surface resorption on the palatal cervical root is suggestive of a prior palatal luxation injury.

References

1. AAE clinical considerations for a regenerative procedure. <https://www.aae.org/specialty/wp-content/uploads/sites/2/2017/06/currentregenerativeendodonticconsiderations.pdf>.
2. Abbott PV, Heithersay GS, Hume WR: Release and diffusion through human tooth roots in vitro of corticosteroid and tetracycline trace molecules from Ledermix paste, *Endod Dent Traumatol* 4:55, 1988.
3. Abbott PV, Hume WR, Heithersay GS: Effects of combining Ledermix and calcium hydroxide pastes on the diffusion of corticosteroid and tetracycline through human tooth roots in vitro, *Endod Dent Traumatol* 5:188, 1989.
4. About I, Murray PE, Franquin JC, et al: The effect of cavity restoration variables on odontoblast cell numbers and dental repair, *J Dent* 29:109, 2001.
5. Accorinte L, Holland R, Reis A, et al: Evaluation of mineral trioxide aggregate and calcium hydroxide cement as pulp-capping agents in human teeth, *J Endod* 34:1, 2008.
6. Deleted in page review.
7. Andersson L, Friskopp J, Blomlof L: Fiber-glass splinting of traumatized teeth, *ASDC J Dent Child* 50:21, 1983.
8. Andersson L, Malmgren B: The problem of dentoalveolar ankylosis and subsequent replacement resorption in the growing patient, *Aust Endod J* 25:57, 1999.
9. Andreasen FM: Transient apical breakdown and its relation to color and sensibility changes after luxation injuries to teeth, *Endod Dent Traumatol* 2:9, 1986.
10. Andreasen FM: Pulpal healing after luxation injuries and root fracture in the permanent dentition, *Endod Dent Traumatol* 5:111, 1989.

11. Andreasen FM, Andreasen JO: Resorption and mineralization processes following root fracture of permanent incisors, *Endod Dent Traumatol* 4:202, 1988.
12. Andreasen FM, Andreasen JO, Bayer T: Prognosis of root-fractured permanent incisors: prediction of healing modalities, *Endod Dent Traumatol* 5:11, 1989.
13. Andreasen FM, Kahler B: Diagnosis of acute dental trauma: the importance of standardized documentation: a review, *Dent Traumatol* 31:340, 2015.
14. Andreasen FM, Kahler B: Pulpal response after acute dental injury in the permanent dentition: clinical implications-a review, *J Endod* 41:299, 2015.
15. Andreasen FM, Pedersen BV: Prognosis of luxated permanent teeth: the development of pulp necrosis, *Endod Dent Traumatol* 1:207, 1985.
16. Andreasen FM, Zhijie Y, Thomsen BL, et al: Occurrence of pulp canal obliteration after luxation injuries in the permanent dentition, *Endod Dent Traumatol* 3:103, 1987.
17. Andreasen JO: Etiology and pathogenesis of traumatic dental injuries: a clinical study of 1,298 cases, *Scand J Dent Res* 78:329, 1970.
18. Andreasen JO: Luxation of permanent teeth due to trauma: a clinical and radiographic follow-up study of 189 injured teeth, *Scand J Dent Res* 78:273, 1970.
19. Andreasen JO: Effect of extra-alveolar period and storage media upon periodontal and pulpal healing after replantation of mature permanent incisors in monkeys, *Int J Oral Surg* 10:43, 1981.
20. Andreasen JO: Review of root resorption systems and models: etiology of root resorption and the homeostatic mechanisms of the periodontal ligament. In Davidovitch Z, editor: *The biological mechanisms of tooth eruption and resorption*, Birmingham, AL, 1989, EB-SCOP Media.
21. Andreasen JO, Andreasen FM, Andersson L: *Textbook and color atlas of traumatic injuries to the teeth*, ed 4, Hoboken, NJ, 2007, Wiley-Blackwell.
22. Andreasen JO, Andreasen FM, Mejare I, et al: Healing of 400 intra-alveolar root fractures. Part 1. Effect of pre-injury and injury factors such as sex, age, stage of root development, fracture type, location of fracture and severity of dislocation, *Dent Traumatol* 20:192, 2004.
23. Andreasen JO, Andreasen FM, Mejare I, et al: Healing of 400 intra-alveolar root fractures. Part 2. Effect of treatment factors such as treatment delay, repositioning, splinting type and period and antibiotics, *Dent Traumatol* 20:203, 2004.
24. Andreasen JO, Farik B, Munksgaard EC: Long-term calcium hydroxide as a root canal dressing may increase risk of root fracture, *Dent Traumatol* 18:134, 2002.
25. Andreasen JO, Hjorting-Hansen E: Replantation of teeth. I. Radiographic and clinical study of 110 human teeth replanted after accidental loss, *Acta Odontol Scand* 24:263, 1966.
26. Andreasen JO, Hjorting-Hansen E: Intraalveolar root fractures: radiographic and histologic study of 50 cases, *J Oral Surg* 25:414, 1967.
27. Andreasen JO, Kristerson L: The effect of limited drying or removal of the periodontal ligament: periodontal healing after replantation of mature permanent incisors in monkeys, *Acta Odontol Scand* 39:1, 1981.
28. Andreasen JO, Ravn JJ: Epidemiology of traumatic dental injuries to primary and permanent teeth in a Danish population sample, *Int J Oral Surg* 1:235, 1972.
29. Arakawa M, Kitasako Y, Otsuki M, et al: Direct pulp capping with an auto-cured sealant resin and a self-etching primer, *Am J Dent* 16:61, 2003.
30. Banchs F, Trope M: Revascularization of immature permanent teeth with apical periodontitis: new treatment protocol?, *J Endod* 30:196, 2004.
31. Barrett AP, Reade PC: Revascularization of mouse tooth isografts and allografts using autoradiography and carbon-perfusion, *Arch Oral Biol* 26:541, 1981.
32. Barrett EJ, Kenny DJ: Avulsed permanent teeth: a review of the literature and treatment guidelines, *Endod Dent Traumatol* 13:153, 1997.
33. Bastone EB, Freer TJ, McNamara JR: Epidemiology of dental trauma: a review of the literature, *Aust Dent J* 45:2, 2000.
34. Becerra P, Ricucci D, Loghin S, et al: Histological study of a human immature permanent premolar with chronic apical abscess after revascularization/revitalization, *J Endod* 40:133, 2014.
35. Belobrov I, Parashos P: Treatment of tooth discoloration after the use of white mineral trioxide aggregate, *J Endod* 37:1017, 2011.
36. Bergenholtz G: Micro-organisms from necrotic pulp of traumatized teeth, *Odontol Revy* 25:347, 1974.
37. Bernardes RA, de Moraes IG, Duarte MAH, et al: Use of cone-beam volumetric tomography in the diagnosis of root fractures, *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 108:270, 2009.
38. Bernardes RA, de Paulo RS, Pereira LO, et al: Comparative study of cone beam computed tomography and intraoral periapical radiographs in diagnosis of lingual-simulated external root resorptions, *Dent Traumatol* 28:268, 2012.
39. Bhaskar SN, Rappaport HM: Dental vitality tests and pulp status, *J Am Dent Assoc* 86:409, 1973.
40. Binnie WH, Rowe AH: A histological study of the periapical tissues of incompletely formed pulpless teeth filled with calcium hydroxide, *J Dent Res* 52:1110, 1973.
41. Bjorvatn K, Selvig KA, Klinge B: Effect of tetracycline and SnF₂ on root resorption in replanted incisors in dogs, *Scand J Dent Res* 97:477, 1989.
42. Blomlof L: Milk and saliva as possible storage media for traumatically exarticulated teeth prior to replantation, *Swed Dent J Suppl* 8:1, 1981.
43. Blomlof L, Lindskog S, Andersson L, et al: Storage of experimentally avulsed teeth in milk prior to replantation, *J Dent Res* 62:912, 1983.
44. Bose R, Nummikoski P, Hargreaves K: A retrospective evaluation of radiographic outcomes in immature teeth with necrotic root canal systems treated with regenerative endodontic procedures, *J Endod* 35:1343, 2009.
- 44a. Bourguignon C, Cohenca N, Lauridsen E, et al: International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: 1. Fractures and luxations, *Dent Traumatol*, 2020. doi:10.1111/edt.12578.
45. Bryson EC, Levin L, Banchs F, et al: Effect of immediate intracanal placement of Ledermix Paste on healing of replanted dog teeth after extended dry times, *Dent Traumatol* 18:316, 2002.
46. Bystrom A, Claesson R, Sundqvist G: The antibacterial effect of camphorated paramonochlorophenol, camphorated phenol and calcium hydroxide in the treatment of infected root canals, *Endod Dent Traumatol* 1:170, 1985.
47. Canakci V, Akgul HM, Akgul N, et al: Prevalence and handedness correlates of traumatic injuries to the permanent incisors in 13 17-year-old adolescents in Erzurum, Turkey, *Dent Traumatol* 19:248, 2003.
48. Chala S, Abouqal R, Rida S: Apexification of immature teeth with calcium hydroxide or mineral trioxide aggregate: systematic review and meta-analysis, *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 112:e36, 2011.
49. Chen H, Teixeira FB, Ritter AL, et al: The effect of intracanal anti-inflammatory medicaments on external root resorption of replanted dog teeth after extended extra-oral dry time, *Dental Traumatol* 24:74, 2008.
50. Cohenca N, Simon JH, Marhtur A, et al: Clinical indications for digital imaging in dento-alveolar trauma. Part 2. Root resorption, *Dent Traumatol* 23:105, 2007.
51. Cohenca N, Simon JH, Roges R, et al: Clinical indications for digital imaging in dento-alveolar trauma. Part 1. Traumatic injuries, *Dent Traumatol* 23:95, 2007.
52. Cotton TP, Geisler TM, Holden DT, et al: Endodontic applications of cone beam volumetric tomography, *J Endod* 33:1121, 2007.
53. Cox CF, Keall CL, Keall HJ, et al: Biocompatibility of surface-sealed dental materials against exposed pulps, *J Prosthet Dent* 57:1, 1987.
54. Cox CF, Tarim B, Kopel H, et al: Technique sensitivity: biological factors contributing to clinical success with various restorative materials, *Adv Dent Res* 15:85, 2001.
55. Cunha RF, Pavarini A, Percinoto C, et al: Influence of surgical repositioning of mature permanent dog teeth following experimental intrusion: a histologic assessment, *Dent Traumatol* 18:304, 2002.
56. Cvek M: Treatment of non-vital permanent incisors with calcium hydroxide. IV. Periodontal healing and closure of the root canal in the coronal fragment of teeth with intra-alveolar fracture and vital apical fragment: a follow-up, *Odontol Revy* 25:239, 1974.
57. Cvek M: A clinical report on partial pulpotomy and capping with calcium hydroxide in permanent incisors with complicated crown fracture, *J Endod* 4:232, 1978.
58. Cvek M: Prognosis of luxated non-vital maxillary incisors treated with calcium hydroxide and filled with gutta-percha: a retrospective clinical study, *Endod Dent Traumatol* 8:45, 1992.

59. Cvek M, Cleaton-Jones P, Austin J, et al: Effect of topical application of doxycycline on pulp revascularization and periodontal healing in reimplanted monkey incisors, *Endod Dent Traumatol* 6:170, 1990.
60. Cvek M, Cleaton-Jones PE, Austin JC, et al: Pulp reactions to exposure after experimental crown fractures or grinding in adult monkeys, *J Endod* 8:391, 1982.
61. Cvek M, Hollender L, Nord CE: Treatment of non-vital permanent incisors with calcium hydroxide. VI. A clinical, microbiological and radiological evaluation of treatment in one sitting of teeth with mature or immature root, *Odontol Revy* 27:93, 1976.
62. Cvek M, Nord CE, Hollender L: Antimicrobial effect of root canal debridement in teeth with immature root: a clinical and microbiologic study, *Odontol Revy* 27:1, 1976.
63. Da Silva AC, Passeri LA, Mazzonetto R, et al: Incidence of dental trauma associated with facial trauma in Brazil: a 1-year evaluation, *Dent Traumatol* 20:6, 2004.
64. Day P, Flores MT, O'Connell A, et al: International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: 3. Injuries in the primary dentition, *Dent Traumatol*, 2020, doi:10.1111/edt.12576.
65. Deutsch AS, Musikant BL, Cavallari J, et al: Root fracture during insertion of prefabricated posts related to root size, *J Prosthet Dent* 53:786, 1985.
66. Durack C, Patel S, Davies J, et al: Diagnostic accuracy of small volume cone beam computed tomography and intraoral periapical radiography for the detection of simulated external inflammatory root resorption, *Int Endod J* 44:136, 2011.
67. Ebeleseder KA, Santler G, Glockner K, et al: An analysis of 58 traumatically intruded and surgically extruded permanent teeth, *Endod Dent Traumatol* 16:34, 2000.
68. Eklund G, Stalhane I, Hedegard B: A study of traumatized permanent teeth in children aged 7-15 years. III. A multivariate analysis of post-traumatic complications of subluxated and luxated teeth, *Sven Tandlak Tidsskr* 69:179, 1976.
69. Endodontics Colleagues for Excellence. Summer 2014. The treatment of traumatic dental injuries. https://www.aae.org/specialty/wp-content/uploads/sites/2/2017/06/ecfe_summer2014-final.pdf
70. Ersin NK, Eronat N: The comparison of a dentin adhesive with calcium hydroxide as a pulp-capping agent on the exposed pulps of human and sheep teeth, *Quintessence Int* 36:271, 2005.
71. Estrela C, Reis Bueno M, Alencar AHG: Method to evaluate inflammatory root resorption by using cone beam computed tomography, *J Endod* 35:1491, 2009.
72. Faraco IM Jr, Holland R: Response of the pulp of dogs to capping with mineral trioxide aggregate or a calcium hydroxide cement, *Dent Traumatol* 17:163, 2001.
73. Filippi A, Pohl Y, von Arx T: Decoronation of an ankylosed tooth for preservation of alveolar bone prior to implant placement, *Dent Traumatol* 17:93, 2001.
74. Filippi A, Pohl Y, von Arx T: Treatment of replacement resorption with Emdogain: preliminary results after 10 months, *Dent Traumatol* 17:134, 2001.
75. Flores MT: Traumatic injuries in the primary dentition, *Dent Traumatol* 18:287, 2002.
- 75a. Fouad AE, Abbott PV, Tsilingaridis G, et al: International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: 2. Avulsion of permanent teeth, *Dent Traumatol*, 2020, doi:10.1111/edt.12573.
76. Frank AL: Therapy for the divergent pulpless tooth by continued apical formation, *J Am Dent Assoc* 72:87, 1966.
77. Fuks AB, Bielak S, Chosak A: Clinical and radiographic assessment of direct pulp capping and pulpotomy in young permanent teeth, *Pediatr Dent* 4:240, 1982.
78. Fuks AB, Cosack A, Klein H, et al: Partial pulpotomy as a treatment alternative for exposed pulps in crown-fractured permanent incisors, *Endod Dent Traumatol* 3:100, 1987.
79. Fulling HJ, Andreasen JO: Influence of maturation status and tooth type of permanent teeth upon electrometric and thermal pulp testing, *Scand J Dent Res* 84:286, 1976.
80. Fuss Z, Trowbridge H, Bender IB, et al: Assessment of reliability of electrical and thermal pulp testing agents, *J Endod* 12:301, 1986.
81. Gazelius B, Olgart L, Edwall B: Restored vitality in luxated teeth assessed by laser Doppler flowmeter, *Endod Dent Traumatol* 4:265, 1988.
82. Gelbier MJ, Winter GB: Traumatized incisors treated by vital pulpotomy: a retrospective study, *Br Dent J* 164:319, 1988.
83. Giuliani V, Baccetti T, Pace R, et al: The use of MTA in teeth with necrotic pulps and open apices, *Dent Traumatol* 18:217, 2002.
84. Goldberg F, Kaplan A, Roitman M, et al: Reinforcing effect of a resin glass ionomer in the restoration of immature roots in vitro, *Dent Traumatol* 18:70, 2002.
85. Granath LE, Hagman G: Experimental pulpotomy in human bicusps with reference cutting technique, *Acta Odontol Scand* 29:155, 1971.
86. Hammarstrom L, Lindsog S: General morphological aspects of resorption of teeth and alveolar bone, *Int Endod J* 18:93, 1985.
87. Hammarstrom L, Pierce A, Blomlof L, et al: Tooth avulsion and replantation: a review, *Endod Dent Traumatol* 2:1, 1986.
88. Hebling J, Giro EM, Costa CA: Biocompatibility of an adhesive system applied to exposed human dental pulp, *J Endodontol* 25:676, 1999.
89. Hedegard B, Stalhane I: A study of traumatized permanent teeth in children 7-15 years. I, *Sven Tandlak Tidsskr* 66:431, 1973.
90. Heide S, Mjor IA: Pulp reactions to experimental exposures in young permanent monkey teeth, *Int Endod J* 16:11, 1983.
91. Heithersay GS: Calcium hydroxide in the treatment of pulpless teeth with associated pathology, *J Br Endod Soc* 8:74, 1975.
92. Heithersay GS: Clinical, radiologic, and histopathologic features of invasive cervical resorption, *Quintessence Int* 30:27, 1999.
93. Heithersay GS: Invasive cervical resorption: an analysis of potential predisposing factors, *Quintessence Int* 30:83, 1999.
94. Heithersay GS: Treatment of invasive cervical resorption: an analysis of results using topical application of trichloroacetic acid, curettage, and restoration, *Quintessence Int* 30:96, 1999.
95. Heithersay GS: Life cycles of traumatized teeth: long-term observations from a cohort of dental trauma victims, *Aust Dent J* 61 Suppl 1: 120, 2016.
96. Heithersay GS: Life cycles of traumatized teeth: long-term observations from a cohort of dental trauma victims-series 2, *Aust Dent J* 61(Suppl 1):317, 2016.
97. Heithersay GS, Kahler B: Healing responses following transverse root fracture: a historical review and case reports showing healing with (a) calcified tissue and (b) dense fibrous connective tissue, *Dent Traumatol* 29:253, 2013.
98. Heller AL, Koenigs JF, Brilliant JD, et al: Direct pulp capping of permanent teeth in primates using a resorbable form of tricalcium phosphate ceramic, *J Endod* 1:95, 1975.
99. Herforth A, Strassburg M: Therapy of chronic apical periodontitis in traumatically injuring front teeth with ongoing root growth, *Dtsch Zahnarztl Z* 32:453, 1977.
100. Hiltz J, Trope M: Vitality of human lip fibroblasts in milk, Hanks Balanced Salt Solution and Viaspan storage media, *Endod Dent Traumatol* 7:69, 1991.
101. Holland R, Bisco Ferreira L, de Souza V, et al: Reaction of the lateral periodontium of dogs' teeth to contaminated and noncontaminated perforations filled with mineral trioxide aggregate, *J Endod* 33:1192, 2007.
102. Holloway GA Jr, Watkins DW: Laser Doppler measurement of cutaneous blood flow, *J Invest Dermatol* 69:306, 1977.
103. Hoshino E, Kurihara-Ando N, Sato I, et al: In-vitro antibacterial susceptibility of bacteria taken from infected root dentine to a mixture of ciprofloxacin, metronidazole and minocycline, *Int Endod J* 29:125, 1996.
104. Iqbal MK, Bamaas N: Effect of enamel matrix derivative (Emdogain) upon periodontal healing after replantation of permanent incisors in beagle dogs, *Dent Traumatol* 17:36, 2001.
105. Iwaya S, Ikawa M, Kubota M: Revascularization of an immature permanent tooth with apical periodontitis and sinus tract, *Dental Traumatol* 17:185, 2001.
106. Jacobsen I: Root fractures in permanent anterior teeth with incomplete root formation, *Scand J Dent Res* 84:210, 1976.
107. Jacobsen I, Kerekes K: Diagnosis and treatment of pulp necrosis in permanent anterior teeth with root fracture, *Scand J Dent Res* 88:370, 1980.
108. Jacobsen I, Zachrisson BU: Repair characteristics of root fractures in permanent anterior teeth, *Scand J Dent Res* 83:355, 1975.
109. Jarvinen S: Fractured and avulsed permanent incisors in Finnish children: a retrospective study, *Acta Odontol Scand* 37:47, 1979.
110. Kahler B, Heithersay GS: An evidence-based appraisal of splinting luxated, avulsed and root-fractured teeth, *Dent Traumatol* 24:2, 2008.

111. Kahler B, Hu JY, Marriot-Smith CS, et al: Splinting of teeth following trauma: a review and a new splinting recommendation, *Aust Dent J* 61(Suppl 1):59, 2016.
112. Kahler B, Mistry S, Moule A, et al: Revascularization outcomes: a prospective analysis of 16 consecutive cases, *J Endod* 40:333, 2014.
113. Kahler B, Rossi-Fedele G: A review of tooth discoloration after regenerative endodontic therapy, *J Endod* 42:563, 2016.
114. Kahler B, Rossi-Fedele G, Chugal N, et al: An evidence-based review of the efficacy of treatment approaches for immature permanent teeth with pulp necrosis, *J Endod* 43:1052, 2017.
115. Kahler SL, Shetty S, Andreasen FM, et al: The effect of long-term dressing with calcium hydroxide on the fracture susceptibility of teeth, *J Endod* 44:464, 2018.
116. Kakehashi S, Stanley HR, Fitzgerald RJ: The effect of surgical exposures on dental pulps in germ-free and conventional laboratory rats, *Oral Surg* 20:340, 1965.
117. Kamburoglu K, Ilker Cebeci AR, Göran Gröndahl H: Effectiveness of limited cone-beam computed tomography in the detection of horizontal root fracture, *Dent Traumatol* 25:256, 2009.
118. Katebzadeh N, Dalton BC, Trope M: Strengthening immature teeth during and after apexification, *J Endod* 24:256, 1998.
119. Kenny DJ, Barrett EJ, Casas MJ: Avulsions and intrusions: the controversial displacement injuries, *J Can Dent Assoc* 69:308, 2003.
120. Kerekes K, Heide S, Jacobsen I: Follow-up examination of endodontic treatment in traumatized juvenile incisors, *J Endod* 6:744, 1980.
121. Kim SG, Malek M, Sigurdsson A, et al: Regenerative endodontics: a comprehensive review, *Int Endod J* 51:1367, 2018.
122. Koliniotou-Koumpia E, Tziakas D: Pulpal responses following direct pulp capping of healthy dog teeth with dentine adhesive systems, *J Dent* 33:639, 2005.
123. Kositbowornchai S, Nuansakul R, Sikram S, et al: Root fracture detection: a comparison of direct digital radiography with conventional radiography, *Dentomaxillofac Radiol* 30:106, 2001.
124. Kwan SC, Johnson JD, Cohen N: The effect of splint material and thickness on tooth mobility after extraction and replantation using a human cadaveric model, *Dent Traumatol* 28:277, 2012.
125. Lengheden A, Blomlof L, Lindsog S: Effect of delayed calcium hydroxide treatment on periodontal healing in contaminated replanted teeth, *Scand J Dent Res* 99:147, 1991.
126. Lindsog S, Pierce AM, Blomlof L, et al: The role of the necrotic periodontal membrane in cementum resorption and ankylosis, *Endod Dent Traumatol* 1:96, 1985.
127. Love RM: Bacterial penetration of the root canal of intact incisor teeth after a simulated traumatic injury, *Endod Dent Traumatol* 12:289, 1996.
128. Love RM, Ponnambalam Y: Dental and maxillofacial skeletal injuries seen at the University of Otago School of Clinicianry, New Zealand, 2000-2004, *Dent Traumatol* 24:170, 2008.
129. Lundin SA, Noren JG, Warfvinge J: Marginal bacterial leakage and pulp reactions in Class II composite resin restorations in vivo, *Swed Dent J* 14:185, 1990.
130. Malmgren B, Andreasen JO, Flores MT, et al: International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: 3. Injuries in the primary dentition, *Dent Traumatol* 28:174, 2012.
131. Maroto M, Barberia E, Planells P, et al: Treatment of a non-vital immature incisor with mineral trioxide aggregate (MTA), *Dent Traumatol* 19:165, 2003.
132. Martin G, Ricucci D, Gibbs JL, et al: Histological findings of revascularized/revitalized immature permanent molar with apical periodontitis using platelet-rich plasma, *J Endod* 39:138, 2013.
133. Masterton JB: The healing of wounds of the dental pulp: an investigation of the nature of the scar tissue and of the phenomena leading to its formation, *Dent Pract Dent Rec* 16:325, 1966.
134. May JJ, Cohen N, Peters OA: Contemporary management of horizontal root fractures to the permanent dentition: diagnosis, radiologic assessment to include cone-beam computed tomography, *Pediatr Dent* 35:120, 2013.
135. Mejare I, Hasselgren G, Hammarstrom LE: Effect of formaldehyde-containing drugs on human dental pulp evaluated by enzyme histochemical technique, *Scand J Dent Res* 84:29, 1976.
136. Mesaros S, Trope M, Maixner W, et al: Comparison of two laser Doppler systems on the measurement of blood flow of premolar teeth under different pulpal conditions, *Int Endod J* 30:167, 1997.
137. Murray PE, Smith AJ, Windsor LJ, et al: Remaining dentine thickness and human pulp responses, *Int Endod J* 33:36, 2003.
138. Myers WC, Fountain SB: Dental pulp regeneration aided by blood and blood substitutes after experimentally induced periapical infection, *Oral Surg Oral Med Oral Pathol* 37:441, 1974.
139. Nair PN, Duncan HF, Pitt Ford TR, et al: Histological, ultrastructural and quantitative investigations on the response of healthy human pulps to experimental capping with mineral trioxide aggregate: a randomized controlled trial, *Int Endod J* 41:128, 2008.
140. Ohman A: Healing and sensitivity to pain in young replanted human teeth: an experimental, clinical and histological study, *Odontol Tidskr* 73:166, 1965.
141. Parirokh M, Kakoei S: Vital pulp therapy of mandibular incisors: a case report with 11-year follow up, *Aust Endod J* 32:75, 2006.
142. Patel S, Dawood A, Pitt Ford T, et al: The potential applications of cone beam computed tomography in the management of endodontic problems, *Int Endod J* 40:818, 2007.
143. Patel S, Durack C, Abella F, et al: Cone beam computed tomography in endodontics: a review, *Int Endod J* 48:3, 2015.
144. Pierce A, Heithersay G, Lindsog S: Evidence for direct inhibition of dentinoclasts by a corticosteroid/antibiotic endodontic paste, *Endod Dent Traumatol* 4:44, 1988.
145. Pierce A, Lindsog S: The effect of an antibiotic/corticosteroid paste on inflammatory root resorption in vivo, *Oral Surg Oral Med Oral Pathol* 64:216, 1987.
146. Pileggi R, Dumsha TC, Myslinksi NR: The reliability of electric pulp test after concussion injury, *Endod Dent Traumatol* 12:16, 1996.
147. Rabie G, Barnett F, Tronstad L: Long-term splinting of maxillary incisor with intra alveolar root fracture, *Endod Dent Traumatol* 4:99, 1988.
148. Ravn JJ: Follow-up study of permanent incisors with complicated crown fractures after acute trauma, *Scand J Dent Res* 90:363, 1982.
149. Ray HA, Trope M: Periapical status of endodontically treated teeth in relation to the technical quality of the root filling and the coronal restoration, *Int Endod J* 28:12, 1995.
150. Ritter AL, Ritter AV, Murrah V, et al: Pulp revascularization of replanted immature dog teeth after treatment with minocycline and doxycycline assessed by laser Doppler flowmetry, radiography, and histology, *Dent Traumatol* 20:75, 2004.
151. Sae-Lim V, Wang CY, Choi GW, et al: The effect of systemic tetracycline on resorption of dried replanted dogs' teeth, *Endod Dent Traumatol* 14:127, 1998.
152. Sae-Lim V, Wang CY, Trope M: Effect of systemic tetracycline and amoxicillin on inflammatory root resorption of replanted dogs' teeth, *Endod Dent Traumatol* 14:216, 1998.
153. Saroglu I, Sonmez H: The prevalence of traumatic injuries treated in the pedodontic clinic of Ankara University, Turkey, during 18 months, *Dent Traumatol* 18:299, 2002.
154. Schjott M, Andreasen JO: Emdogain does not prevent progressive root resorption after replantation of avulsed teeth: a clinical study, *Dental Traumatol* 21:46, 2005.
155. Schroder U: Reaction of human dental pulp to experimental pulpotomy and capping with calcium hydroxide (thesis), *Odont Revy* 24(Suppl 25):97, 1973.
156. Schroder U, Granath LE: Early reaction of intact human teeth to calcium hydroxide following experimental pulpotomy and its significance to the development of hard tissue barrier, *Odontol Revy* 22:379, 1971.
157. Seltzer S: *Endodontology*, Philadelphia, 1988, Lea & Febiger.
158. Seltzer S, Bender IB, Turkenkopf S: Factors affecting successful repair after root canal therapy, *J Am Dent Assoc* 52:651, 1963.
159. Selvig KA, Zander HA: Chemical analysis and microradiography of cementum and dentin from periodontically diseased human teeth, *J Periodontol* 33:103, 1962.
160. Shuping GB, Orstavik D, Sigurdsson A, et al: Reduction of intracanal bacteria using nickel-titanium rotary instrumentation and various medications, *J Endod* 26:751, 2000.
161. Silva GA, Lanza LD, Lopes-Junior N, et al: Direct pulp capping with a dentin bonding system in human teeth: a clinical and histological evaluation, *Oper Dent* 31:297, 2006.
162. Silverman S: The dental structures in primary hyperparathyroidism, *Oral Surg* 15:426, 1962.
163. Sjogren U, Figdor D, Spangberg L, et al: The antimicrobial effect of calcium hydroxide as a short-term intracanal dressing, *Int Endod J* 24:119, 1991.

164. Sjogren U, Hagglund B, Sundqvist G, et al: Factors affecting the long-term results of endodontic treatment, *J Endod* 16:498, 1990.
165. Skaare AB, Jacobsen I: Dental injuries in Norwegians aged 7-18 years, *Dent Traumatol* 19:67, 2003.
166. Skieller V: The prognosis for young teeth loosened after mechanical injuries, *Acta Odontol Scand* 18:171, 1960.
167. Skoglund A, Tronstad L: Pulpal changes in replanted and autotransplanted immature teeth of dogs, *J Endod* 7:309, 1981.
168. Soder PO, Otteskog P, Andreasen JO, et al: Effect of drying on viability of periodontal membrane, *Scand J Dent Res* 85:164, 1977.
169. Spangberg L, Rutberg M, Rydinge E: Biologic effects of endodontic antimicrobial agents, *J Endod* 5:166, 1979.
170. Stalhane I, Hedegard B: Traumatized permanent teeth in children aged 7-15 years, *Sven Tandlak Tidskr* 68:157, 1975.
171. Stanley HR: Diseases of the dental pulp. In Tieck RW, editor: *Oral pathology*, New York, 1965, McGraw-Hill.
172. Stanley HR, Lundy T: Dycal therapy for pulp exposures, *Oral Surg Oral Med Oral Pathol* 34:818, 1972.
173. Subay RK, Demirci M: Pulp tissue reactions to a dentin bonding agent as a direct capping agent, *J Endod* 31:201, 2005.
174. Swift EJ Jr, Trope M: Treatment options for the exposed vital pulp, *Pract Periodontics Aesthet Dent* 11:735, 1999.
175. Tapias MA, Jimenez-Garcia R, Lamas F, et al: Prevalence of traumatic crown fractures to permanent incisors in a childhood population: Mostoles, Spain, *Dent Traumatol* 19:119, 2003.
176. Teixeira FB, Teixeira EC, Thompson JY, et al: Fracture resistance of roots endodontically treated with a new resin filling material, *J Am Dent Assoc* 135:646, 2004.
177. Torabinejad M, Hong CU, McDonald F, et al: Physical and chemical properties of a new root-end filling material, *J Endod* 21:349, 1995.
178. Torabinejad M, Rastegar AF, Kettering JD, et al: Bacterial leakage of mineral trioxide aggregate as a root-end filling material, *J Endod* 21:109, 1995.
179. Trabert KC, Caput AA, Abou-Rass M: Tooth fracture: a comparison of endodontic and restorative treatments, *J Endod* 4:341, 1978.
180. Tronstad L: Reaction of the exposed pulp to Dycal treatment, *Oral Surg Oral Med Oral Pathol* 38:945, 1974.
181. Tronstad L: Root resorption: etiology, terminology and clinical manifestations, *Endod Dent Traumatol* 4:241, 1988.
182. Tronstad L, Andreasen JO, Hasselgren G, et al: pH changes in dental tissues after root canal filling with calcium hydroxide, *J Endod* 7:17, 1980.
183. Tronstad L, Mjor IA: Capping of the inflamed pulp, *Oral Surg Oral Med Oral Pathol* 34:477, 1972.
184. Trope M: Root resorption of dental and traumatic origin: classification based on etiology, *Pract Periodontics Aesthet Dent* 10:515, 1998.
185. Trope M: Clinical management of the avulsed tooth: present strategies and future directions, *Dent Traumatol* 18:1, 2002.
186. Trope M, Friedman S: Periodontal healing of replanted dog teeth stored in Viaspan, milk and Hank's balanced salt solution, *Endod Dent Traumatol* 8:183, 1992.
187. Trope M, Moshonov J, Nissan R, et al: Short vs long-term calcium hydroxide treatment of established inflammatory root resorption in replanted dog teeth, *Endod Dent Traumatol* 11:124, 1995.
188. Trope M, Yesilsoy C, Koren L, et al: Effect of different endodontic treatment protocols on periodontal repair and root resorption of replanted dog teeth, *J Endod* 18:492, 1992.
189. Turley PK, Joiner MW, Hellstrom S: The effect of orthodontic extrusion on traumatically intruded teeth, *Am J Orthod* 85:47, 1984.
190. von Arx T, Filippi A, Buser D: Splinting of traumatized teeth with a new device: TTS (titanium trauma splint), *Dent Traumatol* 17:180, 2001.
191. Wedenberg C, Lindsog S: Experimental internal resorption in monkey teeth, *Endod Dent Traumatol* 1:221, 1985.
192. Wedenberg C, Zetterqvist L: Internal resorption in human teeth: a histological, scanning electron microscopic, and enzyme histochemical study, *J Endod* 13:255, 1987.
193. Weiss M: Pulp capping in older patients, *NY State Dent J* 32:451, 1966.
194. Windley W, Teixeira F, Levin L, et al: Disinfection of immature teeth with a triple antibiotic paste, *J Endod* 31:439, 2005.
195. Witherspoon DE, Small JC, Harris GZ: Mineral trioxide aggregate pulpotomies: a case series outcomes assessment, *J Am Dent Assoc* 137:610, 2006.
196. Yanpiset K, Trope M: Pulp revascularization of replanted immature dog teeth after different treatment methods, *Endod Dent Traumatol* 16:211, 2000.
197. Yanpiset K, Vongsavan N, Sigurdsson A, et al: Efficacy of laser Doppler flowmetry for the diagnosis of revascularization of reimplanted immature dog teeth, *Dent Traumatol* 17:63, 2001.
198. Yoldaş SE, Bani M, Atabek D, et al: Comparison of the potential discoloration effect of Bioaggregate, Biodentine, and white mineral trioxide aggregate on bovine teeth: In vitro research, *J Endod* 42:1815, 2016.
199. Zachrisson BU, Jacobsen I: Long-term prognosis of 66 permanent anterior teeth with root fracture, *Scand J Dent Res* 83:345, 1975.