

1

Diagnosis

LOUIS H. BERMAN and ILAN ROTSTEIN

CHAPTER OUTLINE

Art and Science of Diagnosis

Chief Complaint

Medical History

Dental History

History of Present Dental Problem

Dental History Interview

Examination and Testing

Extraoral Examination

Intraoral Examination

Soft-Tissue Examination

Intraoral Swelling

Intraoral Sinus Tracts

Palpation

Percussion

Mobility

Periodontal Examination

Pulp Tests

Thermal

Electric

Laser Doppler Flowmetry

Pulse Oximetry

Special Tests

Bite Test

Test Cavity

Staining and Transillumination

Selective Anesthesia

Radiographic Examination and Interpretation

Intraoral Radiographs

Digital Radiography

Cone-Beam Computerized Tomography

Magnetic Resonance Imaging

Cracks and Fractures

Crack Types

Vertical Root Fractures

Perforations

Clinical Classification of Pulpal and Periapical Diseases

Pulpal Disease

Normal Pulp

Pulpitis

Reversible Pulpitis

Irreversible Pulpitis

Previously Treated

Previously Initiated Therapy

Apical (Periapical) Disease

Normal Apical Tissues

Periodontitis

Symptomatic Apical Periodontitis

Asymptomatic Apical Periodontitis

Acute Apical Abscess

Chronic Apical Abscess

Referred Pain

Summary

Art and Science of Diagnosis

Diagnosis is the art and science of detecting and distinguishing deviations from health and the cause and nature thereof.⁶ The purpose of a diagnosis is to determine what problem the patient is having and why the patient is having that problem. Ultimately, this will directly relate to what treatment, if any, will be necessary. No appropriate treatment recommendation can be made until all of the *whys* are answered. Therefore, careful data gathering as well as a planned, methodical, and systematic approach to this investigatory process is crucial.

Gathering objective data and obtaining subjective findings are not enough to formulate an accurate clinical diagnosis. The data must be interpreted and processed to determine what information is significant, and what information might be questionable. The facts need to be collected with an active dialogue between the clinician and the patient, with the clinician asking the right questions

and carefully interpreting the answers. In essence, the process of determining the existence of an oral pathosis is the culmination of the art and science of making an accurate diagnosis.

The process of making a diagnosis can be divided into five stages:

1. The patient tells the clinician the reasons for seeking advice.
2. The clinician questions the patient about the symptoms and history that led to the visit.
3. The clinician performs objective clinical tests.
4. The clinician correlates the objective findings with the subjective details and creates a tentative list of differential diagnoses.
5. The clinician formulates a definitive diagnosis.

This information is accumulated by means of an organized and systematic approach that requires considerable clinical judgment. The clinician must be able to approach

the problem by crafting what questions to ask the patient and how to ask these pertinent questions. Careful listening is paramount to begin painting the picture that details the patient's complaint. These subjective findings combined with the results of diagnostic tests provide the critical information needed to establish the diagnosis.

Neither the art nor the science is effective alone. Establishing a differential diagnosis in endodontics requires a unique blend of knowledge, skills, and the ability to interpret and interact with a patient in real time. Questioning, listening, testing, interpreting, and finally answering the ultimate question of *why* will lead to an accurate diagnosis and in turn result in a more successful treatment plan.

CHIEF COMPLAINT

On arrival for a dental consultation, the patient should complete a thorough registration that includes information pertaining to medical and dental history (Figs. 1.1 and 1.2). This should be signed and dated by the patient, as well as initialed by the clinician as verification that all of the submitted information has been reviewed (see Chapter 27 for more information).

The reasons patients give for consulting with a clinician are often as important as the diagnostic tests performed. Their remarks serve as initial important clues that will help the clinician to formulate a correct diagnosis. Without these direct and unbiased comments, objective findings may lead to an incorrect diagnosis. The clinician may find a dental pathosis, but it may not contribute to the pathologic condition that mediates the patient's chief complaint. Investigating these complaints may indicate that the patient's concerns are related to a medical condition or to recent dental treatment. Certain patients may even receive initial emergency treatment for pulpal or periapical symptoms in a general hospital.⁹³ On occasion, the chief complaint is simply that another clinician correctly or incorrectly advised the patient that he or she had a dental problem, with the patient not necessarily having any symptoms or any objective pathosis. Therefore, the clinician must pay close attention to the actual expressed complaint, determine the chronology of events that led to this complaint, and question the patient about other pertinent issues, including medical and dental history. For future reference and in order to ascertain a correct diagnosis, the patient's chief complaint should be properly documented, using *the patient's own words*.

MEDICAL HISTORY

The clinician is responsible for taking a proper medical history from every patient who presents for treatment. Numerous examples of medical history forms are available from a variety of sources, or clinicians may choose to customize their own forms. After the form is completed by the patient, or by the parent or guardian in the case of a minor, the clinician should review the responses with the patient, parent, or guardian, and then initial the medical history form to indicate that this review has been done. The patient "of record" should be questioned at each treatment visit to determine whether there have been any changes in the patient's medical history or medications. A more thorough

and complete update of the patient's medical history should be taken if the patient has not been seen for over a year.^{51,52}

Baseline blood pressure and pulse should be recorded for the patient at each treatment visit. Elevation in blood pressure or a rapid pulse rate may indicate an anxious patient who may require a stress reduction protocol, or it may indicate that the patient has hypertension or other cardiovascular health problems. Referral to a physician or medical facility may be indicated. It is imperative that vital signs be gathered at each treatment visit for any patient with a history of major medical problems. The temperature of patients presenting with subjective fever or any signs or symptoms of a dental infection should be taken.^{57,80,105}

The clinician should evaluate a patient's response to the health questionnaire from two perspectives: (1) those medical conditions and current medications that will necessitate altering the manner in which dental care will be provided and (2) those medical conditions that may have oral manifestations or mimic dental pathosis.

Patients with serious medical conditions may require either a modification in the manner in which the dental care will be delivered or a modification in the dental treatment plan (Box 1.1). In addition, the clinician should be aware if the patient has any drug allergies or interactions, allergies to dental products, an artificial joint prosthesis, organ transplants, or is taking medications that may negatively interact with common local anesthetics, analgesics, sedatives, and antibiotics.⁸⁰ This may seem overwhelming, but it emphasizes the importance of obtaining a thorough and accurate medical history while considering the various medical conditions and dental treatment modifications that may be necessary before dental treatment is provided.

Several medical conditions have oral manifestations, which must be carefully considered when attempting to arrive at an accurate dental diagnosis. Many of the oral soft-tissue changes that occur are more related to the medications used to treat the medical condition rather than to the condition itself. More common examples of medication side effects are stomatitis, xerostomia, petechiae, ecchymoses, lichenoid mucosal lesions, and bleeding of the oral soft tissues.⁸⁰

When developing a dental diagnosis, a clinician must also be aware that some medical conditions can have clinical presentations that mimic oral pathologic lesions.^{13,28,32,74,80,102,107,133} For example, tuberculosis involvement of the cervical and submandibular lymph nodes can lead to a misdiagnosis of lymph node enlargement secondary to an odontogenic infection. Lymphomas can involve these same lymph nodes.⁸⁰ Immunocompromised patients and patients with uncontrolled diabetes mellitus respond poorly to dental treatment and may exhibit recurring abscesses in the oral cavity that must be differentiated from abscesses of dental origin.^{43,76,80,83} Patients with iron deficiency anemia, pernicious anemia, and leukemia frequently exhibit paresthesia of the oral soft tissues. This finding may complicate making a diagnosis when other dental pathosis is present in the same area of the oral cavity. Sickle cell anemia has the complicating factor of bone pain, which mimics odontogenic pain, and loss of trabecular bone pattern on radiographs, which can be confused with radiographic lesions of endodontic origin. Multiple myeloma can result in unexplained mobility of teeth. Radiation therapy

TELL US ABOUT YOUR SYMPTOMS

LAST NAME _____ FIRST NAME _____

1. Are you experiencing any pain at this time? If not, please go to question 6. Yes _____ No _____

2. If yes, can you locate the tooth that is causing the pain? Yes _____ No _____

3. When did you first notice the symptoms? _____

4. Did your symptoms occur suddenly or gradually? _____

5. Please check the frequency and quality of the discomfort, and the number that most closely reflects the intensity of your pain:

LEVEL OF INTENSITY (On a scale of 1 to 10) 1 = Mild 10 = Severe	FREQUENCY	QUALITY
1 _____ 2 _____ 3 _____ 4 _____ 5 _____ 6 _____ 7 _____ 8 _____ 9 _____ 10 _____	_____ Constant	_____ Sharp
	_____ Intermittent	_____ Dull
	_____ Momentary	_____ Throbbing
	_____ Occasional	

Is there anything you can do to relieve the pain? Yes _____ No _____

If yes, what? _____

Is there anything you can do to cause the pain to increase? Yes _____ No _____

If yes, what? _____

When eating or drinking, is your tooth sensitive to: Heat _____ Cold _____ Sweets _____

Does your tooth hurt when you bite down or chew? Yes _____ No _____

Does it hurt if you press the gum tissue around this tooth? Yes _____ No _____

Does a change in posture (lying down or bending over) cause your tooth to hurt? Yes _____ No _____

6. Do you grind or clench your teeth? Yes _____ No _____

7. If yes, do you wear a night guard? Yes _____ No _____

8. Has a restoration (filling or crown) been placed on this tooth recently? Yes _____ No _____

9. Prior to this appointment, has root canal therapy been initiated on this tooth? Yes _____ No _____

10. Is there anything else we should know about your teeth, gums, or sinuses that would assist us in our diagnosis? _____

Signed: Patient or Parent _____ Date _____

Fig. 1.1 Dental history form that also allows the patient to record pain experience in an organized and descriptive manner.

TELL US ABOUT YOUR HEALTH

LAST NAME _____ **FIRST NAME** _____

How would you rate your health?

Please circle one.

Excellent Good Fair Poor

When did you have your last physical exam? _____

If you are under the care of a physician, please give reason(s) for treatment.

Physician's Name, Address, and Telephone Number:

Name _____ Address _____

City _____ State _____ Zip _____ Telephone _____

Have you ever had any kind of surgery?

Yes _____ No _____

If yes, what kind? _____ Date _____

_____ Date _____

Have you ever had any trouble with prolonged bleeding after surgery?

Yes _____ No _____

Do you wear a pacemaker or any other kind of prosthetic device?

Yes _____ No _____

Are you taking any kind of medication or drugs at this time?

Yes _____ No _____

If yes, please give name(s) of the medicine(s) and reason(s) for taking them:

Name _____ Reason _____

Have you ever had an unusual reaction to an anesthetic or drug (like penicillin)?

Yes _____ No _____

If yes, please explain: _____

Please circle any past or present illness you have had:

Alcoholism	Blood pressure	Epilepsy	Hepatitis	Kidney or liver	Rheumatic fever
Allergies	Cancer	Glaucoma	Herpes	Mental	Sinusitis
Anemia	Diabetes	Head/Neck injuries	Immunodeficiency	Migraine	Ulcers
Asthma	Drug dependency	Heart disease	Infectious diseases	Respiratory	Venereal disease

Are you allergic to Latex or any other substances or materials?

Yes _____ No _____

If so, please explain _____

If female, are you pregnant?

Yes _____ No _____

Is there any other information that should be known about your health? _____

Signed: Patient or Parent _____ Date: _____

Fig. 1.2 Succinct, comprehensive medical history form designed to provide insight into systemic conditions that could produce or affect the patient's symptoms, mandate alterations in treatment modality, or change the treatment plan.

Box 1.1 Medical Conditions That Warrant Modification of Dental Care or Treatment

Cardiovascular: High- and moderate-risk categories of endocarditis, pathologic heart murmurs, hypertension, unstable angina pectoris, recent myocardial infarction, cardiac arrhythmias, poorly managed congestive heart failure.^{57,80,105}

Pulmonary: Chronic obstructive pulmonary disease, asthma, tuberculosis.^{80,129}

Gastrointestinal and renal: End-stage renal disease; hemodialysis; viral hepatitis (types B, C, D, and E); alcoholic liver disease; peptic ulcer disease; inflammatory bowel disease; pseudomembranous colitis.^{25,34,48,80}

Hematologic: Sexually transmitted diseases, human immunodeficiency virus (HIV) and acquired immunodeficiency syndrome (AIDS), diabetes mellitus, adrenal insufficiency, hyperthyroidism and hypothyroidism, pregnancy, bleeding disorders, cancer and leukemia, osteoarthritis and rheumatoid arthritis, systemic lupus erythematosus.^{35,43,76,80,83,88,100,135}

Neurologic: Cerebrovascular accident, seizure disorders, anxiety, depression and bipolar disorders, presence or history of drug or alcohol abuse, Alzheimer disease, schizophrenia, eating disorders, neuralgias, multiple sclerosis, Parkinson disease.^{36,44,80}

to the head and neck region can result in increased sensitivity of the teeth and osteoradionecrosis.⁸⁰ Trigeminal neuralgia, referred pain from cardiac angina, and multiple sclerosis can also mimic dental pain (see also Chapter 4). Acute maxillary sinusitis is a common condition that may create diagnostic confusion because it may mimic tooth pain in the maxillary posterior quadrant. In this situation the teeth in the quadrant may be extremely sensitive to cold and percussion, thus mimicking the signs and symptoms of pulpitis. This is certainly not a complete list of all the medical entities that can mimic dental disease, but it should alert the clinician that a medical problem could confuse and complicate the diagnosis of dental pathosis; this issue is discussed in more detail in subsequent chapters.

If, at the completion of a thorough dental examination, the subjective, objective, clinical testing, and radiographic findings do not result in a diagnosis with an obvious dental origin, then the clinician must consider that an existing medical problem could be the true source of the pathosis. In such instances, a consultation with the patient's physician is always appropriate.

DENTAL HISTORY

The chronology of events that lead up to the chief complaint is recorded as the *dental history*. This information will help guide the clinician as to which diagnostic tests are to be performed. The history should include any past and present symptoms, as well as any procedures or trauma that might have evoked the chief complaint. Proper documentation is imperative. It may be helpful to use a premade form to record the pertinent information obtained during the dental history interview and diagnostic examination. Often a SOAP format is used, with the history and findings documented under the categories of Subjective, Objective, Appraisal, and Plan. There are also built-in features within some practice management software packages that allow digital entries into the patient's electronic file for the diagnostic workup (Figs. 1.3 and 1.4).

History of Present Dental Problem

The dialogue between the patient and the clinician should encompass all of the details pertinent to the events that led to the chief complaint. The clinician should direct the conversation in a manner that produces a clear and concise narrative that chronologically depicts all of the necessary information about the patient's symptoms and the development of these symptoms. To help elucidate this information, the patient is first instructed to fill out a dental history form as part of the patient's office registration. This information will help the clinician decide which approach to use when asking the patient questions. The interview first determines *what is going on* in an effort to determine *why it is going on* for the purpose of eventually determining *what is necessary to resolve the chief complaint*.

Dental History Interview

After starting the interview and determining the nature of the chief complaint, the clinician continues the conversation by documenting the sequence of events that initiated the request for an evaluation. The dental history is divided into five basic directions of questioning: localization, commencement, intensity, provocation or attenuation, and duration.

Localization. "Can you point to the offending tooth?" Often the patient can point to or tap the offending tooth. This is the most fortunate scenario for the clinician because it helps direct the interview toward the events that might have caused any particular pathosis in this tooth. In addition, localization allows subsequent diagnostic tests to focus more on this particular tooth. When the symptoms are not well localized, the diagnosis is a greater challenge.

Commencement. "When did the symptoms first occur?" A patient who is having symptoms often remembers when these symptoms started. Sometimes the patient will even remember the initiating event. It may have been spontaneous in nature; it may have begun after a dental visit for a restoration; trauma may be the etiology; biting on a hard object may have initially produced the symptoms; or the initiating event may have occurred concurrently with other symptoms (e.g., sinusitis, headache, chest pain). However, the clinician should resist the tendency to make a premature diagnosis based on these circumstances. The clinician should not simply assume "guilt by association" but instead should use this information to enhance the overall diagnostic process.

Intensity. "How intense is the pain?" It often helps to quantify how much pain the patient is actually having. The clinician might ask, "On a scale from 1 to 10, with 10 the most severe, how would you rate your symptoms?" Hypothetically, a patient could present with "an uncomfortable sensitivity to cold" or "an annoying pain when chewing" but might rate this "pain" only as a 2 or a 3. These symptoms certainly contrast with the type of symptoms that prevent a patient from sleeping at night. Often the intensity can be subjectively measured by what is necessary for the diminution of pain (e.g., acetaminophen versus a narcotic pain reliever). This intensity level may affect the decision to treat or not to treat with endodontic therapy. Pain is now considered a standard vital sign, and documenting pain intensity (scale of 0 to 10) provides a baseline for comparison after treatment.

Name: (Last) _____ (First) _____ Date: _____ Tooth: _____

S. (SUBJECTIVE)

Chief Complaint:

History of Present Illness:

Nature of Pain: None Mild Moderate Severe
Quality: Dull Sharp Throbbing Constant
Onset: Stim Required Intermittent Spontaneous
Location: Localized Diffuse Referred Radiating to:
Duration: Seconds Minutes Hours Constant
Initiated by: Cold Heat Sweet Spontaneous Palpation Mastication Supination Keeps awake at night
Relieved by: Cold Heat OTC-Meds Narc-Meds

O. (OBJECTIVE)

Extraoral:

Facial swelling: Yes No
L Nodes swollen: Yes No

Intraoral:

Soft tissues: WNL
Swelling: Yes No Mild Moderate Severe Location:
Sinus tract: Yes No Closed
Clinical crown: Restn Caries Exposure Fracture

#	Cold	Heat	EPT	Perc	Palp	Mob	Bite Stick	Dis-color	Periodontal Exam								
									MB	B	DB	DL	L	ML	Recess	Furcation	Bleed-Probing

(Normal: N No Response: 0 Mild: + Moderate: ++ Severe: +++ Lingered: L Delayed: D)

Radiographic Findings:

Alveolar Bone: WNL Apical lucency Lateral lucency Ap / Lat opacity Crestal bone loss
Lamina Dura: WNL Obscure Broken Widened
Roots: WNL Curvature Resorption Perforation Dilaceration Fracture Long Sinus / IAN
Pulp Chamber: WNL Calcification Pulp Stone Exposure Resorption Perforation
Pulp Canal: WNL Calcification Bifurcated Resorption Prior RCT Furcation Involvement Perforation
Crown: WNL Caries Restoration Crown Dens in dente
Sinus Tract: Traces to:

A. (Assessment)

Diagnosis: **Pulpal:** WNL Rev Pulpitis Irrev Pulpitis Necrosis Prior RCT / Non-healing Pulpless
Periapical: WNL APP CPP APA CPA Cond Osteitis
Etiology: Caries Restoration Prior RCT Iatrogenic Coronal leakage Trauma Perio Elective Resorptn VRF
Prognosis: Good Fair Poor

P. (PLAN)

Endodontic: Caries control RCT ReTx I&D Apico Apexification/genesis Perf / Resorption Repair
Periodontal: S/RP Crown lengthen Root amp Hemisection Extraction
Restorative: Temp Post space B/U P&C Onlay / Crown Bleach

Fig. 1.3 When taking a dental history and performing a diagnostic examination, often a premade form can facilitate complete and accurate documentation. (Courtesy Dr. Ravi Koka, San Francisco, CA.)

Consultation on 02/10/2005

Chief Complaint
Hurts when chews. Has gotten worse over the last two weeks.

Symptoms
☒ Symptomatic
☐ Asymptomatic

Location
☐ Upper Right
☐ Lower Right
☐ Upper Left
☐ Lower Left
☐ Referred
☐ Radiating

Chronology
☐ Constant
☒ Intermittent
☐ Momentary
☐ Lingering
☐ Referred

Quality
☒ Sharp
☐ Dull
☐ Throbbing
☐ Steady
☐ Enlarging
☐ Spontaneous

Affected By
☐ Hot
☐ Cold
☒ Biting
☒ Chewing
☐ Percussion
☐ Palpitation

Visual Exam
☐ Normal
☐ Discoloration
☐ Caries
☐ Pulp Exposure
☐ Prior Access
☐ Swelling

Probable Tooth
[Dropdown]

Exam Date 02/10/2005

Sinus Tract	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]
Clinical Crown	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]
Discolored	☐	☐	☐	☐	☐	☐	☐
Facial Swelling	☐	☐	☐	☐	☐	☐	☐
Caries	☐	☐	☐	☐	☐	☐	☐
Previous RCT	☐	☐	☐	☐	☐	☐	☐
Large Restore	☐	☐	☐	☐	☐	☐	☐

Radiology

Alveolar Bone	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]
Lamina Dura	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]
Roots	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]
Pulp Chamber	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]
Pulp Canal	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]	[Dropdown]
Straight Canals	0	0	0	0	0	0	0
Curved Canals	0	0	0	0	0	0	0

Pulp Test

Tooth # -> 29 30

Cold	No Response	Hyper	Hyper
Hot	No Response	WNL	No response
Percussion	WNL	Severe	Moderate
EPT	[Dropdown]	[Dropdown]	[Dropdown]
Palpation	[Dropdown]	[Dropdown]	[Dropdown]
Mobility	Normal	Normal	Normal
Bite Stick	[Dropdown]	[Dropdown]	[Dropdown]
Swelling	[Dropdown]	[Dropdown]	[Dropdown]
Sinus Tract	[Dropdown]	[Dropdown]	[Dropdown]
Clinical Crown	[Dropdown]	[Dropdown]	[Dropdown]
Discolored	☐	☐	☐
Facial Swelling	☐	☐	☐
Caries	☐	☐	☐
Previous RCT	☐	☐	☐
Large Restore	☐	☐	☐

Radiology

Alveolar Bone	[Dropdown]	[Dropdown]
---------------	------------	------------

Diagnosis

Date 02/10/2010

Clinical Diagnosis
Tooth 30

Reason for Endodontic Therapy
☒ Pathosis
☐ Restorative
☐ Fracture
☐ Retreatment
☐ Evulsed

Pulpal
☐ Normal
☐ Reversible Pulpitis
☐ Irreversible Pulpitis
☒ Necrotic

Periradicular
☐ Normal
☐ Acute Apical Abscess
☐ Chronic Apical Abscess
☐ Abscess

Anticipated Restoration
☐ Amalgam/Composite
☒ Crown/Bridge
☐ Post & Core
☐ Onlays/Overlays

Referral Preference for Tooth Completion
Post

Prognosis Endo Favorable Perio [Dropdown]

Endodontist TBJ Tom B. Johnson, D.D.S. TBJ

Notes

Fig. 1.4 Several practice management software packages have features for charting endodontic diagnoses using user-defined drop-down menus and areas for specific notations. Note that for legal purposes, it is desirable that all recorded documentation have the ability to be locked, or if any modifications are made after 24 hours, the transaction should be recorded with an automated time/date stamp. This is necessary so the data cannot be fraudulently manipulated. (Courtesy PBS Endo, Cedar Park, TX.)

Provocation or attenuation. “What produces or reduces the symptoms?” Mastication and locally applied temperature changes account for the majority of initiating factors that cause dental pain. The patient may relate that drinking something cold causes the pain or possibly that chewing or biting is the only stimulus that “makes it hurt.” The patient might say that the pain is only reproduced on “release from biting.” On occasion, a patient may present to the dental office with a cold drink in hand and state that the symptoms can only be *reduced* by bathing the tooth in cold water. Non-prescription pain relievers may relieve some symptoms, whereas narcotic medication may be required to reduce others. Note that patients who are using narcotic as well as non-narcotic (e.g., ibuprofen) analgesics may respond differently to questions and diagnostic tests, thereby altering the validity of diagnostic results. Thus, it is important to know what drugs patients have taken in the previous 4 to 6 hours. These provoking and relieving factors may help the clinician to determine which diagnostic tests should be performed to establish a more objective diagnosis.

Duration. “Do the symptoms subside shortly, or do they linger after they are provoked?” The difference between a cold sensitivity that subsides in a few seconds and one that subsides in minutes may determine whether a clinician repairs a defective restoration or provides endodontic treatment. The duration of symptoms after a stimulating event should be recorded to establish how long the patient felt the sensation in terms of seconds or minutes. Clinicians often first test control teeth (possibly including a contralateral “normal” tooth) to define a “normal” response for the patient; thus, “lingering” pain is apparent when comparing the duration between the control teeth and the suspected tooth.

With the dental history interview complete, the clinician has a better understanding of the patient’s chief complaint and can concentrate on making an objective diagnostic evaluation, although the subjective (and artistic) phase of making a diagnosis is not yet complete and will continue

after the more objective testing and scientific phase of the investigatory process.

Examination and Testing

EXTRAORAL EXAMINATION

Basic diagnostic protocol suggests that a clinician observe patients as they enter the operatory. Signs of physical limitations may be present, as well as signs of facial asymmetry that result from facial swelling. Visual and palpation examinations of the face and neck are warranted to determine whether swelling is present. Many times a facial swelling can be determined only by palpation when a unilateral “lump or bump” is present. The presence of bilateral swellings may be a normal finding for any given patient; however, it may also be a sign of a systemic disease or the consequence of a developmental event. Palpation allows the clinician to determine whether the swelling is localized or diffuse, firm or fluctuant. These latter findings will play a significant role in determining the appropriate treatment.

Palpation of the cervical and submandibular lymph nodes is an integral part of the examination protocol. If the nodes are found to be firm and tender along with facial swelling and an elevated temperature, there is a high probability that an infection is present. The disease process has moved from a localized area immediately adjacent to the offending tooth to a more widespread systemic involvement.

Extraoral facial swelling of odontogenic origin typically is the result of endodontic etiology because diffuse facial swelling resulting from a periodontal abscess is rare. Swellings of nonodontogenic origin must always be considered in the differential diagnosis, especially if an obvious dental pathosis is not found.⁷⁷ This situation is discussed in subsequent chapters.

A subtle visual change such as loss of definition of the nasolabial fold on one side of the nose may be the earliest sign of a canine space infection (Fig. 1.5). Pulpal necrosis



Fig. 1.5 **A**, Canine space swelling of the left side of the face extending into and involving the left eye. **B**, Swelling of the upper lip and the loss of definition of the nasolabial fold on the patient’s left side, which indicates an early canine space infection.

and periradicular disease associated with a maxillary canine should be suspected as the source of the problem. Extremely long maxillary central incisors also may be associated with a canine space infection, but most extraoral swellings associated with the maxillary centrals express themselves as a swelling of the upper lip and base of the nose.

If the buccal space becomes involved, the swelling will be extraoral in the area of the posterior cheek (Fig. 1.6). These swellings are generally associated with infections originating from the buccal root apices of the maxillary premolar and molar teeth and the mandibular premolar (Fig. 1.7) and first molar teeth. The mandibular second and third molars also may be involved, but infections associated with these two teeth are just as likely to exit to the lingual where other spaces would be involved. For infections associated with these teeth, the root apices of the maxillary teeth must lie superior to the attachment of the buccinator muscle to the maxilla, and the apices of the mandibular teeth must be inferior to the buccinator muscle attachment to the mandible.⁷⁷

Extraoral swelling associated with mandibular incisors will generally exhibit itself in the submental (Fig. 1.8) or submandibular space. Infections associated with any mandibular teeth, which exit the alveolar bone on the lingual and are inferior to the mylohyoid muscle attachment, will



Fig. 1.6 Buccal space swelling associated with an acute periradicular abscess from the mandibular left second molar.

be noted as swelling in the submandibular space. Further discussions of fascial space infections may be found in Chapter 16.

Sinus tracts of odontogenic origin may also open through the skin of the face (Figs. 1.9 and 1.10).^{2,56,64} These openings in the skin will generally close once the offending tooth is treated and healing occurs. A scar is more likely to be visible on the skin surface in the area of the sinus tract stoma than on the oral mucosal tissues (see Fig. 1.10, C and D). Many patients with extraoral sinus tracts give a history of being treated by general physicians, dermatologists, oncologists, or plastic surgeons with systemic or topical antibiotics or surgical procedures in attempts to heal the extraoral stoma. In these particular cases, after multiple treatment failures, the patients may finally be referred to a dental clinician to determine whether there is a dental cause. Raising the awareness of physicians to such cases will aid in more accurate diagnosis and faster referral to the dentist or endodontist.

INTRAORAL EXAMINATION

The intraoral examination may give the clinician insight as to which intraoral areas may need a more focused evaluation. Any abnormality should be carefully examined for either prevention or early treatment of associated pathosis.^{4,30,75,113,110,126} Swelling, localized lymphadenopathy, or a sinus tract should provoke a more detailed assessment of related and proximal intraoral structures.

Soft-Tissue Examination

As with any dental examination, there should be a routine evaluation of the intraoral soft tissues. The gingiva and mucosa should be dried with either a low-pressure air syringe or a 2-by-2-inch gauze pad. By retracting the tongue and cheek, all of the soft tissue should be examined for abnormalities in color or texture. Any raised lesions or ulcerations should be documented and, when necessary, evaluated with a biopsy or referral.⁸²

Intraoral Swelling

Intraoral swellings should be visualized and palpated to determine whether they are diffuse or localized and whether




Fig. 1.7 **A**, Buccal space swelling of the left side of the patient's face. Note the asymmetry of the left side of the face. **B**, Intraoral view of another patient shows swelling present in the left posterior mucobuccal fold. **C**, This buccal space infection was associated with periradicular disease from the mandibular left first molar. Note on the radiograph the periradicular radiolucency and incomplete endodontic treatment. (B and C, Courtesy Dr. Jaydeep S. Talim, Los Angeles, CA.)



Fig. 1.8 Swelling of the submental space associated with periradicular disease from the mandibular incisors.

they are firm or fluctuant. These swellings may be present in the attached gingiva, alveolar mucosa, mucobuccal fold, palate, or sublingual tissues. Other testing methods are required to determine whether the origin is endodontic, periodontic, or a combination of these two or whether it is of nonodontogenic origin.

Swelling in the anterior part of the palate (**Fig. 1.11**) is most frequently associated with an infection present at the apex of the maxillary lateral incisor or the palatal root of the maxillary first premolar. More than 50% of the maxillary lateral incisor root apices deviate in the distal or palatal directions. A swelling in the posterior palate (**Fig. 1.12**) is most likely associated with the palatal root of one of the maxillary molars.⁷⁷

Intraoral swelling present in the mucobuccal fold (**Fig. 1.13**) can result from an infection associated with the apex of the root of any maxillary tooth that exits the alveolar bone on the facial aspect and is inferior to the muscle attachment present in that area of the maxilla (see also Chapter 15). The same is true with the mandibular teeth if the root apices are superior to the level of the muscle attachments and the infection exits the bone on the facial. Intraoral swelling can also occur in the sublingual space if the infection from the root apex spreads to the lingual and exits the alveolar bone superior to the attachment for the mylohyoid muscle. The tongue will be elevated and the swelling will be bilateral because the sublingual space is contiguous with no midline separation. If the infection exits the alveolar bone to the lingual with mandibular molars and is inferior to the attachment of the mylohyoid muscle, the swelling will be noted in the submandibular space. Severe infections involving the maxillary and mandibular molars can extend into the parapharyngeal space, resulting in intraoral swelling of the tonsillar and pharyngeal areas. This can be life threatening if the patient's airway becomes obstructed.^{77,80}

Intraoral Sinus Tracts

On occasion, a chronic endodontic infection will drain through an intraoral communication to the gingival surface and is known as a *sinus tract*.¹² This pathway, which is sometimes lined with epithelium, extends directly from the source of the infection to a surface opening, or *stoma*, on




Fig. 1.9 **A**, Extraoral drainage associated with periradicular disease from the mandibular right canine. Note the parulis on the right anterior side of the face. **B**, Initial scar associated with the extraoral drainage incision after the parulis was drained and root canal therapy performed on the canine. **C**, Three-month follow-up shows healing of the incision area. Note the slight inversion of the scar tissue.

the attached gingival surface. As previously described, it can also extend extraorally. The term *fistula* is often inappropriately used to describe this type of drainage. The fistula, by definition, is actually an abnormal communication pathway between two internal organs or from one epithelium-lined surface to another epithelium-lined surface.⁶




Fig. 1.10 **A**, Extraoral sinus tract opening onto the skin in the central chin area. **B**, Radiograph showing large radiolucency associated with the mandibular incisors. **C**, A culture is obtained from the drainage of the extraoral sinus tract. **D**, The healed opening of the extraoral sinus tract 1 month after root canal therapy was completed. Note the slight skin concavity in the area of the healed sinus tract.




Fig. 1.11 Fluctuant swelling in the anterior palate associated with periapical disease from the palatal root of the maxillary first premolar.

Histologic studies have found that most sinus tracts are not lined with epithelium throughout their entire length. One study found that only 1 out of the 10 sinus tracts examined were lined with epithelium, whereas the other 9 specimens were lined with granulation tissue.⁵⁵ Another study, with a larger sample size, found that two thirds of the specimens did not have epithelium extending beyond the level of the surface mucosa rete ridges.¹² The remaining specimens had some epithelium that extended from the oral mucosa surface to the periradicular lesion.¹² The presence or absence of an epithelial lining does not seem to prevent closure of the tract as long as the source of the problem is properly diagnosed and adequately treated and the endodontic lesion has healed. Failure of a sinus tract to heal after treatment will necessitate further diagnostic procedures to determine whether other sources of infection are present or whether a misdiagnosis occurred.

In general, a periapical infection that has an associated sinus tract is not painful, although often there is a history of varying magnitudes of discomfort before sinus tract



Fig. 1.12 Fluctuant swelling in the posterior palate associated with periradicular disease from the palatal root of the maxillary first molar.



Fig. 1.13 Fluctuant swelling in the mucobuccal fold associated with periradicular disease from the maxillary central incisor.

development. Besides providing a conduit for the release of infectious exudate and the subsequent relief of pain, the sinus tract can also provide a useful aid in determining the source of a given infection. Sometimes objective evidence as to the origin of an odontogenic infection is lacking. The stoma of the sinus tract may be located directly adjacent to or at a distant site from the infection. Tracing the sinus tract will provide objectivity in diagnosing the location of the problematic tooth. To trace the sinus tract, a size #25 or #30 gutta-percha cone is threaded into the opening of the

sinus tract. Although this may be slightly uncomfortable to the patient, the cone should be inserted until resistance is felt. After a periapical radiograph is exposed, the origin of the sinus tract is determined by following the path taken by the gutta-percha cone (**Fig. 1.14**). This will direct the




Fig. 1.14 **A**, To locate the source of an infection, the sinus tract can be traced by threading the stoma with a gutta-percha point. **B**, Radiograph of the area shows an old root canal in a maxillary second premolar and a questionable radiolucent area associated with the first premolar, with no clear indication of the etiology of the sinus tract. **C**, After tracing the sinus tract, the gutta-percha is seen to be directed to the source of pathosis, the apex of the maxillary first premolar.

clinician to the tooth involved and, more specifically, to the part of the root of the tooth that is the source of the pathosis. Once the causative factors related to the formation of the sinus tract are removed, the stoma and the sinus tract will close within several days.

The stomata of intraoral sinus tracts may open in the alveolar mucosa, in the attached gingiva, or through the furcation or gingival crevice. They may exit through either the facial or the lingual tissues depending on the proximity of the root apices to the cortical bone. If the opening is in the gingival crevice, it is normally present as a narrow defect in one or two isolated areas along the root surface. When a narrow defect is present, the differential diagnosis must include the opening of a periradicular endodontic lesion, a vertical root fracture, or the presence of a developmental groove on the root surface. This type of sinus tract can be differentiated from a primary periodontal lesion because the latter generally presents as a pocket with a broad coronal opening and more generalized alveolar bone loss around the root. Other pulp testing methods may assist in verifying the source of infection.^{111,112,121}

Palpation

In the course of the soft-tissue examination, the alveolar hard tissues should also be palpated. Emphasis should be placed on detecting any soft-tissue swelling or bony expansion, especially noting how it compares with and relates to the adjacent and contralateral tissues. In addition to objective findings, the clinician should question the patient about any areas that feel unusually sensitive during this palpation part of the examination.

A palpation test is performed by applying firm digital pressure to the mucosa covering the roots and apices. The index finger is used to press the mucosa against the underlying cortical bone. This will detect the presence of periradicular abnormalities or specific areas that produce painful response to digital pressure. A positive response to palpation may indicate an active periradicular inflammatory process. However, this test does not indicate whether the inflammatory process is of endodontic or periodontal origin.

Percussion

Referring back to the patient's chief complaint may indicate the importance of percussion testing for this particular case. If the patient is experiencing acute sensitivity or pain on mastication, this response can typically be duplicated by individually percussing the teeth, which often isolates the symptoms to a particular tooth. Pain to percussion does not indicate that the tooth is vital or nonvital but is rather an indication of inflammation in the periodontal ligament (i.e., symptomatic apical periodontitis). This inflammation may be secondary to physical trauma, occlusal prematurities, periodontal disease, or the extension of pulpal disease into the periodontal ligament space. The indication of where the pain originates is interpreted by the mesencephalic nucleus, receiving its information from proprioceptive nerve receptors. Although subject to debate, the general consensus is that there are relatively few proprioceptors in the dental pulp; however, they are prevalent in the periodontal ligament spaces.²⁴ This is why it may be difficult for the patient to discriminate the location of dental pain in the earlier stages of pathosis, when only the C fibers are stimulated.

Once the disease state extends into the periodontal ligament space, the pain may become more localized for the patient; therefore, the affected tooth will be more identifiable with percussion and mastication testing.

Before percussing any teeth, the clinician should tell the patient what will transpire during this test. Because the presence of acute symptoms may create anxiety and possibly alter the patient's response, properly preparing the patient will lead to more accurate results. The contralateral tooth should first be tested as a control, as should several adjacent teeth that are certain to respond normally. The clinician should advise the patient that the sensation from this tooth is normal and ask to be advised of any tenderness or pain from subsequent teeth.

Percussion is performed by tapping on the incisal or occlusal surfaces of the teeth either with the finger or with a blunt instrument. The testing should initially be done gently, with light pressure being applied digitally with a gloved finger tapping. If the patient cannot detect significant difference between any of the teeth, the test should be repeated using the blunt end of an instrument, like the back end of a mirror handle (Fig. 1.15). The tooth crown is tapped vertically and horizontally. The tooth should first be percussed occlusally, and if the patient discerns no difference, the test should be repeated, percussing the buccal and lingual aspects of the teeth. For any heightened responses, the test should be repeated as necessary to determine that it is accurate and reproducible, and the information should be documented.

Although this test does not disclose the condition of the pulp, it indicates the presence of a periradicular inflammation. An abnormal positive response indicates inflammation of the periodontal ligament that may be of either pulpal or periodontal origin. The sensitivity of the proprioceptive fibers in an inflamed periodontal ligament will help identify the location of the pain. This test should be done gently, especially in highly sensitive teeth. It should be repeated several times and compared with control teeth.

Mobility

Like percussion testing, an increase in tooth mobility is not an indication of pulp vitality. It is merely an indication of a



Fig. 1.15 Percussion testing of a tooth, using the back end of a mirror handle.

compromised periodontal attachment apparatus. This compromise could be the result of acute or chronic physical trauma, occlusal trauma, parafunctional habits, periodontal disease, root fractures, rapid orthodontic movement, or the extension of pulpal disease, specifically an infection, into the periodontal ligament space. Tooth mobility is directly proportional to the integrity of the attachment apparatus or to the extent of inflammation in the periodontal ligament. Often the mobility reverses to normal after the initiating factors are repaired or eliminated. Because determining mobility by simple finger pressure can be visually subjective, the back ends of two mirror handles should be used, one on the buccal aspect and one on the lingual aspect of the tooth (Fig. 1.16). Pressure is applied in a facial-lingual direction as well as in a vertical direction and the tooth mobility is scored (Box 1.2). Any mobility that exceeds +1 should be considered abnormal. However, the teeth should be evaluated on the basis of how mobile they are relative to the adjacent and contralateral teeth.

Periodontal Examination

Periodontal probing is an important part of any intraoral diagnosis. The measurement of periodontal pocket depth is an indication of the depth of the gingival sulcus, which corresponds to the distance between the height of the free gingival margin and the height of the attachment apparatus below. Using a calibrated periodontal probe, the clinician should record the periodontal pocket depths on the mesial, middle, and distal aspects of both the buccal and lingual sides of the tooth, noting the depths in millimeters. The periodontal probe is “stepped” around the long axis of



Fig. 1.16 Mobility testing of a tooth, using the back ends of two mirror handles.

Box 1.2 Recording Tooth Mobility

- +1 *mobility*: The first distinguishable sign of movement greater than normal
- +2 *mobility*: Horizontal tooth movement no greater than 1 mm
- +3 *mobility*: Horizontal tooth movement greater than 1 mm, with or without the visualization of rotation or vertical depressability

the tooth, progressing in 1-mm increments. Periodontal bone loss that is wide, as determined by a wide span of deep periodontal probing, is generally considered to be of periodontal origin and is typically more generalized in other areas of the mouth. However, isolated areas of vertical bone loss may be of an endodontic origin, specifically from a non-vital tooth whose infection has extended from the periapex to the gingival sulcus. Again, proper pulp testing is imperative, not just for the determination of a diagnosis but also for the development of an accurate prognosis assessment. For example, a periodontal pocket of endodontic origin may resolve after endodontic treatment, but if the tooth was originally vital with an associated deep periodontal pocket, endodontic treatment will not improve the periodontal condition. In addition, as discussed in Chapter 22, a vertical root fracture may often cause a localized narrow periodontal pocket that extends deep down the root surface. Characteristically, the adjacent periodontium is usually within normal limits.

Furcation bone loss can be secondary to periodontal or pulpal disease. The amount of furcation bone loss, as observed both clinically and radiographically, should be documented (Box 1.3). Results of pulp tests (described later) will aid in diagnosis.

PULP TESTS

Pulp test (pulp sensibility test) is a diagnostic procedure to determine pulp status. It can be performed with electrical, mechanical, or thermal stimuli, or by the assessment of the blood supply to the tooth.⁶ It involves attempting to make a determination of the responsiveness of pulpal sensory neurons.^{62,63} It aims to obtain a subjective response from the patient (i.e., to determine whether the pulpal nerves are functional), or the tests may involve a more objective approach using devices that detect the integrity of the pulpal vasculature. Unfortunately, the quantitative evaluation of the status of pulp tissue can only be determined histologically, as it has been shown that there is not necessarily a good correlation between the objective clinical signs and symptoms and the pulpal histology.^{122,123}

Thermal

Various methods and materials have been used to test the pulp's response to thermal stimuli. The baseline or normal response to either cold or hot is a patient's report that a sensation is felt but disappears immediately upon removal of the thermal stimulus. Abnormal responses include a lack of response to the stimulus, a lingering or intensification of a painful sensation after the stimulus is removed, or an immediate, excruciatingly painful sensation as soon as the stimulus is placed on the tooth.

Box 1.3 Recording Furcation Defects

- Class I furcation defect*: The furcation can be probed but not to a significant depth.
- Class II furcation defect*: The furcation can be entered into but cannot be probed completely through to the opposite side.
- Class III furcation defect*: The furcation can be probed completely through to the opposite side.

Cold testing is the primary pulp testing method used by many clinicians today. It is especially useful for patients presenting with porcelain jacket crowns or porcelain-fused-to-metal crowns where no natural tooth surface (or much metal) is accessible. If a clinician chooses to perform this test with sticks of ice, then the use of a rubber dam is recommended, because melting ice will run onto adjacent teeth and gingiva, yielding potentially false-positive responses.

Frozen carbon dioxide (CO_2), also known as *dry ice* or *carbon dioxide snow*, or CO_2 stick, has been found to be reliable in eliciting a positive response if vital pulp tissue is present in the tooth.^{46,98,99} One study found that vital teeth would respond to both frozen CO_2 and skin refrigerant, with skin refrigerant producing a slightly quicker response.⁶⁶ Frozen carbon dioxide has also been found to be effective in evaluating the pulpal response in teeth with full coverage crowns for which other tests such as electric pulp testing is not possible.¹¹ For testing purposes, a solid stick of CO_2 is prepared by delivering CO_2 gas into a specially designed plastic cylinder (Fig. 1.17). The resulting CO_2 stick is applied to the facial surface of either the natural tooth structure or crown. Several teeth can be tested with a single CO_2 stick. The teeth should be isolated and the oral soft tissues should be protected with a 2-by-2-inch gauze or

cotton roll so the frozen CO_2 will not come into contact with these structures. Because of the extremely cold temperature of the frozen CO_2 (-69°F to -119°F ; -56°C to -98°C), burns of the soft tissues can occur. It has been demonstrated on extracted teeth that frozen CO_2 application has resulted in a significantly greater intrapulpal temperature decrease than either skin refrigerant or ice.¹¹ Also, it appears that the application of CO_2 to teeth does not result in any irreversible damage to the pulp tissues or cause any significant enamel crazing.^{61,104}

The most popular method of performing cold testing is with a refrigerant spray. It is readily available, easy to use, and provides test results that are reproducible, reliable, and equivalent to that of frozen CO_2 .^{46,66,96,141} One of the current products contains 1,1,1,2-tetrafluoroethane, which has zero ozone depletion potential and is environmentally safe. It has a temperature of -26.2°C .⁶⁶ The spray is most effective for testing purposes when it is applied to the tooth on a large #2 cotton pellet (Fig. 1.18). In one study,⁶⁵ a significantly lower intrapulpal temperature was achieved when a #2 cotton pellet was dipped or sprayed with the refrigerant compared with the result when a small #4 cotton pellet or cotton applicator was used. The sprayed cotton pellet should be applied to the midfacial area of the tooth or crown. As with any other pulp testing method, adjacent or contralateral “normal” teeth should also be tested to establish a baseline response. It appears that frozen CO_2 and refrigerant spray are superior to other cold testing methods and equivalent or superior to the electric pulp tester for assessing pulp vitality.^{11,46} However, one study found that periodontal attachment loss and gingival recession may influence the reported pain response with cold stimuli.¹¹⁶

To be most reliable, cold testing should be used in conjunction with an electric pulp tester (described later in this chapter) so that the results from one test will verify the findings of the other test. If a mature, nontraumatized tooth does not respond to both cold testing and electric pulp testing, then the pulp can be considered necrotic.^{23,98,141} However, a multirrooted tooth, with at least one root containing vital pulp tissue, may respond to a cold test and electric pulp test even if one or more of the roots contain necrotic pulp tissue.⁹⁸

Another thermal testing method involves the use of heat. Heat testing is most useful when a patient's chief complaint is intense dental pain on contact with any hot liquid or food. When a patient is unable to identify which tooth is sensitive, a heat test is appropriate. Starting with the most posterior tooth in that area of the mouth, each tooth is individually isolated with a dental dam. An irrigating syringe is filled with a liquid (most commonly plain water) that has a temperature similar to that which would cause the painful sensation. The liquid is then expressed from the syringe onto the isolated tooth to determine whether the response is normal or abnormal. The clinician moves forward in the quadrant, isolating each individual tooth until the offending tooth is located. That tooth will exhibit an immediate, intense painful response to the heat. With heat testing, a delayed response may occur, so waiting 10 seconds between each heat test will allow sufficient time for the onset of symptoms. This method can also be used to apply cold water to the entire crown for cases in which cold is the precipitating stimulus.




Fig. 1.17 **A**, Carbon dioxide tank with apparatus attached to form solid CO_2 stick/pencil. **B**, CO_2 gas being transformed into a solid stick/pencil. **C**, CO_2 stick/pencil extruded from end of a plastic carrier and ready for use.




Fig. 1.18 **A**, Refrigerant spray container. **B**, A large cotton pellet made of a cotton roll, or a ready-made size #2 (large) cotton pellet, can be used to apply the refrigerant spray to the tooth surface. The small #4 cotton pellet does not provide as much surface area as the #2 cotton pellet, and therefore should not be used to deliver the refrigerant to the tooth surface. **C**, A large cotton pellet sprayed with the refrigerant and ready to be applied to the tooth surface. (A, Courtesy Coltène/Whaledent, Cuyahoga Falls, OH.)

Another method for heat testing is to apply heated gutta-percha or compound stick to the surface of the tooth. If this method is used, a light layer of lubricant should be placed onto the tooth surface before applying the heated material to prevent the hot gutta-percha or compound from adhering to the dry tooth surface. Heat can also be generated by the friction created when a dry rubber-polishing wheel is run at a high speed against the dry surface of a tooth. However, this latter method is not recommended. Another approach is the use of electronic heat-testing instruments.²⁰

If the heat test confirms the results of other pulp testing procedures, emergency care can then be provided. Often a tooth that is sensitive to heat may also be responsible for some spontaneous pain. The patient may present with cold liquids in hand just to minimize the pain (Fig. 1.19). In such cases, the application of cold to a specific tooth may eliminate the pain and greatly assist in the diagnosis. Typically, a tooth that responds to heat and then is relieved by cold is found to be necrotic.

Electric

Assessment of pulp neural responses (*sensibility*) can also be accomplished by electric pulp testing.⁷⁹ Electric pulp testers of different designs and manufacturers have been used for this purpose. Electric pulp testers should be an integral part of any dental practice. It should be noted that the sensibility of the pulp is determined by the intactness and health of the vascular supply, not by the status of the pulpal nerve fibers. Even though advances are being made with regard to determining the status of the pulp on the basis of




Fig. 1.19 Irreversible pulpitis associated with the mandibular right second molar. Patient has found that the only way to alleviate the pain is to place a jar filled with ice water against the right side of his face.

the blood supply, this technology has not been perfected enough at this time to be used on a routine basis in a clinical setting.

The electric pulp tester has some limitations in providing predictable information about the status of the pulp. The

Box 1.4 Potential Common Interpretation Errors of Responses Obtained From Electric Pulp Testing

False-Positive Responses

Partial pulp necrosis
Patient's high anxiety
Ineffective tooth isolation
Contact with metal restorations

False-Negative Responses

Calcific obliterations in the root canals
Recently traumatized teeth
Immature apex
Drugs that increase patient's threshold for pain
Poor contact of pulp tester to tooth

response of the pulp to electric testing does not reflect the histologic health or disease status of the pulp.^{122,123} A response by the pulp to the electric current only denotes that some viable nerve fibers are present in the pulp and are capable of responding. Numeric readings on the pulp tester have significance only if the number differs significantly from the readings obtained from a control tooth tested on the same patient with the electrode positioned at a similar area on both teeth. However, in most cases, the response is scored as either present or absent. Studies^{122,123} have shown that electric pulp test results are most accurate when no response is obtained to any amount of electric current. This lack of response has been found most frequently when a necrotic pulp is present. In addition, false-positive and false-negative responses can occur (Box 1.4), and the clinician must take it into account when formulating the final diagnosis.

The electric pulp tester will not work unless the probe can be placed in contact with or be bridged to the natural tooth structure.⁹⁵ With the advent of universal precautions for infection control, the use of rubber gloves prevents the clinician from completing the circuit.⁷ Some pulp testers may require the patient to place a finger, or fingers, on the tester probe to complete the electric circuit; however, the use of lip clips is an alternative to having patients hold the tester. Proper use of the electric pulp tester requires the evaluated teeth to be carefully isolated and dried. A control tooth of similar tooth type and location in the arch should be tested first in order to establish a baseline response and to inform the patient as to what a "normal" sensation is. The suspected tooth should be tested at least twice to confirm the results. The tip of the testing probe that will be placed in contact with the tooth structure must be coated with a water- or petroleum-based medium.⁸⁶ The most commonly used medium is toothpaste. The coated probe tip is placed in the incisal third of the facial or buccal area of the tooth to be tested.¹⁵ Once the probe is in contact with the tooth, the patient is asked to touch or grasp the tester probe, unless a lip clip is used (Fig. 1.20, A). This completes the circuit and initiates the delivery of an electric current to the tooth. The patient is instructed to remove his or her finger(s) from the probe when a "tingling" or "warming" sensation is felt in the tooth. The readings from the pulp tester are recorded (Fig. 1.20, B) and will be evaluated once all the appropriate




Fig. 1.20 A, Electric pulp tester with probe. The probe tip will be coated with a conductive medium, such as toothpaste, and placed in contact with the tooth surface. The patient will activate the unit by placing a finger on the metal shaft of the probe. B, View of the electric pulp tester control panel; the knob on the front right of the unit controls the rate at which the electric current is delivered to the tooth. The plastic panel on the left front displays the digital numerical reading obtained from the pulp test. The digital scale runs from 0 to 80. (Courtesy SybronEndo, Orange, CA.)

teeth have been tested by the electric pulp tester and the other pulp testing methods.

If a complete coverage crown or extensive restoration is present, a bridging technique can be attempted to deliver the electric current to any exposed natural tooth structure.⁹⁵ The tip of an endodontic explorer is coated with toothpaste or other appropriate medium and placed in contact with the natural tooth structure. The tip of the electric pulp tester probe is coated with a small amount of toothpaste and placed in contact with the side of the explorer. The patient completes the circuit and the testing proceeds as described previously. If no natural tooth structure is available, then an alternative pulp testing method, such as cold, should be used.

One study compared the ability of thermal and electric pulp testing methods to register the presence of vital pulp tissue.⁹⁹ The *sensibility*, which is the ability of a test to identify teeth that are diseased, was 0.83 for the cold test, 0.86 for heat test, and 0.72 for the electric test. This means the cold test correctly identified 83% of the teeth that had a necrotic pulp, whereas heat tests were correct 86% of the time and electric pulp tests were correct only 72% of the

time. This same study evaluated the *specificity* of these three tests. Specificity relates to the ability of a test to identify teeth without disease. Ninety-three percent of teeth with healthy pulps were correctly identified by both the cold and electric pulp tests, whereas only 41% of the teeth with healthy pulps were identified correctly by the heat test. From the results of the testing, it was found that the cold test had an accuracy of 86%, the electric pulp test 81%, and the heat test 71%.

Some studies have indicated there might not be a significant difference between pulp testing results obtained by electric pulp tester and those obtained by the thermal methods.^{46,98,99} However, cold tests have been shown to be more reliable than electric pulp tests in younger patients with less developed root apices.^{5,42,98} This is the reason to verify the results obtained by one testing method and compare them with results obtained by other methods. Until such time that the testing methods used to assess the vascular supply of the pulp become less time consuming and technique sensitive, thermal and electric pulp testing will continue to be the primary methods for determining pulp sensibility.

Laser Doppler Flowmetry

Laser Doppler flowmetry (LDF) is a method used to assess blood flow in microvascular systems. Attempts are being made to adapt this technology to assess pulpal blood flow. A diode is used to project an infrared light beam through the crown and pulp chamber of a tooth. The infrared light beam is scattered as it passes through the pulp tissue. The Doppler principle states that the light beam's frequency will shift when hitting moving red blood cells but will remain unshifted as it passes through static tissue. The average Doppler frequency shift will measure the velocity at which the red blood cells are moving.¹¹⁴

Several studies^{40,60,69,84,114,115,117} have found LDF to be an accurate, reliable, and reproducible method of assessing pulpal blood flow. One of the great advantages of pulp testing with devices such as the LDF is that the collected data are based on objective findings rather than subjective patient responses. As is discussed in Chapter 21, certain luxation injuries will cause inaccuracies in the results of electric and thermal pulp testing. LDF has been shown to be a great indicator for pulpal vitality in these cases.¹³⁰ However, this technology is not being used routinely in the dental practice.

Pulse Oximetry

The pulse oximeter is another noninvasive device (Fig. 1.21). Widely used in medicine, it is designed to measure the oxygen concentration in the blood and the pulse rate. A pulse oximeter works by transmitting two wavelengths of light, red and infrared, through a translucent portion of a patient's body (e.g., a finger, earlobe, or tooth). Some of the light is absorbed as it passes through the tissue; the amount absorbed depends on the ratio of oxygenated to deoxygenated hemoglobin in the blood. On the opposite side of the targeted tissue, a sensor detects the absorbed light. On the basis of the difference between the light emitted and the light received, a microprocessor calculates the pulse rate and oxygen concentration in the blood.¹¹⁸ The transmission of light to the sensor requires that there be no obstruction from restorations, which can sometimes limit the usefulness of pulse oximetry to test the status of the pulp tissue.




Fig. 1.21 Nellcor OxiMax N-600x pulse oximeter. (Courtesy Nellcor Puritan Bennett, Boulder, CO; now part of Covidien.)

Custom-made sensors have been developed and were found to be more accurate than electric and thermal pulp tests.^{31,54} This sensor has been especially useful in evaluating teeth that have been subjected to traumatic injuries, as such teeth tend to present, especially in the short term, with questionable neural response using conventional pulp testing methods.^{8,31,53}

Studies regarding the ability of pulse oximetry to diagnose pulp health have drawn various conclusions. Several studies have found pulse oximetry to be a reliable method for assessing pulp health.^{69,70,118,125,140} Others have stated that in its present form the pulse oximeter may not be predictable for such diagnoses.¹⁴⁰ Most of the problems appear to be related to the currently available technology. Some investigators have concluded that the devices used for pulp testing are too cumbersome and complicated to be used on a routine basis in a dental practice.^{68,118,140}

SPECIAL TESTS

Bite Test

Bite tests and percussion tests are indicated when a patient presents with pain while biting. On occasion, the patient may not know which tooth is sensitive to biting pressure, and percussion and bite tests may help to localize the tooth involved. The tooth may be sensitive to biting when the pulpal pathosis has extended into the periodontal ligament space, creating a *symptomatic apical periodontitis*, or the sensitivity may be present secondary to a crack in the tooth. The clinician can often differentiate between periradicular periodontitis and a cracked tooth or fractured cusp. If periradicular periodontitis is present, the tooth will respond with pain to percussion and biting tests regardless of where the pressure is applied to the coronal part of the tooth. A cracked tooth or fractured cusp will typically elicit pain only when the bite or percussion test is applied in a certain direction to one cusp or section of the tooth.^{22,108}

For the bite test to be meaningful, a device should be used that allows the clinician to apply pressure to individual cusps or areas of the tooth. A variety of devices have been used for bite tests, including cotton tip applicators, toothpicks, orangewood sticks, and rubber polishing wheels. There are several devices specifically designed to perform this test. The Tooth Slooth (Professional Results, Laguna Niguel, CA) (Fig. 1.22) and FracFinder (Hu-Friedy, Oakbrook, IL) are just two of the commercially available devices



Fig. 1.22 To determine which tooth, or tooth part, is sensitive to mastication, having the patient bite on a specially designed bite stick is often helpful.

used for the bite test. As with all pulp tests, adjacent and contralateral teeth should be used as controls so that the patient is aware of the “normal” response to these tests. The small cupped-out area on these instruments is placed in contact with the cusp to be tested. The patient is then asked to apply biting pressure with the opposing teeth to the flat surface on the opposite side of the device. The biting pressure should be applied slowly until full closure is achieved. Firm pressure should be applied for a few seconds; the patient is then asked to release the pressure quickly. Each individual cusp on a tooth can be tested in a similar manner. The clinician should note whether the pain is elicited during the pressure phase or on quick release of the pressure. A common finding with a fractured cusp or cracked tooth is the frequent presence of pain upon release of biting pressure.

Test Cavity

The test cavity method for assessing pulpal response is not routinely used since, by definition, it is an invasive irreversible test. This method is used only when all other test methods are deemed impossible or the results of the other tests are inconclusive. An example of a situation in which this method can be used is when the tooth suspected of having pulpal disease has a full coverage crown. If no sound tooth structure is available to use a bridging technique with the electric pulp tester, and the cold test results are inconclusive, a small class I cavity preparation is made through the occlusal surface of the crown. This is accomplished with a high-speed #1 or #2 round bur with proper air and water coolant. The patient is not anesthetized while this procedure is performed, and the patient is asked to respond if any painful sensation is felt during the drilling procedure. If the patient feels pain once the bur contacts sound dentin, the procedure is terminated and the class I cavity preparation is restored. This sensation signifies only that there is some viable nerve tissue remaining in the pulp, not that the pulp is totally healthy. If the patient fails to feel any sensation when the bur reaches the dentin, this is a good indication that the pulp is necrotic and root canal therapy is indicated.

The patient should be given a full explanation and reassurance of the procedure before the test cavity is attempted.

Staining and Transillumination

To determine the presence of a crack in the surface of a tooth, the application of a stain to the area is often of great assistance. It may be necessary to remove the restoration in the tooth to better visualize a crack or fracture. Methylene blue dye, when painted on the tooth surface with a cotton tip applicator, will penetrate into cracked areas. The excess dye may be removed with a moist application of 70% isopropyl alcohol. The remaining dye will indicate the possible location of the crack.

Transillumination using a bright fiberoptic light probe to the surface of the tooth may be very helpful (Fig. 1.23). Directing a high-intensity light directly on the exterior surface of the tooth at the cementum-enamel junction (CEJ) may reveal the extent of the fracture. Teeth with fractures block transilluminated light. The part of the tooth that is proximal to the light source will absorb this light and glow, whereas the area beyond this fracture will not have light transmitted to it and will show dark by comparison.¹⁰¹ Although the presence of a fracture may be evident using dyes and transillumination, the full extent of the fracture cannot always be determined by these tests alone.

Selective Anesthesia

When symptoms are not localized or referred, the diagnosis may be challenging. Sometimes the patient may not even be able to specify whether the symptoms are emanating from the maxillary or mandibular arch. In these instances, when pulp testing is inconclusive, *selective anesthesia* may be helpful.

If the patient cannot determine which arch the pain is coming from, then the clinician should first selectively anesthetize the maxillary arch. This should be accomplished by using a periodontal ligament (intraalveolar) injection. The injection is administered to the most posterior tooth in the quadrant of the arch that may be suspected, starting from the distal sulcus. The anesthesia is subsequently administered in an anterior direction, one tooth at a time, until the pain is eliminated. If the pain is not eliminated after an appropriate period of time, then the clinician should similarly repeat this technique on the mandibular teeth below. It should be understood that periodontal ligament injections may anesthetize an adjacent tooth and thus are more useful for identifying the arch rather than the specific tooth.

RADIOGRAPHIC EXAMINATION AND INTERPRETATION

Intraoral Radiographs

The radiographic interpretation of a potential endodontic pathosis is an integral part of endodontic diagnosis and prognosis assessment. Few diagnostic tests provide as much useful information as dental radiography. For this reason, the clinician is sometimes tempted to prematurely make a definitive diagnosis based solely on radiographic interpretation. However, the image should be used only as one sign, providing important clues in the diagnostic investigation. When not coupled with a proper history and clinical examination




Fig. 1.23 Sometimes there is no clear indication of why a tooth is symptomatic. This radiograph shows a mandibular second molar with a moderately deep restoration (**A**); the pulp tests nonvital. Without any transillumination, a fracture cannot be detected (**B**). However, by placing a high-intensity light source on the tooth surface, a root fracture can be observed on the buccal surface (**C**) and the distal-lingual surface (**D**).

and testing, the radiograph alone can lead to a misinterpretation of normality and pathosis (Fig. 1.24). Because treatment planning will ultimately be based on the diagnosis, the potential for inappropriate treatment may frequently exist if the radiograph alone is used for making final diagnosis. The clinician should not subject the patient to unnecessary multiple radiation exposures; two pretreatment images from different angulations are often sufficient. However, under extenuating circumstances—especially when the diagnosis is difficult—additional exposures may be necessary to determine the presence of multiple roots, multiple canals, resorptive defects, caries, restoration defects, root fractures, and the extent of root maturation and apical development.

The radiographic appearance of endodontic pathosis can sometimes be highly subjective. In a study by Goldman and colleagues, there was only 50% agreement among interpreters for the radiographic presence of pathosis.⁴⁹ When the cases were reevaluated several months later, the same evaluators agreed with their own original diagnosis less than 85% of the time.⁵⁰ This further emphasizes the necessity for additional objective diagnostic tests, as well as the importance of obtaining and comparing older radiographs of the same area of the patient.

For standard two-dimensional radiography, clinicians basically project x-radiation through an object and capture the image on a recording medium, either x-ray film or a digital sensor. Much like casting a shadow from a light source, the image appearance may vary greatly depending on how the radiographic source is directed. Thus, the three-dimensional interpretation of the resulting two-dimensional image requires not only knowledge of normality and pathosis but also advanced knowledge of how the radiograph was exposed. By virtue of “casting a shadow,” the anatomic features that are closest to the film (or sensor) will move the least when there is a change in the horizontal or vertical angulation of the radiation source (Fig. 1.25). This may be helpful in determining the existence of additional roots, the location of pathosis, and the unmasking of anatomic structures. Changes in the horizontal or vertical angulation may help elucidate valuable anatomic and pathologic information; it also has the potential to hide important information. An incorrect vertical angulation may cause the buccal roots of a maxillary molar to be masked by the zygomatic arch. An incorrect horizontal angulation may cause roots to overlap with the roots of adjacent teeth, or it may incorrectly create



Fig. 1.24 Radiograph showing what appears to be a mandibular lateral incisor associated with periapical lesion of a nonvital tooth. Although pulp necrosis can be suspected, the tooth tested vital. In this case, the appearance of apical bone loss is secondary to a cementoma.

the appearance of a one-rooted tooth, when two roots are actually present.

In general, when endodontic pathosis appears radiographically, it appears as a radiolucency in the area of the periapex. The pathosis may present merely as a widening or break in the lamina dura—the most consistent radiographic finding when a tooth is nonvital⁶⁷—or it may present as a radiolucent area at the apex of the root or in the alveolar bone adjacent to the exit of a lateral or furcation accessory canal. On occasion no radiographic change can be seen at all, even in the presence of a disease process in the alveolar bone. This is mainly due to the fact that the disease process did not reach the cortical plate of the bone.

Two-dimensional dental radiography has two basic shortcomings: the lack of early detection of pathosis in the cancellous bone, because of the density of the cortical plates, and the influence of the superimposition of anatomic structures. Variability in the radiographic expression of an osseous pathosis has much to do with the relative location of the root of the tooth and how it is oriented with respect to the cortical and cancellous bone. Radiographic changes from bone loss will not be detected if the loss is only in cancellous bone.¹⁶ However, the radiographic evidence of pathosis will be observed once this bone loss extends to the junction of the cortical and cancellous bone. In addition, certain teeth are more prone to exhibit radiographic changes than others, depending on their anatomic location.¹⁷ The radiographic appearance of endodontic pathosis




Fig. 1.25 Radiographic images are only two-dimensional, and often it is difficult to discriminate the relative location of overlapping objects. **A**, When the source of radiation is directly perpendicular to overlapping objects, the image is captured without much separation of the objects. However, when the radiation source is at an angle to offset the overlapping objects, the image is captured with the objects being viewed as separated. **B**, The object that is closest to the film (or sensor) will move the least, with the object closest to the radiation source appearing farthest away.

is correlated with the relationship of the periapex of the tooth and its juxtaposition to the cortical-cancellous bone junction. The apices of most anterior and premolar teeth are located close to the cortical-cancellous bone junction. Therefore, periapical pathosis from these teeth is exhibited sooner on the radiograph. By comparison, the distal roots of mandibular first molars and both roots of mandibular second molars are generally positioned more centrally within the cancellous bone, as are maxillary molars, especially the palatal roots. Periapical lesions from these roots must expand more before they reach the cortical-cancellous bone junction and are recognized as radiographic pathosis. For these reasons, it is important not to exclude the possibility of pulpal pathosis in situations in which there are no radiographic changes.

Many factors can influence the quality of the radiographic interpretation, including the ability of the person exposing the radiograph, the quality of the radiographic film, the quality of the exposure source, the quality of the film processing, and the skill with which the film is viewed. Controlling all of these variables can be a difficult challenge but is paramount for obtaining an accurate radiographic interpretation.

Digital Radiography

Digital radiography has been available since the late 1980s and has recently been refined and popularized with better hardware and a more user-friendly interface. It has the

ability to capture, view, magnify, enhance, and store radiographic images in an easily reproducible format that does not degrade over time. Significant advantages of digital radiographs over conventional radiographs include lower radiation doses, instant viewing, convenient manipulation, efficient transmission of an image via the Internet, simple duplication, and easy archiving.

Digital radiography uses no x-ray film and requires no chemical processing. Instead, a *sensor* is used to capture the image created by the radiation source. This sensor is either directly or wirelessly attached to a local computer, which interprets this signal and, using specialized software, translates the signal into a two-dimensional digital image that can be displayed, enhanced, and analyzed. The image is stored in the patient's file, typically in a dedicated network server, and can be recalled as needed. Further information about digital radiography may be found in Chapter 2.

The viewing of a digital radiographic image on a high-resolution monitor allows for rapid and easy interpretation for both the clinician and the patient. The image appears almost instantly, with no potential for image distortion from improper chemical processing. The clinician can magnify different areas on the radiograph and then digitally enhance the image in order to better visualize certain anatomic structures; in some cases, the image can even be colorized, which is a useful tool for patient education (Fig. 1.26).

In the past, x-ray film has had a slightly better resolution than most digital radiography images, at about 16 line pairs per millimeter (lp/mm).⁸⁷ However, current sensors offer resolutions beyond that of conventional film. Under the best of circumstances, the human eye can see only about 10 lp/mm, which is the lowest resolution for most dental digital radiography systems. Digital sensors are much more sensitive to radiation than conventional x-ray film and thus require 50% to 90% less radiation in order to acquire an image, an important feature for generating greater patient acceptance of dental radiographs.




Fig. 1.26 Digital radiography has an advantage over conventional film in that the image can be enhanced and colorized—a useful tool for patient education.

The diagnostic quality of this technology has been shown to be comparable to conventional film-based radiography.^{39,73,97} The interpretation of a digital radiograph can be subjective, similar to that of the conventional film.¹³⁴ Factors that appear to have the most impact on the interpretation of the image are the years of experience of the examiner and familiarity of the operator with the given digital system.¹³⁴

Cone-Beam Computerized Tomography

Limitations in conventional two-dimensional radiography promulgated a need for three-dimensional imaging, known as *cone-beam computerized tomography* (CBCT) (also known as *cone-beam volumetric tomography* [CBVT]) or as *cone-beam volumetric imaging* [CBVI]. Although a form of this technology has existed since the early 1980s,¹⁰⁶ specific devices for dental use first appeared almost two decades later.⁹⁰ Most of these machines are similar to a dental panoramic radiographic device, whereby the patient stands or sits as a cone-shaped radiographic beam is directed to the target area with a reciprocating capturing sensor on the opposite side (Fig. 1.27). The resulting information is digitally reconstructed and interpreted to create an interface whereby the clinician can three-dimensionally interpret “slices” of the patient's tissues in a multitude of planes (Figs. 1.28 and 1.29).^{37,33} The survey of the scans can be interpreted immediately after the scan. Various software applications have been used to enable the images to be sent to other clinicians. This is accomplished either in printed format or with portable and transferable software that can be used interactively by another clinician.

In general, many dental applications only require a limited field of vision, confining the study to the maxilla and mandible. However, many devices have the ability to



Fig. 1.27 Cone-beam volumetric tomography, using the 3D Accutomo 80. (Courtesy J. Morita USA, Irvine, CA.)



Fig. 1.28 Cone-beam volumetric tomography has the ability to capture, store, and present radiographic images in various horizontal and vertical planes. (Courtesy J. Morita USA, Irvine, CA)




Fig. 1.29 Cone-beam volumetric tomography has the advantage of being able to detect pathosis in the bone or associated with the teeth without the obstruction of anatomic structures. The planes of vision may be axial, sagittal, or coronal. (Courtesy J. Morita USA, Irvine, CA)

provide a full field of vision for viewing more regional structures. Clinicians should thoroughly understand the ethical and medical-legal ramifications of doing scans with full fields of view. Incidental nondental findings have been seen from these scans, such as intracranial aneurysms, which, when undetected, could be life threatening.⁹¹

The radiation source of CBCT is different from that of conventional two-dimensional dental imaging in that the radiation beam created is conical in shape. Also, conventional digital dental radiography is captured and interpreted as *pixels*, a series of dots that collectively produces an image of the scanned structure. For CBCT, the image is instead captured as a series of three-dimensional pixels, known as *voxels*. Combining these voxels gives a three-dimensional image that can be “sliced” into various planes, allowing for specific evaluations without a necropsy (Fig. 1.30). One of the advantages of using a device that has a limited field of vision is that the voxel size can be less than half that of a device using a full field of vision, thereby increasing the resolution of the resulting image




Fig. 1.30 The radiation source in cone-beam volumetric tomography is conical. The receiving sensor captures the image as “voxels,” or three-dimensional pixels of information, allowing digital interpretation.

and providing for a more accurate interpretation of anatomic structures and pathologic conditions. The development of limited field of vision devices has also contributed to decreasing the radiation and cost of these machines, making them more practical for endodontic use.⁴¹

Compared with two-dimensional radiographs, CBCT can clearly visualize the interior of the cancellous bone without the superimposition of the cortical bone. Studies show that CBCT is much more predictable and efficient in demonstrating anatomic landmarks, bone density, bone loss, periapical lesions, root fractures, root perforations, and root resorptions.^{1,21,26,27,38,47,71,78,81,85,92,94,128,131,142}

The superimposition of anatomic structures can also mask the interpretation of alveolar defects. Specifically, the maxillary sinus, zygoma, incisive canal and foramen, nasal bone, orbit, mandibular oblique ridge, mental foramen, mandibular mentalis, sublingual salivary glands, tori, and the overlap of adjacent roots may either obscure bone loss or mimic bone loss, making an accurate interpretation of conventional radiography sometimes difficult or impossible.

Several studies have demonstrated the advantages of CBCT in the differential diagnosis of such structures from pathologic conditions.^{21,29,71,137}

CBCT should not be seen as a replacement for conventional dental radiography, but rather as a diagnostic adjunct. The advantage of conventional dental radiography is that it can visualize most of the structures in one image. CBCT can show great detail in many planes of vision but can also leave out important details if the “slice” is not in the area of existing pathosis (Fig. 1.31). There is a promising future for the use of CBCT for endodontic diagnosis and treatment. It has already proven invaluable in the detection of dental and nondental pathoses (Fig. 1.32). For a further review of CBCT and radiography, see Chapter 2.

Magnetic Resonance Imaging

Magnetic resonance imaging (MRI) has also been suggested for dental diagnosis. It may offer simultaneous three-dimensional hard- and soft-tissue imaging of teeth without ionizing radiation.⁵⁸ The use of MRI in endodontics is still limited.

CRACKS AND FRACTURES

The wide variety of types of cracks and fractures in teeth and their associated signs and symptoms often make their diagnosis difficult. The extensiveness of the crack or fracture

line may directly alter the prognosis assessment for a given tooth and should be examined before treatment decision making. Certain types of cracks may be as innocent as a superficial enamel craze line, or they may be as prominent as a fractured cusp. The crack may progress into the root system to involve the pulp, or it may split the entire tooth into two separate segments. The crack may be oblique, extending cervically, such that once the coronal segment is removed the tooth may or may not be restorable. Any of these situations may present with mild, moderate, or severe symptoms or possibly no symptoms at all.

Crack Types

There have been many suggestions in the literature of how to classify cracks in teeth. By defining the type of crack present, an assessment of the prognosis may be determined and treatment alternatives can be planned. Unfortunately, it is often extremely difficult to determine how extensive a crack is until the tooth is extracted.

Cracks in teeth can be divided into three basic categories:

- Craze lines
- Fractures (also referred to as *cracks*)
- Split tooth/roots

Craze lines are merely cracks in the enamel that do not extend into the dentin and either occur naturally or develop after trauma. They are more prevalent in adult teeth and




Fig. 1.31 **A**, This standard two-dimensional radiographic image reveals recurrent caries under the mesial margin of the maxillary first molar. However, the patient localized pain to mastication on the maxillary second molar. **B**, Cone-beam volumetric tomography revealed an apical radiolucency associated with the maxillary second molar. The bone loss was obscured on the two-dimensional radiograph by the maxillary sinus, zygoma, and cortical bone.




Fig. 1.32 **A**, Periapical radiograph showing a large apical radiolucency associated with the mandibular second molar. Apical pathosis should be ruled out. **B**, Cone-beam volumetric tomography revealed salivary indentation of the mandible in the area apical and lingual to the mandibular second molar, consistent with a Stafne defect. **C**, Enlargement of coronal section in the area of the mandibular second molar and the Stafne defect located on the lingual aspect of the mandible.

usually occur more in the posterior teeth. If light is transilluminated through the crown of such a tooth, these craze lines may show up as fine lines in the enamel with light being able to transmit through them, indicating that the crack is only superficial. The use of optical coherence tomography (OCT) has also been suggested for the detection

of enamel cracks.⁵⁹ Craze lines typically will not manifest with symptoms. No treatment is necessary for craze lines unless they create a cosmetic issue.

Fractures extend deeper into the dentin than superficial craze lines and primarily extend mesially to distally, involving the marginal ridges. Dyes and transillumination are helpful for visualizing potential root fractures.

Symptoms from a fractured tooth range from none to severe pain. A fracture in the tooth does not necessarily indicate that the tooth has split into two pieces; however, left alone or especially with provocations such as occlusal prematurities, the fracture may progress into a split root. A fractured tooth may be treated by a simple restoration, endodontics (nonsurgical or surgical), or even extraction, depending on the extent and orientation of the fracture, the degree of symptoms, and whether the symptoms can be eliminated. This makes the clinical management of fractured teeth difficult and sometimes unpredictable.

A definitive combination of factors, signs, and symptoms that, when collectively observed, allows the clinician to conclude the existence of a specific disease state is termed a *syndrome*. However, given the multitude of signs and symptoms that fractured roots can present with, it is often difficult to achieve an objective definitive diagnosis. For this reason, the terminology of *cracked tooth syndrome*^{22,108} should be avoided.⁶ The subjective and objective factors seen in cases of fractured teeth will generally be diverse; therefore, a tentative diagnosis of a fractured tooth will most likely be more of a prediction. Once this prediction is made, the patient must be properly informed as to any potential decrease in prognosis of the pending dental treatment. Because treatment options for repairing fractured teeth have only a limited degree of success, early detection and prevention, and proper informed consent, are crucial.^{9,10,72,119,120,124,132}

Split tooth/roots occur when a fracture extends from one separating them into two segments. If the split is more oblique, it is possible that once the smaller separated segment is removed, the tooth might still be restorable—for example, a fractured cusp. However, if the split extends below the osseous level, the tooth may not be restorable and endodontic treatment may not result in a favorable prognosis.

Proper prognosis assessment is imperative before any dental treatment but is often difficult in cases of cracked teeth. Because of the questionable long-term success from treating cases of suspected or known fractures, the clinician should be cautious in making the decision to continue with treatment and should avoid endodontic treatment in cases of a definitive diagnosis of a split tooth or root.

Vertical Root Fractures

One of the more common reasons for recurrent endodontic pathosis is the *vertical root fracture*, a severe crack in the tooth that extends longitudinally down the long axis of the root (Figs. 1.33 and 1.34). Often it extends through the pulp and to the periodontium. It tends to be more centrally located within the tooth, as opposed to being more oblique, and typically traverses through the marginal ridges. These fractures may be present before endodontic treatment, secondary to endodontic treatment, or may develop after endodontic treatment has been completed. Because diagnosing




Fig. 1.33 Poorly fitting intracoronal restorations can place stresses within the tooth that can cause a vertical root fracture. **A**, This radiograph of a mandibular second premolar (with a gold inlay) reveals extensive periapical and periradicular bone loss, especially on the distal aspect. **B**, The tooth pulp tested nonvital, and there was an associated 12-mm-deep, narrow, isolated periodontal pocket on the buccal aspect of the tooth. After the tooth was extracted, the distal aspect was examined. **C**, On magnification ($\times 16$) the distal aspect of the root revealed an oblique vertical root fracture. Similarly, the placement of an ill-fitting post may exert intraradicular stresses on a root that can cause a fracture to occur vertically. **D**, This radiograph depicts a symmetrical space between the obturation and the canal wall, suggesting a vertical root fracture. **E**, After the tooth is extracted, the root fracture can be easily observed.

these vertical root fractures may be difficult, they often go unrecognized. Therefore, diagnosing the existence and extent of a vertical root fracture is imperative before any restorative or endodontic treatment is done, as it can dramatically affect the overall success of treatment.

A patient who consents to endodontic treatment must be informed if the tooth has a questionable prognosis. The clinician must be able to interpret the subjective and objective findings that suggest a vertical root fracture or split tooth,

be able to make a prediction as to the eventual potential of healing, and convey this information to the patient. A more detailed discussion on vertical root fractures is described in Chapter 22.

PERFORATIONS

Root perforations are clinical complications that may lead to treatment failure. When root perforation occurs,



Fig. 1.34 Physical trauma from sports-related injuries or seizure-induced trauma, if directed accordingly, may cause a vertical root fracture in a tooth. This fracture occurred in a 7-year-old child secondary to trauma from a grand mal seizure.

communications between the root canal system and either periradicular tissues or the oral cavity may reduce the prognosis of treatment. Root perforations may result from extensive carious lesions, resorption, or operator error occurring during root canal instrumentation or post preparation.

The treatment prognosis of root perforations depends on the size, location, time of diagnosis and treatment, degree of periodontal damage, as well as the sealing ability and biocompatibility of the repair material.⁴⁵ It has been recognized that treatment success depends mainly on immediate sealing of the perforation and appropriate infection control. Among the materials that are commonly used to seal root perforations are mineral trioxide aggregate (MTA), super ethoxybenzoic acid (EBA) cement, intermediate restorative material (IRM), glass ionomer cements, and composites. The topic of perforations is further discussed in Chapter 20.

Clinical Classification of Pulpal and Periapical Diseases

Many attempts have been made over the years to develop classifications of pulpal and periapical disease. However, studies have shown that making a correlation between clinical signs and symptoms and the histopathology of a given clinical condition is challenging.^{122,123} Therefore *clinical* classifications have been developed in order to formulate treatment plan options. In the most general terms,

the objective and subjective findings are used to classify the suspected pathosis, with the assigned designations merely representing the presence of healthy or diseased tissue.

The terminology and classifications that follow are based on those suggested by the American Association of Endodontists in 2016.⁶

PULPAL DISEASE

Normal Pulp

This is a clinical diagnostic category in which the pulp is symptom-free and normally responsive to pulp testing.⁶ Teeth with normal pulp do not usually exhibit any spontaneous symptoms. The symptoms produced from pulp tests are mild, do not cause the patient distress, and result in a transient sensation that resolves in seconds. Radiographically, there may be varying degrees of pulpal calcification but no evidence of resorption, caries, or mechanical pulp exposure. No endodontic treatment is indicated for these teeth.


Pulpitis

This is a clinical and histologic term denoting inflammation of the dental pulp, clinically described as reversible or irreversible and histologically described as acute, chronic, or hyperplastic.⁶

Reversible Pulpitis

This is a clinical diagnosis based on subjective and objective findings indicating that the inflammation should resolve and the pulp return to normal.⁶ When the pulp within the tooth is irritated so that the stimulation is uncomfortable to the patient but reverses quickly after irritation, it is classified as *reversible pulpitis*. Causative factors include caries, exposed dentin, recent dental treatment, and defective restorations. Conservative removal of the irritant will resolve the symptoms. Confusion can occur when there is exposed dentin, without evidence of pulp pathosis, which can sometimes respond with sharp, quickly reversible pain when subjected to thermal, evaporative, tactile, mechanical, osmotic, or chemical stimuli. This is known as *dentin* (or *dentinal*) *sensitivity* (or *hypersensitivity*). Exposed dentin in the cervical area of the tooth accounts for most of the cases diagnosed as dentin sensitivity.¹⁰³

As described in Chapter 14, fluid movement within dentinal tubules can stimulate the odontoblasts and associated fast-conducting A-delta nerve fibers in the pulp, which in turn produce sharp, quickly reversible dental pain (Fig. 1.35). The more open these tubules are (e.g., from a newly exposed preparation, dentin decalcification, periodontal scaling, tooth-bleaching materials, or coronal tooth fractures), the more the tubule fluid will move and, subsequently, the more the tooth will display dentin sensitivity when stimulated. When making a diagnosis, it is important to discriminate this dentin sensitivity sensation from that of reversible pulpitis, which would be secondary to caries, trauma, or new or defective restorations. Detailed questioning about recent dental treatment and a thorough clinical and radiographic examination will help to separate dentin sensitivity from other pulpal pathosis, as the treatment modalities for each are completely different.¹⁸



Dentin tubule fluid movement

Fig. 1.35 Dentinal tubules are filled with fluid that, when stimulated, will cause sensation. Temperature changes, air, and osmotic changes can provoke the odontoblastic process to induce the stimulation of underlying A-delta fibers.

Irreversible Pulpitis

As the disease state of the pulp progresses, the inflammatory condition of the pulp can change to *irreversible pulpitis*. At this stage, treatment to remove the diseased pulp will be necessary. This condition can be divided into the subcategories of *symptomatic* and *asymptomatic* irreversible pulpitis.

Symptomatic Irreversible Pulpitis. This is a clinical diagnosis based on subjective and objective findings indicating that the vital inflamed pulp is incapable of healing.⁶ Teeth that are classified as having *symptomatic irreversible pulpitis* exhibit intermittent or spontaneous pain. Rapid exposure to dramatic temperature changes (especially to cold stimuli) will elicit heightened and prolonged episodes of pain even after the thermal stimulus has been removed. The pain in these cases may be sharp or dull, localized, diffuse, or referred. Typically, there are minimal or no changes in the radiographic appearance of the periradicular bone. With advanced irreversible pulpitis, a thickening of the periodontal ligament may become apparent on the radiograph, and there may be some evidence of pulpal irritation by virtue of extensive pulp chamber or root canal space calcification. Deep restorations, caries, pulp exposure, or any other direct or indirect insult to the pulp, recently or historically, may be present. It may be seen radiographically or clinically or may be suggested from a complete dental history. Patients who present with symptomatic anterior teeth for which there are no obvious etiologic factors also should be questioned regarding past general anesthesia or endotracheal intubation procedures.^{3,127,138} In addition, patients should be questioned about a history of orthodontic treatment. Typically, when symptomatic irreversible pulpitis remains untreated, the pulp will eventually become necrotic.^{109,139}

Asymptomatic Irreversible Pulpitis. This is a clinical diagnosis based on subjective and objective findings indicating that the vital inflamed pulp is incapable of healing.⁶

However, the patient does not complain of any symptoms. On occasion, deep caries will not produce any symptoms, even though clinically or radiographically the caries may extend well into the pulp. Left untreated, the tooth may become symptomatic or the pulp will become necrotic. In cases of *asymptomatic irreversible pulpitis*, endodontic treatment should be performed as soon as possible so that symptomatic irreversible pulpitis or necrosis does not develop and cause the patient severe pain and distress.

Pulp Necrosis

This is a clinical diagnostic category indicating death of the dental pulp. The pulp is usually nonresponsive to pulp testing.⁶ When *pulpal necrosis* (or *nonvital pulp*) occurs, the pulpal blood supply is nonexistent and the pulpal nerves are nonfunctional. It is the only clinical classification that directly attempts to describe the histologic status of the pulp (or lack thereof). This condition is subsequent to symptomatic or asymptomatic irreversible pulpitis. After the pulp becomes completely necrotic, the tooth will typically become asymptomatic until such time when there is an extension of the disease process into the periradicular tissues. With pulp necrosis, the tooth will usually not respond to electric pulp tests or to cold stimulation. However, if heat is applied for an extended period of time, the tooth may respond to this stimulus. This response could possibly be related to remnants of fluid or gases in the pulp canal space expanding and extending into the periapical tissues.

Pulpal necrosis may be partial or complete and it may not involve all of the canals in a multiradical tooth. For this reason, the tooth may present with confusing symptoms. Pulp testing over one root may give no response, whereas over another root it may give a positive response. The tooth may also exhibit symptoms of symptomatic irreversible pulpitis. Pulp necrosis, in the absence of restorations, caries, or luxation injuries, is likely caused by a longitudinal fracture extending from the occlusal surface and into the pulp.¹⁹

After the pulp becomes necrotic, microbial growth can be sustained within the canal. When this infection (or its microbial byproducts) extends into the periodontal ligament space, the tooth may become symptomatic to percussion or exhibit spontaneous pain. Radiographic changes may occur, ranging from a thickening of the periodontal ligament space to the appearance of a periapical radiolucent lesion. The tooth may become hypersensitive to heat, even to the warmth of the oral cavity, and is often relieved by applications of cold. As previously discussed, this may be helpful in attempting to localize a necrotic tooth (i.e., by the application of cold one tooth at a time) when the pain is referred or not well localized.

Previously Treated

This is a clinical diagnostic category indicating that the tooth has been endodontically treated and the canals are obturated with various filling materials other than intracanal medicaments.⁶ In this situation, the tooth may or may not present with signs or symptoms but will require additional nonsurgical or surgical endodontic procedures to retain the tooth. In most such situations, there will no longer be any vital or necrotic pulp tissue present to respond to pulp testing procedures.

Previously Initiated Therapy

This is a clinical diagnostic category indicating that the tooth has been previously treated by partial endodontic therapy (e.g., pulpotomy, pulpectomy).⁶ In most instances, the partial endodontic therapy was performed as an emergency procedure for symptomatic or asymptomatic irreversible pulpitis cases. In other situations, these procedures may have been performed as part of vital pulp therapy procedures, traumatic tooth injuries, apexification, or apexogenesis therapy. At the time these cases present for root canal therapy it would not be possible to make an accurate pulpal diagnosis because all, or part, of the pulp tissue has already been removed.

APICAL (PERIAPICAL) DISEASE

Normal Apical Tissues

This classification is the standard against which all of the other apical disease processes are compared. In this category the patient is asymptomatic and the tooth responds normally to percussion and palpation testing. The radiograph reveals an intact lamina dura and periodontal ligament space around all the root apices.

Periodontitis

This classification refers to an inflammation of the periodontium.⁶ When located in the periapical tissues it is referred to as apical periodontitis. Apical periodontitis can be subclassified to symptomatic apical periodontitis and asymptomatic apical periodontitis.

Symptomatic Apical Periodontitis

This condition is defined as an inflammation, usually of the apical periodontium, producing clinical symptoms including a painful response to biting or percussion or palpation. It might or might not be associated with an apical radiolucent area.⁶ This tooth may or may not respond to pulp sensibility tests, and the radiograph or image of the tooth will typically exhibit at least a widened periodontal ligament space and may or may not show an apical radiolucency associated with one or all of the roots.

Asymptomatic Apical Periodontitis

This condition is defined as inflammation and destruction of apical periodontium that is of pulpal origin; it appears as an apical radiolucent area, and does not produce clinical symptoms.⁶ This tooth does not usually respond to pulp sensibility tests, and the radiograph or image of the tooth will exhibit an apical radiolucency. The tooth is generally not sensitive to biting pressure but may “feel different” to the patient on percussion. Manifestation of persistent apical periodontitis may vary among patients.^{8,9}

Acute Apical Abscess

This condition is defined as an inflammatory reaction to pulpal infection and necrosis characterized by *rapid onset*, spontaneous pain, tenderness of the tooth to pressure, pus formation, and swelling of associated tissues.⁶ A tooth with an *acute apical abscess* will be acutely painful to biting pressure, percussion, and palpation. This tooth will not respond to any pulp sensibility tests and will exhibit varying degrees of mobility. The radiograph or image can exhibit anything

from a widened periodontal ligament space to an apical radiolucency. Swelling will be present intraorally and the facial tissues adjacent to the tooth will almost always present with some degree of swelling. The patient will frequently be febrile, and the cervical and submandibular lymph nodes may exhibit tenderness to palpation.

Chronic Apical Abscess

This condition is defined as an inflammatory reaction to pulpal infection and necrosis characterized by *gradual onset*, little or no discomfort, and the intermittent discharge of pus through an associated sinus tract.⁶ The sinus tract can appear intraorally or extraorally. In general, a tooth with a *chronic apical abscess* will not present with clinical symptoms. The tooth will not respond to pulp sensibility tests, and the radiograph or image will exhibit an apical radiolucency. Usually the tooth is not sensitive to biting pressure but can “feel different” to the patient on percussion. This entity is distinguished from asymptomatic apical periodontitis because it will exhibit intermittent drainage through an associated sinus tract.

Referred Pain

The perception of pain in one part of the body that is distant from the actual source of the pain is known as *referred pain*. Whereas pain of nonodontogenic origin can refer pain to the teeth, teeth may also refer pain to other teeth as well as to other anatomic areas of the head and neck. This may create a diagnostic challenge in that the patient may insist that the pain is from a certain tooth or even from an ear when, in fact, it is originating from a distant tooth with pulpal pathosis. Using electronic pulp testers, investigators found that patients could localize *which* tooth was being stimulated only 37.2% of the time and could narrow the location to three teeth only 79.5% of the time, illustrating that patients may have a difficult time discriminating the exact location of pulpal pain.⁴⁴

Referred pain from a tooth is usually provoked by an intense stimulation of pulpal C fibers, the slow conducting nerves that, when stimulated, cause an intense, slow, dull pain. Anterior teeth seldom refer pain to other teeth or to opposite arches, whereas posterior teeth may refer pain to the opposite arch or to the periauricular area but seldom to the anterior teeth.¹⁴ Mandibular posterior teeth tend to transmit referred pain to the periauricular area more often than maxillary posterior teeth. One study showed that when second molars were stimulated with an electric pulp tester, patients could discriminate accurately which arch the sensation was coming from only 85% of the time, compared with an accuracy level of 95% with first molars and 100% with anterior teeth.¹³⁶ The investigators also pointed out that when patients first feel the sensation of pain, they are more likely to accurately discriminate the origin of the pain. With higher levels of discomfort, patients have less ability to accurately determine the source of the pain. Therefore, in cases of diffuse or referred pain, the history of where the patient first felt the pain may be significant.

Because referred pain can complicate a dental diagnosis, the clinician must be sure to make an accurate diagnosis to protect the patient from unnecessary dental or medical

treatment. After all the testing procedures are complete and if it is determined that the pain is not of odontogenic origin, then the patient should be referred to an orofacial pain clinic for further testing. For further information on pain of nonodontogenic origin, see Chapter 4.

Summary

Endodontics is a multifaceted specialty, with much emphasis on how cases are clinically treated. Clinicians have increased their ability to more accurately perform endodontic procedures by way of increased visualization using the operating microscope, precise apical foramen detection using electronic apex locators, enhanced imaging techniques using digital radiography, three-dimensional imaging, and more. Practices have incorporated more refined canal cleaning and shaping techniques by using ultrasonics and rotary-driven nickel titanium files facilitated with computer-assisted electronic handpieces. Many other advancements also have been introduced with the objective of achieving an optimal result during endodontic treatment. However, these advancements are useless if an incorrect diagnosis is made. Before the clinician ever considers performing any endodontic treatment, the following questions must be answered:

- Is the existing problem of dental origin?
- Are the pulpal tissues within the tooth pathologically involved?
- Why is the pulpal pathosis present?
- What is the prognosis?
- What is the appropriate form of treatment?

Testing, questioning, and reasoning are combined to achieve an accurate diagnosis and to ultimately form an appropriate treatment plan. The art and science of making this diagnosis are the first steps that must be taken before initiating any endodontic treatment.

References

1. Abella F, Patel S, Duran-Sindreu F, et al: Evaluating the periapical status of teeth with irreversible pulpitis by using cone-beam computed tomography scanning and periapical radiographs, *J Endod* 38:1588, 2012.
2. Abuabara A, Zielak JC, Schramm CA, et al: Dental infection simulating skin lesion, *An Bras Dermatol* 87:619, 2012.
3. Adolphs N, Kessler B, von Heymann C, et al: Dentoalveolar injury related to general anaesthesia: a 14 years review and a statement from the surgical point of view based on a retrospective analysis of the documentation of a university hospital, *Dent Traumatol* 27:10, 2011.
4. Al-Hezaimi K, Naghshbandi J, Simon JH, et al: Successful treatment of a radicular groove by intentional replantation and emdogain therapy: four years follow-up, *Oral Surg Oral Med Oral Pathol Oral Radiol Endodon* 107:e82, 2009.
5. Alomari FA, Al-Hababeh R, Alsakarna BK: Responses of pulp sensibility tests during orthodontic treatment and retention, *Int Endod J* 44:635, 2011.
6. American Association of Endodontists: *Glossary of endodontic terms*, ed 9, Chicago, 2016, American Association of Endodontists.
7. Anderson RW, Pantera EA: Influence of a barrier technique on electric pulp testing, *J Endod* 14:179, 1988.
8. Andreasen JD, Andreasen FM, Andersson L, editors: *Textbook and color atlas of traumatic injuries to the teeth*, ed 5, Philadelphia, 2019, Wiley Blackwell.
9. Andreasen JO, Ahrensburg SS, Tsilingaridis G: Root fractures: the influence of type of healing and location of fracture on tooth survival rates: an analysis of 492 cases, *Dent Traumatol* 28:404, 2012.
10. Arakawa S, Cobb CM, Rapley JW, et al: Treatment of root fracture by CO₂ and Nd:YAG lasers: an in vitro study, *J Endod* 22:662, 1996.
11. Augsburger RA, Peters DD: In vitro effects of ice, skin refrigerant, and CO₂ snow on intrapulpal temperature, *J Endod* 7:110, 1981.
12. Baumgartner JC, Picket AB, Muller JT: Microscopic examination of oral sinus tracts and their associated periapical lesions, *J Endod* 10:146, 1984.
13. Beltes C, Zachou E: Endodontic management in a patient with vitamin D-resistant rickets, *J Endod* 38:255, 2012.
14. Bender IB: Pulpal pain diagnosis: a review, *J Endod* 26:175, 2000.
15. Bender IB, Landau MA, Fonseca S, et al: The optimum placement-site of the electrode in electric pulp testing of the 12 anterior teeth, *J Am Dent Assoc* 118:305, 1989.
16. Bender IB, Seltzer S: Roentgenographic and direct observation of experimental lesions in bone. Part I, *J Am Dent Assoc* 62:152, 1961.
17. Bender IB, Seltzer S: Roentgenographic and direct observation of experimental lesions in bone. Part II, *J Am Dent Assoc* 62:708, 1961.
18. Berman LH: Dentinal sensation and hypersensitivity: a review of mechanisms and treatment alternatives, *J Periodontol* 56:216, 1984.
19. Berman LH, Kuttler S: Fracture necrosis: diagnosis, prognosis assessment, and treatment recommendations, *J Endod* 36:442, 2010.
20. Bierma MK, McClanahan S, Baisden MK, et al: Comparison of heat-testing methodology, *J Endod* 38:1106, 2012.
21. Bornstein MM, Lauber R, Sendi P, et al: Comparison of periapical radiography and limited cone-beam computed tomography in mandibular molars for analysis of anatomical landmarks before apical surgery, *J Endod* 37:151, 2011.
22. Cameron CE: The cracked tooth syndrome: additional findings, *J Am Dent Assoc* 93:971, 1981.
23. Chen E, Abbott PV: Evaluation of accuracy, reliability, and repeatability of five dental pulp tests, *J Endod* 37:1619, 2011.
24. Chiego DJ, Cox CF, Avery JK: H-3 HRP analysis of the nerve supply to primate teeth, *Dent Res* 59:736, 1980.
25. Cleveland JL, Gooch BF, Shearer BG, et al: Risk and prevention of hepatitis C virus infection, *J Am Dent Assoc* 130:641, 1999.
26. Costa FF, Gaia BF, Umetsubo OS, et al: Detection of horizontal root fracture with small-volume cone-beam computed tomography in the presence and absence of intracanal metallic post, *J Endod* 37:1456, 2011.
27. Costa FF, Gaia BF, Umetsubo OS, et al: Use of large-volume cone-beam computed tomography in identification and localization of horizontal root fracture in the presence and absence of intracanal metallic post, *J Endod* 38:856, 2012.
28. Costa FWG, Rodrigues RR, Batista ACB: Multiple radiopaque mandibular lesions in a patient with Apert syndrome, *J Endod* 38:1639, 2012.
29. Cotton TP, Geisler TM, Holden DT, et al: Endodontic applications of cone-beam volumetric tomography, *J Endod* 33:1121, 2007.
30. Dankner E, Harari D, Rotstein I: Dens evaginatus of anterior teeth: literature review and radiographic survey of 15,000 teeth, *Oral Surg Oral Med Oral Pathol* 81:472, 1996.
31. Dastmalchi N, Jafarzadeh H, Moradi S: Comparison of the efficacy of a custom-made pulse oximeter probe with digital electric pulp tester, cold spray, and rubber cup for assessing pulp vitality, *J Endod* 38:1182, 2012.
32. Davido N, Rigolet A, Kerner S, et al: Case of Ewing's sarcoma misdiagnosed as a periapical lesion of maxillary incisor, *J Endod* 37:259, 2011.
33. Deepak BS, Subash TS, Narmatha VJ, et al: Imaging techniques in endodontics: an overview, *J Clin Imaging Sci* 2:13, 2012.
34. DeRossi SS, Glick M: Dental considerations for the patient with renal disease receiving hemodialysis, *J Am Dent Assoc* 127:211, 1996.
35. DeRossi SS, Glick M: Lupus erythematosus: considerations for dentistry, *J Am Dent Assoc* 129:330, 1998.
36. Dirks SJ, Paunovich ED, Terezhalmay GT, et al: The patient with Parkinson's disease, *Quint Int* 34:379, 2003.
37. Durack C, Patel S: Cone beam computed tomography in endodontics, *Braz Dent J* 23:179, 2012.
38. Edlund M, Nair MK, Nair UP: Detection of vertical root fractures by using cone-beam computed tomography: a clinical study, *J Endod* 37:768, 2011.

39. Eikener S, Vandre R: Comparison of digital dental x-ray systems with self-developing film and manual processing for endodontic file length determination. *J Endod* 26:65, 2000.
40. Evans D, Reid J, Strang R, et al: A comparison of laser Doppler flowmetry with other methods of assessing the vitality of traumatized anterior teeth. *Endod Dent Traumatol* 15:284, 1999.
41. Farman AG, Levato CM, Scarfe WC: A primer on cone beam CT. *Inside Dentistry* 1:90, 2007.
42. Filippatos CG, Tsatsoulis IN, Floratos S, et al: A variability of electric pulp response threshold in premolars: a clinical study. *J Endod* 38:144, 2012.
43. Fouad AF: Diabetes mellitus as a modulating factor of endodontic infections. *J Dent Educ* 67:459, 2003.
44. Friend LA, Glenwright HD: An experimental investigation into the localization of pain from the dental pulp. *Oral Surg Oral Med Oral Pathol* 25:765, 1968.
45. Fuss Z, Trope M: Root perforations: classification and treatment choices based on prognostic factors. *Endod Dent Traumatol* 12:255, 1996.
46. Fuss Z, Trowbridge H, Bender IB, et al: Assessment of reliability of electrical and thermal pulp testing agents. *J Endod* 12:301, 1986.
47. Ganz SD: Cone beam computed tomography-assisted treatment planning concepts. *Dent Clin North Am* 55:515, 2011.
48. Gillcrist JA: Hepatitis viruses A, B, C, D, E and G: implications for dental personnel. *J Am Dent Assoc* 130:509, 1999.
49. Goldman M, Pearson A, Darzenta N: Endodontic success: who is reading the radiograph? *Oral Surg Oral Med Oral Pathol* 33:432, 1972.
50. Goldman M, Pearson A, Darzenta N: Reliability of radiographic interpretations. *Oral Surg Oral Med Oral Pathol* 38:287, 1974.
51. Goodchild JH, Glick M: A different approach to medical risk assessment. *Endod Topics* 4:1, 2003.
52. Goon WW, Jacobsen PL: Prodromal odontalgia and multiple devitalized teeth caused by a herpes zoster infection of the trigeminal nerve: report of case. *J Am Dent Assoc* 116:500, 1988.
53. Gopikrishna V, Tinagupta K, Kandaswamy D: Comparison of electrical, thermal and pulse oximetry methods for assessing pulp vitality in recently traumatized teeth. *J Endod* 33:531, 2007.
54. Gopikrishna V, Tinagupta K, Kandaswamy D: Evaluation of efficacy of a new custom-made pulse oximeter dental probe in comparison with electrical and thermal tests for assessing pulp vitality. *J Endod* 33:411, 2007.
55. Harrison JW, Larson WJ: The epithelized oral sinus tract. *Oral Surg Oral Med Oral Pathol* 42:511, 1976.
56. Heling I, Rotstein I: A persistent oronasal sinus tract of endodontic origin. *J Endod* 15:132, 1989.
57. Herman WW, Konzelman JL, Prisant LM: New national guidelines on hypertension. *J Am Dent Assoc* 135:576, 2004.
58. Idiyatullin D, Corum C, Moeller S, et al: Dental magnetic resonance imaging: making the invisible visible. *J Endod* 37:745, 2011.
59. Imai K, Shimada Y, Sadr A, et al: Noninvasive cross-sectional visualization of enamel cracks by optical coherence tomography *in vitro*. *J Endod* 38:1269, 2012.
60. Ingolfsson AER, Tronstad L, Riva CE: Reliability of laser Doppler flowmetry in testing vitality of human teeth. *Endod Dent Traumatol* 10:185, 1994.
61. Ingram TA, Peters DD: Evaluation of the effects of carbon dioxide used as a pulp test. Part 2: in vivo effect on canine enamel and pulpal tissues. *J Endod* 9:296, 1983.
62. Jafarzadeh H, Abbott PV: Review of pulp sensibility tests. Part I: general information and thermal tests. *Int Endod J* 43:738, 2010.
63. Jafarzadeh H, Abbott PV: Review of pulp sensibility tests. Part II: electric pulp tests and test cavities. *Int Endod J* 43:945, 2010.
64. Johnson BR, Remeikis NA, Van Cura JE: Diagnosis and treatment of cutaneous facial sinus tracts of dental origin. *J Am Dent Assoc* 130:832, 1999.
65. Jones DM: Effect of the type carrier used on the results of dichlorodifluoromethane application to teeth. *J Endod* 25:692, 1999.
66. Jones VR, Rivera EM, Walton RE: Comparison of carbon dioxide versus refrigerant spray to determine pulpal responsiveness. *J Endod* 28:531, 2002.
67. Kaffe I, Gratt BM: Variations in the radiographic interpretation of the periapical dental region. *J Endod* 14:330, 1988.
68. Kahan RS, Gulabivala K, Snook M, et al: Evaluation of a pulse oximeter and customized probe for pulp vitality testing. *J Endod* 22:105, 1996.
69. Karayilmaz H, Kirzioglu Z: Comparison of the reliability of laser Doppler flowmetry, pulse oximetry and electric pulp tester in assessing the pulp vitality of human teeth. *J Oral Rehabil* 38:340, 2011.
70. Kataoka SH, Setzer FC, Gondim-Junior E, et al: Pulp vitality in patients with intraoral and oropharyngeal malignant tumors undergoing radiation therapy assessed by pulse oximetry. *J Endod* 37:1197, 2011.
71. Katz J, Chaushu G, Rotstein I: Stafne's bone cavity in the anterior mandible: a possible diagnosis challenge. *J Endod* 27:304, 2001.
72. Kawai K, Masaka N: Vertical root fracture treated by bonding fragments and rotational replantation. *Dent Traumatol* 18:42, 2002.
73. Khocht A, Janal M, Harasty L, et al: Comparison of direct digital and conventional intraoral radiographs in detecting alveolar bone loss. *J Am Dent Assoc* 134:1468, 2003.
74. Koivisto T, Bowles WR, Rohrer M: Frequency and distribution of radiolucent jaw lesions: a retrospective analysis of 9,723 cases. *J Endod* 38:729, 2012.
75. Kusgoz A, Yildirim T, Kayipmaz S, et al: Nonsurgical endodontic treatment of type III dens invaginatus in maxillary canine: an 18-month follow up. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 107:e103, 2009.
76. Lalla RV, D'Ambrosio JA: Dental management considerations for the patient with diabetes mellitus. *J Am Dent Assoc* 132:1425, 2001.
77. Laskin DM: Anatomic considerations in diagnosis and treatment of odontogenic infections. *J Am Dent Assoc* 69:308, 1964.
78. Liang YH, Li G, Wessellink PR, et al: Endodontic outcome predictors identified with periapical radiographs and cone-beam computed tomography scans. *J Endod* 37:326, 2011.
79. Lin J, Chandler NP: Electric pulp testing: a review. *Int Endod J* 41:365, 2008.
80. Little JW, Miller CS, Rhodus NL, editors: *Dental management of the medically compromised patient*, ed 9. St. Louis, 2018, Elsevier.
81. Lofthag-Hansen S, Huuonen S, Gröndahl K, et al: Limited cone-beam CT and intraoral radiography for the diagnosis of periapical pathology. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 103:114, 2007.
82. Marder MZ: The standard of care for oral diagnosis as it relates to oral cancer. *Comp Contin Educ Dent* 19:569, 1998.
83. Mattson JS, Cerutis DR: Diabetes mellitus: a review of the literature and dental implications. *Comp Cont Educ Dent* 22:757, 2001.
84. Mesaros S, Trope M, Maixner W, et al: Comparison of two laser Doppler systems on the measurement of blood flow of premolar teeth under different pulpal conditions. *Int Endod J* 30:167, 1997.
85. Metska ME, Aartman IHA, Wessellink PR, et al: Detection of vertical root fractures *in vivo* in endodontically treated teeth by cone-beam computed tomography scans. *J Endod* 38:1344, 2012.
86. Michaelson RE, Seidberg BH, Guttuso J: An *in vivo* evaluation of interface media used with the electric pulp tester. *J Am Dent Assoc* 91:118, 1975.
87. Miles DA, VanDis ML: Advances in dental imaging. *Dent Clin North Am* 37:531, 1993.
88. Miller CS, Little JW, Falace DA: Supplemental corticosteroids for dental patients with adrenal insufficiency: reconsideration of the problem. *J Am Dent Assoc* 132:1570, 2001.
89. Morsani JM, Aminoshariae A, Han YW: Genetic predisposition to persistent apical periodontitis. *J Endod* 37:455, 2011.
90. Mozzo P, Proccacci A, Tacconi, et al: A new volumetric CT machine for dental imaging based on the cone-beam technique: preliminary results. *Eur Radiol* 8:1558, 1998.
91. Nair M, Pettigrew J, Mancuso A: Intracranial aneurysm as an incidental finding. *Dentomaxillofac Radiol* 36:107, 2007.
92. Nakata K, Naitob M, Izumi M, et al: Effectiveness of dental computed tomography in diagnostic imaging of periradicular lesion of each root of a multirooted tooth: a case report. *J Endod* 32:583, 2007.
93. Nalliah RP, Allareddy V, Elangovan S, et al: Hospital emergency department visits attributed to pulpal and periapical disease in the United States in 2006. *J Endod* 37:6, 2011.
94. Özer SY: Detection of vertical root fractures by using cone beam computed tomography with variable voxel sizes in an *in vitro* model. *J Endod* 37:75, 2011.
95. Pantera EA, Anderson RW, Pantera CT: Use of dental instruments for bridging during electric pulp testing. *J Endod* 18:37, 1992.
96. Pantera EA, Anderson RW, Pantera CT: Reliability of electric pulp testing after pulpal testing with dichlorodifluoromethane. *J Endod* 19:312, 1993.

97. Paurazas SM, Geist JR, Pink FE: Comparison of diagnostic accuracy of digital imaging using CCD and CMOS-APS sensors with E-speed film in the detection of periapical bony lesions, *Oral Surg Oral Med Oral Pathol Oral Radiology Endodon* 44:249, 2000.
98. Peters DD, Baumgartner JC, Lorton L: Adult pulpal diagnosis. 1. Evaluation of the positive and negative responses to cold and electric pulp tests, *J Endod* 20:506, 1994.
99. Petersson K, Soderstrom C, Kiani-Anaraki M, et al: Evaluation of the ability of thermal and electric tests to register pulp vitality, *Endod Dent Traumatol* 15:127, 1999.
100. Pinto A, Glick M: Management of patients with thyroid disease: oral health considerations, *J Am Dent Assoc* 133:849, 2002.
101. Pitts DL, Natkin E: Diagnosis and treatment of vertical root fractures, *J Endod* 9:338, 1983.
102. Poeschl PW, Crepaz V, Russmueller G, et al: Endodontic pathogens causing deep neck space infections: clinical impact of different sampling techniques and antibiotic susceptibility, *J Endod* 37:1201, 2011.
103. Rees JS, Addy M: A cross-sectional study of dentine hypersensitivity, *J Clin Periodontol* 29:997, 2002.
104. Rickoff B, Trowbridge H, Baker J, et al: Effects of thermal vitality tests on human dental pulp, *J Endod* 14:482, 1988.
105. Riley CK, Terezhalmay GT: The patient with hypertension, *Quint Int* 32:671, 2001.
106. Robb RA, Sinak LJ, Hoffman EA, et al: Dynamic volume imaging of moving organs, *J Med Syst* 6:539, 1982.
107. Rodrigues CD, Villar-Neto MJC, Sobral APV, et al: Lymphangioma mimicking apical periodontitis, *J Endod* 37:91, 2011.
108. Rosen H: Cracked tooth syndrome, *J Prosthet Dent* 47:36, 1982.
109. Rotstein I, Engel G: Conservative management of a combined endodontic-orthodontic lesion, *Endod Dent Traumatol* 7:266, 1991.
110. Rotstein I, Moshonov J, Cohenca N: Endodontic therapy for a fused mandibular molar, *Endod Dent Traumatol* 13:149, 1997.
111. Rotstein I: Interaction between endodontics and periodontics, *Periodontol* 2000 74:11, 2017.
112. Rotstein I, Zabalegui B, Al-Hezaimi K: Endodontic-periodontal interrelationship. In Rotstein I, Ingle JI, editors: *Ingle's endodontics*, ed 7, Raleigh, North Carolina, 2019, PMPH USA, p 1111.
113. Rotstein I, Stabholz A, Heling I, et al: Clinical considerations in the treatment of dens invaginatus, *Endod Dent Traumatol* 3:249, 1987.
114. Roykens H, Van Maele G, DeMoor R, et al: Reliability of laser Doppler flowmetry in a 2-probe assessment of pulpal blood flow, *Oral Surg Oral Med Oral Pathol Oral Radiol Endodon* 87:742, 1999.
115. Rud J, Omnell KA: Root fractures due to corrosion: diagnostic aspects, *Scand J Dent Res* 78:397, 1970.
116. Rutsatz C, Baumhardt SG, Feldens CA, et al: Response of pulp sensibility test is strongly influenced by periodontal attachment loss and gingival recession, *J Endod* 38:580, 2012.
117. Sasano T, Nakajima I, Shohi N, et al: Possible application of transmitted laser light for the assessment of human pulpal vitality, *Endod Dent Traumatol* 13:88, 1997.
118. Schnettler JM, Wallace JA: Pulse oximetry as a diagnostic tool of pulp vitality, *J Endod* 17:488, 1991.
119. Schwartz RS: Mineral trioxide aggregate: a new material for endodontics, *J Am Dent Assoc* 130:967, 1999.
120. Selden HS: Repair of incomplete vertical root fractures in endodontically treated teeth: in vivo trials, *J Endod* 22:426, 1996.
121. Seltzer S, Bender IB, Nazimov H: Differential diagnosis of pulp conditions, *Oral Surg Oral Med Oral Pathol* 19:383, 1965.
122. Seltzer S, Bender IB, Zientz M: The dynamics of pulp inflammation: correlations between diagnostic data and actual histologic findings in the pulp. Part I, *Oral Surg Oral Med Oral Pathol* 16:846, 1963.
123. Seltzer S, Bender IB, Zientz M: The dynamics of pulp inflammation: correlations between diagnostic data and actual histologic findings in the pulp. Part II, *Oral Surg Oral Med Oral Pathol* 16:969, 1963.
124. Seo DG, Yi YA, Shin AJ, et al: Analysis of factors associated with cracked teeth, *J Endod* 38:288, 2012.
125. Setzer FC, Kataoka SH, Natrielli F, et al: Clinical diagnosis of pulp inflammation based on pulp oxygenation rates measured by pulse oximetry, *J Endod* 38:880, 2012.
126. Simon JHS, Dogan H, Ceresa LM, et al: The radicular groove: it's potential clinical significance, *J Endod* 26:295, 2000.
127. Simon JHS, Lies J: Silent trauma, *Endod Dent Traumatol* 15:145, 1999.
128. Shemesh H, Cristescu RC, Wesslink PR, et al: The use of cone-beam computed tomography and digital periapical radiographs to diagnose root perforations, *J Endod* 37:513, 2011.
129. Steinbacher DM, Glick M: The dental patient with asthma: an update and oral health considerations, *J Am Dent Assoc* 132:1229, 2001.
130. Stroblitt H, Gojer G, Norer B, et al: Assessing revascularization of avulsed permanent maxillary incisors by laser Doppler flowmetry, *J Am Dent Assoc* 134:1597, 2003.
131. Suebnukarn S, Rhenmora P, Haddawy P: The use of cone-beam computed tomography and virtual reality simulation for pre-surgical practice in endodontic microsurgery, *Int Endod J* 45:627, 2012.
132. Sugaya T, Kawanami M, Noguchi H, et al: Periodontal healing after bonding treatment of vertical root fracture, *Dent Traumatol* 17:174, 2001.
133. Tatlidil R, Gözübüyük MM: Mucinous adenocarcinoma of lung presenting as oral metastases: a case report and literature review, *J Endod* 37:110, 2011.
134. Tewary S, Luzzo J, Hartwell G: Endodontic radiography: who is reading the digital radiograph, *J Endod* 37:919, 2011.
135. Treister N, Glick M: Rheumatoid arthritis: a review and suggested dental care considerations, *J Am Dent Assoc* 130:689, 1999.
136. Van Hassel HJ, Harrington GW: Localization of pulpal sensation, *Oral Surg Oral Med Oral Pathol* 28:753, 1969.
137. Velvart P, Hecker H, Tillinger G: Detection of the apical lesion and the mandibular canal in conventional radiography and computed tomography, *Oral Surg Oral Med Oral Pathol Oral Radiol Endodon* 92:682, 2001.
138. Vogel J, Stubinger S, Kaufmann M: Dental injuries resulting from tracheal intubation: a retrospective study, *Dent Traumatol* 25:73, 2009.
139. Von Böhl M, Ren Y, Fudalej PS, et al: Pulpal reactions to orthodontic force application in humans: a systematic review, *J Endod* 38:1463, 2012.
140. Wallace JA, Schnettler JM: Pulse oximetry as a diagnostic tool of pulpal vitality, *J Endod* 17:488, 1993.
141. Weisleder R, Yamauchi S, Caplan DJ, et al: The validity of pulp testing: a clinical study, *J Am Dent Assoc* 140:1013, 2009.
142. Zou X, Liu D, Yue L, et al: The ability of cone-beam computerized tomography to detect vertical root fractures in endodontically treated and nonendodontically treated teeth: a report of 3 cases, *Oral Surg Oral Med Oral Pathol Oral Radiol Endodon* 111:797, 2011.