

CSE 6740 A/ISyE 6740: Computational Data Analysis: Introductory lecture

Nisha Chandramoorthy

August 24, 2023

Last time

- ▶ Goal in this class: understand the foundations (“why”s and “how”s) of ML

Last time

- ▶ Goal in this class: understand the foundations (“why”s and “how”s) of ML
- ▶ Supervised, unsupervised, self-supervised, semi-supervised overview

Last time

- ▶ Goal in this class: understand the foundations (“why”s and “how”s) of ML
- ▶ Supervised, unsupervised, self-supervised, semi-supervised overview
- ▶ Empirical risk minimization, finite hypothesis classes

Last time

- ▶ Goal in this class: understand the foundations (“why”s and “how”s) of ML
- ▶ Supervised, unsupervised, self-supervised, semi-supervised overview
- ▶ Empirical risk minimization, finite hypothesis classes
- ▶ Overfitting, inductive bias, Intro to PAC learning

Supervised learning framework

- ▶ Distribution \mathcal{D} over the joint distribution of random variables $Z = (X, Y)$, where X is an input and Y is a label/output.

Supervised learning framework

- ▶ Distribution \mathcal{D} over the joint distribution of random variables $Z = (X, Y)$, where X is an input and Y is a label/output.
- ▶ Labeled training data, $S = \{z_i = (x_i, y_i)\}, 1 \leq i \leq m$. Generally iid from \mathcal{D}^m .

Supervised learning framework

- ▶ Distribution \mathcal{D} over the joint distribution of random variables $Z = (X, Y)$, where X is an input and Y is a label/output.
- ▶ Labeled training data, $S = \{z_i = (x_i, y_i)\}, 1 \leq i \leq m$. Generally iid from \mathcal{D}^m .
- ▶ Learner's output: predicted function or hypothesis, a transformation h from X to Y .

Supervised learning framework

- ▶ Distribution \mathcal{D} over the joint distribution of random variables $Z = (X, Y)$, where X is an input and Y is a label/output.
- ▶ Labeled training data, $S = \{z_i = (x_i, y_i)\}, 1 \leq i \leq m$. Generally iid from \mathcal{D}^m .
- ▶ Learner's output: predicted function or hypothesis, a transformation h from X to Y .
- ▶ Loss function, measure of risk: $\ell(z, h) \in \mathbb{R}$. e.g., $\ell(z, h) = \mathbb{1}_{h(x) \neq y}$. (classification)

Supervised learning framework

- ▶ Distribution \mathcal{D} over the joint distribution of random variables $Z = (X, Y)$, where X is an input and Y is a label/output.
- ▶ Labeled training data, $S = \{z_i = (x_i, y_i)\}, 1 \leq i \leq m$. Generally iid from \mathcal{D}^m .
- ▶ Learner's output: predicted function or hypothesis, a transformation h from X to Y .
- ▶ Loss function, measure of risk: $\ell(z, h) \in \mathbb{R}$. e.g., $\ell(z, h) = \mathbb{1}_{h(x) \neq y}$. (classification)
- ▶ Generalization error or risk:

$$R(h) = E_{z \sim \mathcal{D}} \ell(z, h)$$

Empirical Risk Minimization

- ▶ Take classifier h , $R(h) = \mathcal{D}(\{z : h(x) \neq y\})$.

Empirical Risk Minimization

- ▶ Take classifier h , $R(h) = \mathcal{D}(\{z : h(x) \neq y\})$.
- ▶ When you have finite amount of data, empirical risk or training loss

$$\hat{R}_S(h) = \frac{1}{m} \sum_{z \in S} \ell(z) = \frac{|\{z \in S : h(x) \neq y\}|}{m}.$$

Empirical Risk Minimization

- ▶ Take classifier h , $R(h) = \mathcal{D}(\{z : h(x) \neq y\})$.
- ▶ When you have finite amount of data, empirical risk or training loss

$$\hat{R}_S(h) = \frac{1}{m} \sum_{z \in S} \ell(z) = \frac{|\{z \in S : h(x) \neq y\}|}{m}.$$

- ▶ ERM: find h that minimizes $\hat{R}_S(h)$.

Empirical Risk Minimization

- ▶ Take classifier h , $R(h) = \mathcal{D}(\{z : h(x) \neq y\})$.
- ▶ When you have finite amount of data, empirical risk or training loss

$$\hat{R}_S(h) = \frac{1}{m} \sum_{z \in S} \ell(z) = \frac{|\{z \in S : h(x) \neq y\}|}{m}.$$

- ▶ ERM: find h that minimizes $\hat{R}_S(h)$.
- ▶ Theory of supervised learning suggests that ERM leads to small generalization error with high probability

Empirical Risk Minimization

- ▶ Take classifier h , $R(h) = \mathcal{D}(\{z : h(x) \neq y\})$.
- ▶ When you have finite amount of data, empirical risk or training loss

$$\hat{R}_S(h) = \frac{1}{m} \sum_{z \in S} \ell(z) = \frac{|\{z \in S : h(x) \neq y\}|}{m}.$$

- ▶ ERM: find h that minimizes $\hat{R}_S(h)$.
- ▶ Theory of supervised learning suggests that ERM leads to small generalization error with high probability
- ▶ Now we will prove this result formally.

Empirical Risk Minimization

- ▶ Take classifier h , $R(h) = \mathcal{D}(\{z : h(x) \neq y\})$.
- ▶ When you have finite amount of data, empirical risk or training loss

$$\hat{R}_S(h) = \frac{1}{m} \sum_{z \in S} \ell(z) = \frac{|\{z \in S : h(x) \neq y\}|}{m}.$$

- ▶ ERM: find h that minimizes $\hat{R}_S(h)$.
- ▶ Theory of supervised learning suggests that ERM leads to small generalization error with high probability
- ▶ Now we will prove this result formally.
- ▶ After that: Linear models.

Linear models

$$\mathcal{H} = \{h(\cdot, w, b) : h(x, w, b) = w \cdot \Phi(x) + b, w \in \mathbb{R}^d, b \in \mathbb{R}\} \quad (1)$$

Linear models

$$\mathcal{H} = \{h(\cdot, w, b) : h(x, w, b) = w \cdot \Phi(x) + b, w \in \mathbb{R}^d, b \in \mathbb{R}\} \quad (1)$$

► $\Phi : \mathbb{X} \rightarrow \mathbb{R}^d$ is a set of *features*.

Linear models

$$\mathcal{H} = \{h(\cdot, w, b) : h(x, w, b) = w \cdot \Phi(x) + b, w \in \mathbb{R}^d, b \in \mathbb{R}\} \quad (1)$$

- ▶ $\Phi : \mathbb{X} \rightarrow \mathbb{R}^d$ is a set of *features*.
- ▶ Linear regression seeks ERM solution for square loss

Linear models

$$\mathcal{H} = \{h(\cdot, w, b) : h(x, w, b) = w \cdot \Phi(x) + b, w \in \mathbb{R}^d, b \in \mathbb{R}\} \quad (1)$$

- ▶ $\Phi : \mathbb{X} \rightarrow \mathbb{R}^d$ is a set of *features*.
- ▶ Linear regression seeks ERM solution for square loss
- ▶

$$\operatorname{argmin}_{w,b} \frac{1}{m} (w^\top \Phi(x_i) + b - y_i)^2$$

Linear models

$$\mathcal{H} = \{h(\cdot, w, b) : h(x, w, b) = w \cdot \Phi(x) + b, w \in \mathbb{R}^d, b \in \mathbb{R}\} \quad (1)$$

- ▶ $\Phi : \mathbb{X} \rightarrow \mathbb{R}^d$ is a set of *features*.
- ▶ Linear regression seeks ERM solution for square loss



$$\operatorname{argmin}_{w,b} \frac{1}{m} (w^\top \Phi(x_i) + b - y_i)^2$$

- ▶ Equivalently, where X is $m \times (d + 1)$ matrix with rows $X_i = (\Phi(x_i)^\top, 1)$, $W = [w_1, \dots, w_d, 1]^\top$, $Y = [y_1, \dots, y_m]^\top$,

Linear models

$$\mathcal{H} = \{h(\cdot, w, b) : h(x, w, b) = w \cdot \Phi(x) + b, w \in \mathbb{R}^d, b \in \mathbb{R}\} \quad (1)$$

- ▶ $\Phi : \mathbb{X} \rightarrow \mathbb{R}^d$ is a set of *features*.
- ▶ Linear regression seeks ERM solution for square loss



$$\operatorname{argmin}_{w,b} \frac{1}{m} (w^\top \Phi(x_i) + b - y_i)^2$$

- ▶ Equivalently, where X is $m \times (d + 1)$ matrix with rows $X_i = (\Phi(x_i)^\top, 1)$, $W = [w_1, \dots, w_d, 1]^\top$, $Y = [y_1, \dots, y_m]^\top$,

$$\operatorname{argmin}_W \frac{1}{m} \|XW - Y\|^2$$

- ▶ Features may be defined by kernels

Least squares solutions: the optimization way

- ▶ Convex, differentiable function of W – composition of convex, differentiable functions

Least squares solutions: the optimization way

- ▶ Convex, differentiable function of W – composition of convex, differentiable functions
- ▶ Global minimum is the extremal point where derivative vanishes

Least squares solutions: the optimization way

- ▶ Convex, differentiable function of W – composition of convex, differentiable functions
- ▶ Global minimum is the extremal point where derivative vanishes
- ▶ $\nabla \frac{1}{m} \|XW - Y\|^2 = \frac{2}{m} X^T (XW - Y)$

Least squares solutions: the optimization way

- ▶ Convex, differentiable function of W – composition of convex, differentiable functions
- ▶ Global minimum is the extremal point where derivative vanishes
- ▶ $\nabla \frac{1}{m} \|XW - Y\|^2 = \frac{2}{m} X^T (XW - Y)$
- ▶ $X^T XW = X^T Y$. Can also get this by differentiating before writing in matrix form

Least squares solutions: the optimization way

- ▶ Convex, differentiable function of W – composition of convex, differentiable functions
- ▶ Global minimum is the extremal point where derivative vanishes
- ▶ $\nabla \frac{1}{m} \|XW - Y\|^2 = \frac{2}{m} X^T (XW - Y)$
- ▶ $X^T XW = X^T Y$. Can also get this by differentiating before writing in matrix form
- ▶ When is $X^T X = \sum_{i=1}^m \Phi(x_i) \Phi(x_i)^T$ invertible? When the training features span \mathbb{R}^d .

Least squares solutions: the linear algebraic way

- ▶ Case 1: $d + 1 = m$, X is invertible. $W = X^{-1} Y$

Least squares solutions: the linear algebraic way

- ▶ Case 1: $d + 1 = m$, X is invertible. $W = X^{-1} Y$
- ▶ Case 2: $d + 1 > m$, underdetermined/overparameterized. If X has full row rank, then, min norm solution

$$W = X^{\top} (X X^{\top})^{-1} Y$$

Least squares solutions: the linear algebraic way

- ▶ Case 1: $d + 1 = m$, X is invertible. $W = X^{-1} Y$
- ▶ Case 2: $d + 1 > m$, underdetermined/overparameterized. If X has full row rank, then, min norm solution

$$W = X^{\top} (X X^{\top})^{-1} Y$$

- ▶ Case 3: $m > d + 1$, overdetermined. If X has full col rank, then, many solutions. Min norm solution

$$W = (X^{\top} X)^{-1} X^{\top} Y$$

Least squares solutions: the linear algebraic way

- ▶ Case 1: $d + 1 = m$, X is invertible. $W = X^{-1} Y$
- ▶ Case 2: $d + 1 > m$, underdetermined/overparameterized. If X has full row rank, then, min norm solution

$$W = X^{\top} (X X^{\top})^{-1} Y$$

- ▶ Case 3: $m > d + 1$, overdetermined. If X has full col rank, then, many solutions. Min norm solution

$$W = (X^{\top} X)^{-1} X^{\top} Y$$

- ▶ Can solve normal equations above directly, or use iterative methods for linear systems. Cost $\mathcal{O}(d^3)$

Gauss Markov theorem

- ▶ Take noisy $y_i = x_i^\top W + \epsilon_i$, with $E_i = 0$ and $\text{Var}(\epsilon_i) = \sigma^2$; x_i is non-random.

Gauss Markov theorem

- ▶ Take noisy $y_i = x_i^\top W + \epsilon_i$, with $E_i = 0$ and $\text{Var}(\epsilon_i) = \sigma^2$; x_i is non-random.
- ▶ Statement: the OLS estimator is the best linear unbiased estimator (blue). It has the lowest variance.

Gauss Markov theorem

- ▶ Take noisy $y_i = x_i^\top W + \epsilon_i$, with $E_i = 0$ and $\text{Var}(\epsilon_i) = \sigma^2$; x_i is non-random.
- ▶ Statement: the OLS estimator is the best linear unbiased estimator (blue). It has the lowest variance.
- ▶ Proof: consider another linear estimator $W' = CY$, for some $(d+1) \times n$ matrix $C = (X^\top X)^{-1} X^\top + D$.

Gauss Markov theorem

- ▶ Take noisy $y_i = x_i^\top W + \epsilon_i$, with $E_i = 0$ and $\text{Var}(\epsilon_i) = \sigma^2$; x_i is non-random.
- ▶ Statement: the OLS estimator is the best linear unbiased estimator (blue). It has the lowest variance.
- ▶ Proof: consider another linear estimator $W' = CY$, for some $(d+1) \times n$ matrix $C = (X^\top X)^{-1} X^\top + D$.
- ▶ For W' to be unbiased, show $DX = 0$. Then show, $\text{Var}(W') = \text{Var}(W) + \sigma^2 DD^\top$.

Gauss Markov theorem

- ▶ Take noisy $y_i = x_i^\top W + \epsilon_i$, with $E_i = 0$ and $\text{Var}(\epsilon_i) = \sigma^2$; x_i is non-random.
- ▶ Statement: the OLS estimator is the best linear unbiased estimator (blue). It has the lowest variance.
- ▶ Proof: consider another linear estimator $W' = CY$, for some $(d+1) \times n$ matrix $C = (X^\top X)^{-1}X^\top + D$.
- ▶ For W' to be unbiased, show $DX = 0$. Then show, $\text{Var}(W') = \text{Var}(W) + \sigma^2 DD^\top$.
- ▶ Since DD^\top is positive semi-definite, qed.

Ridge regression

- ▶ Motivation: unbiased estimation does not mean least mean-squared error

Ridge regression

- ▶ Motivation: unbiased estimation does not mean least mean-squared error
- ▶ Let true $h_W \in \mathcal{H}$. Mean-squared error of statistical estimator of W , \hat{W} , is its variance + bias-squared

$$E[(\hat{W}_i - W_i)^2] = \text{Var}(\hat{W}_i) + (E[\hat{W}_i] - W_i)^2.$$



$$\operatorname{argmin}_W \frac{1}{m} \|XW - Y\|^2 + \lambda \|W\|^2.$$

Ridge regression

- ▶ Motivation: unbiased estimation does not mean least mean-squared error
- ▶ Let true $h_W \in \mathcal{H}$. Mean-squared error of statistical estimator of W , \hat{W} , is its variance + bias-squared

$$E[(\hat{W}_i - W_i)^2] = \text{Var}(\hat{W}_i) + (E[\hat{W}_i] - W_i)^2.$$



$$\operatorname{argmin}_W \frac{1}{m} \|XW - Y\|^2 + \lambda \|W\|^2.$$

- ▶ penalizes l^2 norm of W . Still convex problem.

Ridge regression

- ▶ Motivation: unbiased estimation does not mean least mean-squared error
- ▶ Let true $h_W \in \mathcal{H}$. Mean-squared error of statistical estimator of W , \hat{W} , is its variance + bias-squared

$$E[(\hat{W}_i - W_i)^2] = \text{Var}(\hat{W}_i) + (E[\hat{W}_i] - W_i)^2.$$



$$\operatorname{argmin}_W \frac{1}{m} \|XW - Y\|^2 + \lambda \|W\|^2.$$

- ▶ penalizes l^2 norm of W . Still convex problem.
- ▶ to derive OLS, also can take derivative and set it to zero. Similarly here.

Ridge regression

- ▶ Motivation: unbiased estimation does not mean least mean-squared error
- ▶ Let true $h_W \in \mathcal{H}$. Mean-squared error of statistical estimator of W , \hat{W} , is its variance + bias-squared

$$E[(\hat{W}_i - W_i)^2] = \text{Var}(\hat{W}_i) + (E[\hat{W}_i] - W_i)^2.$$



$$\operatorname{argmin}_W \frac{1}{m} \|XW - Y\|^2 + \lambda \|W\|^2.$$

- ▶ penalizes l^2 norm of W . Still convex problem.
- ▶ to derive OLS, also can take derivative and set it to zero. Similarly here.
- ▶ Equivalent formulation: $\min_W \sum_{i=1}^m (w^\top \Phi(x_i) - y_i)^2$ subject to $\|w\|^2 \leq \Lambda^2$

Ridge regression solution and interpretation

- ▶ Revisit convex optimization. Derive solution using KKT conditions.

Ridge regression solution and interpretation

- ▶ Revisit convex optimization. Derive solution using KKT conditions.
- ▶ Now simply use convexity and differentiability to obtain global minimum:

$$W = (X^T X + \lambda I)^{-1} X^T Y.$$

Ridge regression solution and interpretation

- ▶ Revisit convex optimization. Derive solution using KKT conditions.
- ▶ Now simply use convexity and differentiability to obtain global minimum:

$$W = (X^T X + \lambda I)^{-1} X^T Y.$$

- ▶ LASSO: with l^1 regularization.

Ridge regression solution and interpretation

- ▶ Revisit convex optimization. Derive solution using KKT conditions.
- ▶ Now simply use convexity and differentiability to obtain global minimum:

$$W = (X^T X + \lambda I)^{-1} X^T Y.$$

- ▶ LASSO: with l^1 regularization.
- ▶ Generalization bounds for bounded regression problems.

Ridge regression solution and interpretation

- ▶ Revisit convex optimization. Derive solution using KKT conditions.
- ▶ Now simply use convexity and differentiability to obtain global minimum:

$$W = (X^T X + \lambda I)^{-1} X^T Y.$$

- ▶ LASSO: with l^1 regularization.
- ▶ Generalization bounds for bounded regression problems.
- ▶ Shrinkage by l^2 regularization.