

Lecture 22: Spectral clustering, EM algorithm

Nisha Chandramoorthy

November 14, 2023

Lloyd's algorithm

- ▶ Randomly choose k centers $\mu_1, \dots, \mu_k \in \mathbb{R}^d$.

Lloyd's algorithm

- ▶ Randomly choose k centers $\mu_1, \dots, \mu_k \in \mathbb{R}^d$.
- ▶ Given centers $\mu_1, \dots, \mu_k \in \mathbb{R}^d$, assign each point x_i to the closest center. That is,

$$C_j = \{x_i : j \in \operatorname{argmin}_l \|x_i - \mu_l\|\}.$$

Lloyd's algorithm

- ▶ Randomly choose k centers $\mu_1, \dots, \mu_k \in \mathbb{R}^d$.
- ▶ Given centers $\mu_1, \dots, \mu_k \in \mathbb{R}^d$, assign each point x_i to the closest center. That is,

$$C_j = \{x_i : j \in \operatorname{argmin}_l \|x_i - \mu_l\|\}.$$

- ▶ Given clusters C_1, \dots, C_k , update centers $\mu_1, \dots, \mu_k \in \mathbb{R}^d$ as

$$\mu_j = \frac{1}{|C_j|} \sum_{x_i \in C_j} x_i.$$

k-means algorithm (Lloyd's algorithm)

- ▶ Lloyd's algorithm is an approximate method to solve the ERM problem:

$$\min_{C_1, \dots, C_k} \sum_{j=1}^k \sum_{x_i \in C_j} \|x_i - \mu(C_j)\|^2.$$

k-means algorithm (Lloyd's algorithm)

- ▶ Lloyd's algorithm is an approximate method to solve the ERM problem:

$$\min_{C_1, \dots, C_k} \sum_{j=1}^k \sum_{x_i \in C_j} \|x_i - \mu(C_j)\|^2.$$

- ▶ here, $\mu(C_j) = \frac{1}{|C_j|} \sum_{x_i \in C_j} x_i = \operatorname{argmin}_{\mu \in \mathbb{R}^d} \sum_{x_i \in C_j} \|x_i - \mu\|^2$ is the mean of the points in cluster C_j .

k-means algorithm (Lloyd's algorithm)

- ▶ Lloyd's algorithm is an approximate method to solve the ERM problem:

$$\min_{C_1, \dots, C_k} \sum_{j=1}^k \sum_{x_i \in C_j} \|x_i - \mu(C_j)\|^2.$$

- ▶ here, $\mu(C_j) = \frac{1}{|C_j|} \sum_{x_i \in C_j} x_i = \operatorname{argmin}_{\mu \in \mathbb{R}^d} \sum_{x_i \in C_j} \|x_i - \mu\|^2$ is the mean of the points in cluster C_j .
- ▶ Lloyd's algorithm is a heuristic. It is not guaranteed to converge to the global optimum or even a local minimum.



Lloyd's algorithm properties

- ▶ k-means algorithm is sensitive to initialization of the centers.

Lloyd's algorithm properties



- ▶ k-means algorithm is sensitive to initialization of the centers.
- ▶ Complexity: $O(mdk)$ per iteration, where m is the number of points, d is the dimension, and k is the number of clusters.

k-means failure modes



Source: [sklearn's toy examples](#)

k-means failure modes contd



Source: [sklearn's toy examples](#)

Spectral clustering

- ▶ Given distance d or similarity matrix, $W \in \mathbb{R}^{m \times m}$, partition the points into k clusters.

Spectral clustering

- ▶ Given distance d or similarity matrix, $W \in \mathbb{R}^{m \times m}$, partition the points into k clusters.
- ▶ W is symmetric and non-negative.

Spectral clustering

- ▶ Given distance d or similarity matrix, $W \in \mathbb{R}^{m \times m}$, partition the points into k clusters.
- ▶ W is symmetric and non-negative.
- ▶ W is a weighted adjacency matrix of a graph.

Spectral clustering

- ▶ Given distance d or similarity matrix, $W \in \mathbb{R}^{m \times m}$, partition the points into k clusters.
- ▶ W is symmetric and non-negative.
- ▶ W is a weighted adjacency matrix of a graph.
- ▶ ERM problem: $\min_{C_1, \dots, C_k} \sum_{j=1}^k \sum_{x_i \in C_j} \sum_{x_l \notin C_j} w_{il}$. Graph min-cut problem.

RatioCut problem: spectral clustering solution

► RatioCut problem: $\min_{C_1, \dots, C_k} \sum_{j=1}^k \frac{\sum_{x_i \in C_j} \sum_{x_l \notin C_j} w_{il}}{|C_j|}.$

RatioCut problem: spectral clustering solution

- ▶ RatioCut problem: $\min_{C_1, \dots, C_k} \sum_{j=1}^k \frac{\sum_{x_i \in C_j} \sum_{x_l \notin C_j} w_{il}}{|C_j|}$.
- ▶ Normalization by $|C_j|$ penalizes small clusters.

RatioCut problem: spectral clustering solution

- ▶ RatioCut problem: $\min_{C_1, \dots, C_k} \sum_{j=1}^k \frac{\sum_{x_i \in C_j} \sum_{x_l \notin C_j} w_{il}}{|C_j|}$.
- ▶ Normalization by $|C_j|$ penalizes small clusters.

RatioCut objective

- ▶ Lemma 22.3 (Ben-David and Shalev Shwartz) RatioCut objective = $\text{Tr}(H^{\top} L H)$

RatioCut objective

- ▶ Lemma 22.3 (Ben-David and Shalev Shwartz) RatioCut objective = $\text{Tr}(H^\top L H)$
- ▶ $L = D - W$ is the graph Laplacian, where D is the diagonal matrix with $D_{ii} = \sum_{j=1}^m w_{ij}$.

RatioCut objective

- ▶ Lemma 22.3 (Ben-David and Shalev Shwartz) RatioCut objective = $\text{Tr}(H^\top L H)$
- ▶ $L = D - W$ is the graph Laplacian, where D is the diagonal matrix with $D_{ii} = \sum_{j=1}^m w_{ij}$.
- ▶ $H \in \mathbb{R}^{m \times k}$ is the indicator matrix of the clusters.
 $H_{ij} = 1/\sqrt{|C_j|}$ if $x_i \in C_j$ and 0 otherwise.

RatioCut objective

- ▶ Lemma 22.3 (Ben-David and Shalev Schwartz) RatioCut objective = $\text{Tr}(H^\top L H)$
- ▶ $L = D - W$ is the graph Laplacian, where D is the diagonal matrix with $D_{ii} = \sum_{j=1}^m w_{ij}$.
- ▶ $H \in \mathbb{R}^{m \times k}$ is the indicator matrix of the clusters.
 $H_{ij} = 1/\sqrt{|C_j|}$ if $x_i \in C_j$ and 0 otherwise.
- ▶ h_i (i th column of H) is nonzero at row j if x_j is in cluster i .

RatioCut objective

- ▶ Lemma 22.3 (Ben-David and Shalev Schwartz) RatioCut objective = $\text{Tr}(H^\top L H)$
- ▶ $L = D - W$ is the graph Laplacian, where D is the diagonal matrix with $D_{ii} = \sum_{j=1}^m w_{ij}$.
- ▶ $H \in \mathbb{R}^{m \times k}$ is the indicator matrix of the clusters.
 $H_{ij} = 1/\sqrt{|C_j|}$ if $x_i \in C_j$ and 0 otherwise.
- ▶ h_i (i th column of H) is nonzero at row j if x_j is in cluster i .
- ▶ H has orthonormal columns.

Recall: graphical representation of X

- ▶ Choose weighting, such as, $w_{ij} = \exp(-\|x_i - x_j\|^2 / 2\sigma^2)$. As $\sigma \rightarrow 0$, $w_{ij} \rightarrow \mathbb{1}_{i=j}$. The $m \times m$ matrix W is the adjacency matrix of a graph.

Recall: graphical representation of X

- ▶ Choose weighting, such as, $w_{ij} = \exp(-\|x_i - x_j\|^2 / 2\sigma^2)$. As $\sigma \rightarrow 0$, $w_{ij} \rightarrow \mathbb{1}_{i=j}$. The $m \times m$ matrix W is the adjacency matrix of a graph.
- ▶ Let D be the diagonal matrix with $D_{ii} = \sum_{j=1}^m w_{ij}$.

Recall: graphical representation of X

- ▶ Choose weighting, such as, $w_{ij} = \exp(-\|x_i - x_j\|^2 / 2\sigma^2)$. As $\sigma \rightarrow 0$, $w_{ij} \rightarrow \mathbb{1}_{i=j}$. The $m \times m$ matrix W is the adjacency matrix of a graph.
- ▶ Let D be the diagonal matrix with $D_{ii} = \sum_{j=1}^m w_{ij}$.
- ▶ Graph laplacian: $L = D - W$.

Recall: graphical representation of X

- ▶ Choose weighting, such as, $w_{ij} = \exp(-\|x_i - x_j\|^2 / 2\sigma^2)$. As $\sigma \rightarrow 0$, $w_{ij} \rightarrow \mathbb{1}_{i=j}$. The $m \times m$ matrix W is the adjacency matrix of a graph.
- ▶ Let D be the diagonal matrix with $D_{ii} = \sum_{j=1}^m w_{ij}$.
- ▶ Graph laplacian: $L = D - W$.
- ▶ Detects local structure / clusters in data.

Lemma proof: RatioCut objective and graph laplacian connection

► RatioCut objective(C_1, \dots, C_k)

$$:= \sum_{j=1}^k \frac{\sum_{x_i \in C_j} \sum_{x_l \notin C_j} w_{il}}{|C_j|}.$$

Lemma proof: RatioCut objective and graph laplacian connection

- ▶ RatioCut objective(C_1, \dots, C_k)

$$:= \sum_{j=1}^k \frac{\sum_{x_i \in C_j} \sum_{x_l \notin C_j} w_{il}}{|C_j|}.$$

- ▶ Need to show equal to $\text{Tr}(H^T L H)$.

Laplacian eigenmaps

- ▶ Want to solve: $\min_{y_1, \dots, y_m} \sum_{i=1}^m \sum_{j=1}^m w_{ij} \|y_i - y_j\|^2$.

Laplacian eigenmaps

- ▶ Want to solve: $\min_{y_1, \dots, y_m} \sum_{i=1}^m \sum_{j=1}^m w_{ij} \|y_i - y_j\|^2$.
- ▶ optimal embeddings: $y_i = E(x_i) = U[i, -n :]$ where U is the matrix of eigenvectors of L .

Laplacian eigenmaps

- ▶ Want to solve: $\min_{y_1, \dots, y_m} \sum_{i=1}^m \sum_{j=1}^m w_{ij} \|y_i - y_j\|^2$.
- ▶ optimal embeddings: $y_i = E(x_i) = U[i, -n :]$ where U is the matrix of eigenvectors of L .
- ▶ For any vector v , $v^\top L v = (1/2) \sum_{i,j=1}^m w_{ij} (v_i - v_j)^2$.

Laplacian eigenmaps

- ▶ Want to solve: $\min_{y_1, \dots, y_m} \sum_{i=1}^m \sum_{j=1}^m w_{ij} \|y_i - y_j\|^2$.
- ▶ optimal embeddings: $y_i = E(x_i) = U[i, -n :]$ where U is the matrix of eigenvectors of L .
- ▶ For any vector v , $v^\top L v = (1/2) \sum_{i,j=1}^m w_{ij} (v_i - v_j)^2$.
- ▶ L is positive semi-definite.

Bottom n eigenvectors

- ▶ Rayleigh quotient optimality

Bottom n eigenvectors

- ▶ Rayleigh quotient optimality
- ▶ Another interpretation: top n eigenvectors of L^\dagger . L_{ij}^\dagger represents expected time for random walk $i \rightarrow j \rightarrow i$.

Bottom n eigenvectors

- ▶ Rayleigh quotient optimality
- ▶ Another interpretation: top n eigenvectors of L^\dagger . L_{ij}^\dagger represents expected time for random walk $i \rightarrow j \rightarrow i$.
- ▶ Kernel PCA with $K = L^\dagger$ is equivalent to Laplacian eigenmaps.

Combining dimension reduction and k-means

- ▶ Spectral clustering algorithm uses Laplacian eigenmaps on m -dimensional data.

Combining dimension reduction and k-means

- ▶ Spectral clustering algorithm uses Laplacian eigenmaps on m -dimensional data.
- ▶ Uses $v_i, i = 1, 2, \dots, k$ eigenvectors of L corresponding to the k smallest eigenvalues.

Combining dimension reduction and k-means

- ▶ Spectral clustering algorithm uses Laplacian eigenmaps on m -dimensional data.
- ▶ Uses $v_i, i = 1, 2, \dots, k$ eigenvectors of L corresponding to the k smallest eigenvalues.
- ▶ Perform k-means on rows of v_i . to obtain clusters

Gaussian mixtures

- ▶ Suppose we want to cluster data that is generated from a mixture of Gaussians.

Gaussian mixtures

- ▶ Suppose we want to cluster data that is generated from a mixture of Gaussians.
- ▶ $x_i \sim \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$.

Gaussian mixtures

- ▶ Suppose we want to cluster data that is generated from a mixture of Gaussians.
- ▶ $x_i \sim \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$.
- ▶ Frequentist view: there is a true (unknown) parameter $\theta = (\pi_1, \dots, \pi_k, \mu_1, \dots, \mu_k, \Sigma_1, \dots, \Sigma_k)$ that generated the data.

- ▶ Clustering objective: maximize log likelihood of the data.

- ▶ Clustering objective: maximize log likelihood of the data.
- ▶ $\ell(x, \theta) = \log p_{\theta}(x) = \log \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$.

- ▶ Clustering objective: maximize log likelihood of the data.
- ▶ $\ell(x, \theta) = \log p_{\theta}(x) = \log \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$.
- ▶ $\hat{R}_S(\theta) = \sum_{i=1}^m \log \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$.

- ▶ Clustering objective: maximize log likelihood of the data.
- ▶ $\ell(x, \theta) = \log p_{\theta}(x) = \log \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$.
- ▶ $\hat{R}_S(\theta) = \sum_{i=1}^m \log \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$.
- ▶ More generally, $\hat{R}_S(\theta) = \sum_{i=1}^m \log \sum_{j=1}^k q_{\theta}(z_j) p_{\theta}(x_i | z_j)$.

- ▶ Clustering objective: maximize log likelihood of the data.
- ▶ $\ell(x, \theta) = \log p_{\theta}(x) = \log \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$.
- ▶ $\hat{R}_S(\theta) = \sum_{i=1}^m \log \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$.
- ▶ More generally, $\hat{R}_S(\theta) = \sum_{i=1}^m \log \sum_{j=1}^k q_{\theta}(z_j) p_{\theta}(x_i | z_j)$.
- ▶ The joint distribution $p_{\theta}(x, z) = q_{\theta}(z) p_{\theta}(x | z)$ is parametrized by θ .

- ▶ Clustering objective: maximize log likelihood of the data.
- ▶ $\ell(x, \theta) = \log p_{\theta}(x) = \log \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$.
- ▶ $\hat{R}_S(\theta) = \sum_{i=1}^m \log \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$.
- ▶ More generally, $\hat{R}_S(\theta) = \sum_{i=1}^m \log \sum_{j=1}^k q_{\theta}(z_j) p_{\theta}(x_i | z_j)$.
- ▶ The joint distribution $p_{\theta}(x, z) = q_{\theta}(z) p_{\theta}(x | z)$ is parametrized by θ .
- ▶ Z is a latent variable, e.g., Z is the cluster assignment of X .

Maximizing log likelihood

- ▶ Distribution q of the latent variable is unknown.

Maximizing log likelihood

- ▶ Distribution q of the latent variable is unknown.
- ▶ Thus, we want to solve:

$$\max_{\theta} \max_q \sum_{i=1}^m \log \sum_{j=1}^k q_{\theta}(z_j) p_{\theta}(x_i | z_j). \quad (1)$$

- ▶ Lemma: For fixed θ , optimal $q_{\theta} \equiv p_{\theta}(X|Z)$ is the posterior distribution of Z given X .

Proof: derivation of ELBO

- ▶ Fix some x and θ .

Proof: derivation of ELBO

- ▶ Fix some x and θ .
- ▶ $\ell(x, \theta) = \log p_{\theta}(x) = \log \sum_{j=1}^k p_{\theta}(x, z_j) = \log \sum_{j=1}^k q_{\theta}(z_j) \frac{p_{\theta}(x, z_j)}{q_{\theta}(z_j)}.$

Proof: derivation of ELBO

- ▶ Fix some x and θ .
- ▶ $\ell(x, \theta) = \log p_{\theta}(x) = \log \sum_{j=1}^k p_{\theta}(x, z_j) = \log \sum_{j=1}^k q_{\theta}(z_j) \frac{p_{\theta}(x, z_j)}{q_{\theta}(z_j)}$.
- ▶ Use Jensen's inequality: $E \log Z \leq \log EZ$ for any random variable Z .

Proof: derivation of ELBO

- ▶ Fix some x and θ .
- ▶ $\ell(x, \theta) = \log p_{\theta}(x) = \log \sum_{j=1}^k p_{\theta}(x, z_j) = \log \sum_{j=1}^k q_{\theta}(z_j) \frac{p_{\theta}(x, z_j)}{q_{\theta}(z_j)}.$
- ▶ Use Jensen's inequality: $E \log Z \leq \log EZ$ for any random variable Z .
- ▶ Thus, $\ell(x, \theta) \geq \sum_{j=1}^k q_{\theta}(z_j) \log \frac{p_{\theta}(x, z_j)}{q_{\theta}(z_j)}.$

Proof: derivation of ELBO

- ▶ Fix some x and θ .
- ▶ $\ell(x, \theta) = \log p_\theta(x) = \log \sum_{j=1}^k p_\theta(x, z_j) = \log \sum_{j=1}^k q_\theta(z_j) \frac{p_\theta(x, z_j)}{q_\theta(z_j)}$.
- ▶ Use Jensen's inequality: $E \log Z \leq \log EZ$ for any random variable Z .
- ▶ Thus, $\ell(x, \theta) \geq \sum_{j=1}^k q_\theta(z_j) \log \frac{p_\theta(x, z_j)}{q_\theta(z_j)}$.
- ▶ This holds for any probability distribution q_θ .

Proof: derivation of ELBO

- ▶ Fix some x and θ .
- ▶ $\ell(x, \theta) = \log p_\theta(x) = \log \sum_{j=1}^k p_\theta(x, z_j) = \log \sum_{j=1}^k q_\theta(z_j) \frac{p_\theta(x, z_j)}{q_\theta(z_j)}$.
- ▶ Use Jensen's inequality: $E \log Z \leq \log EZ$ for any random variable Z .
- ▶ Thus, $\ell(x, \theta) \geq \sum_{j=1}^k q_\theta(z_j) \log \frac{p_\theta(x, z_j)}{q_\theta(z_j)}$.
- ▶ This holds for any probability distribution q_θ .
- ▶ $\text{ELBO}(q, \theta) = \sum_{j=1}^k q(z_j) \log \frac{p_\theta(x, z_j)}{q(z_j)}$.

Proof: derivation of ELBO

- ▶ Fix some x and θ .
- ▶ $\ell(x, \theta) = \log p_{\theta}(x) = \log \sum_{j=1}^k p_{\theta}(x, z_j) = \log \sum_{j=1}^k q_{\theta}(z_j) \frac{p_{\theta}(x, z_j)}{q_{\theta}(z_j)}$.
- ▶ Use Jensen's inequality: $E \log Z \leq \log EZ$ for any random variable Z .
- ▶ Thus, $\ell(x, \theta) \geq \sum_{j=1}^k q_{\theta}(z_j) \log \frac{p_{\theta}(x, z_j)}{q_{\theta}(z_j)}$.
- ▶ This holds for any probability distribution q_{θ} .
- ▶ $\text{ELBO}(q, \theta) = \sum_{j=1}^k q(z_j) \log \frac{p_{\theta}(x, z_j)}{q(z_j)}$.
- ▶ Thus, we have shown, $\ell(x, \theta) \geq \text{ELBO}(q, \theta)$ for any q .

► $\text{ELBO}(q, \theta) = -D_{KL}(q(z) || p_{\theta}(z|x)) + \log p_{\theta}(x).$

- ▶ $\text{ELBO}(q, \theta) = -D_{KL}(q(z)||p_{\theta}(z|x)) + \log p_{\theta}(x)$.
- ▶ When $q(z) = p_{\theta}(z|x)$, $\text{ELBO}(q, \theta) = \log p_{\theta}(x)$.

- ▶ $\text{ELBO}(q, \theta) = -D_{KL}(q(z)||p_{\theta}(z|x)) + \log p_{\theta}(x)$.
- ▶ When $q(z) = p_{\theta}(z|x)$, $\text{ELBO}(q, \theta) = \log p_{\theta}(x)$.
- ▶ $\text{ELBO}(p_{\theta}(Z|X), \theta) = \sum_{j=1}^k p_{\theta}(z_j|x) \log \frac{p_{\theta}(z_j|x)p_{\theta}(x)}{p_{\theta}(z_j|x)}$.

- ▶ $\text{ELBO}(q, \theta) = -D_{KL}(q(z)||p_{\theta}(z|x)) + \log p_{\theta}(x)$.
- ▶ When $q(z) = p_{\theta}(z|x)$, $\text{ELBO}(q, \theta) = \log p_{\theta}(x)$.
- ▶ $\text{ELBO}(p_{\theta}(Z|X), \theta) = \sum_{j=1}^k p_{\theta}(z_j|x) \log \frac{p_{\theta}(z_j|x)p_{\theta}(x)}{p_{\theta}(z_j|x)}$.
- ▶ $\text{ELBO}(p_{\theta}(Z|X), \theta) = \sum_{j=1}^k p_{\theta}(z_j|x) \log p_{\theta}(x) = \log p_{\theta}(x)$.

Expectation Maximization (EM) algorithm

- ▶ Iteratively maximize $\text{ELBO}(q, \theta)$ w.r.t. q and θ .

Expectation Maximization (EM) algorithm

- ▶ Iteratively maximize $\text{ELBO}(q, \theta)$ w.r.t. q and θ .
- ▶ E step: Fix θ and maximize $\text{ELBO}(q, \theta)$ w.r.t. q . That is, set $q = p_{\theta}(Z|X)$.

Expectation Maximization (EM) algorithm

- ▶ Iteratively maximize $\text{ELBO}(q, \theta)$ w.r.t. q and θ .
- ▶ E step: Fix θ and maximize $\text{ELBO}(q, \theta)$ w.r.t. q . That is, set $q = p_{\theta}(Z|X)$.
- ▶ M step: Fix q and maximize $\text{ELBO}(q, \theta)$ w.r.t. θ .

EM algorithm

- ▶ E step: $q_{t+1}(Z|X) = p_{\theta_t}(Z|X)$.

EM algorithm

- ▶ E step: $q_{t+1}(Z|X) = p_{\theta_t}(Z|X)$.
- ▶ M step: $\theta_{t+1} = \operatorname{argmax}_{\theta} \operatorname{ELBO}(q_{t+1}, \theta)$.

EM algorithm

- ▶ E step: $q_{t+1}(Z|X) = p_{\theta_t}(Z|X)$.
- ▶ M step: $\theta_{t+1} = \operatorname{argmax}_{\theta} \operatorname{ELBO}(q_{t+1}, \theta)$.
- ▶ Repeat until convergence.

EM algorithm properties

- ▶ EM algorithm decreases $\hat{R}_S(\theta) = \sum_{i=1}^m \ell(x_i, \theta)$ at each iteration.
- ▶ $\ell(x, \theta_{t+1}) = \text{ELBO}(q_{t+2}, \theta_{t+1}) \geq \text{ELBO}(q_{t+1}, \theta_{t+1}) \geq \text{ELBO}(q_{t+1}, \theta_t) = \ell(x, \theta_t)$.

Example: mixture of Gaussians

- ▶ $x_i \sim \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$. Take $\Sigma_i = I$ for simplicity.

Example: mixture of Gaussians

- ▶ $x_i \sim \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$. Take $\Sigma_j = I$ for simplicity.
- ▶ That is, $\theta = (\pi_1, \dots, \pi_k, \mu_1, \dots, \mu_k)$.

Example: mixture of Gaussians

- ▶ $x_i \sim \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$. Take $\Sigma_i = I$ for simplicity.
- ▶ That is, $\theta = (\pi_1, \dots, \pi_k, \mu_1, \dots, \mu_k)$.
- ▶ $p_\theta(x|z_j) = \mathcal{N}(\mu_j, \Sigma_j)$; $p_\theta(z_j) = \pi_j$.

Example: mixture of Gaussians

- ▶ $x_i \sim \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$. Take $\Sigma_j = I$ for simplicity.
- ▶ That is, $\theta = (\pi_1, \dots, \pi_k, \mu_1, \dots, \mu_k)$.
- ▶ $p_\theta(x|z_j) = \mathcal{N}(\mu_j, \Sigma_j)$; $p_\theta(z_j) = \pi_j$.
- ▶ E step: $q_{t+1}(z|x) = p_{\theta_t}(z|x) = \frac{p_{\theta_t}(x|z)p_{\theta_t}(z)}{p_{\theta_t}(x)}$.

Example: mixture of Gaussians

- ▶ $x_i \sim \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$. Take $\Sigma_j = I$ for simplicity.
- ▶ That is, $\theta = (\pi_1, \dots, \pi_k, \mu_1, \dots, \mu_k)$.
- ▶ $p_\theta(x|z_j) = \mathcal{N}(\mu_j, \Sigma_j)$; $p_\theta(z_j) = \pi_j$.
- ▶ E step: $q_{t+1}(z|x) = p_{\theta_t}(z|x) = \frac{p_{\theta_t}(x|z)p_{\theta_t}(z)}{p_{\theta_t}(x)}$.
- ▶ M step: $\theta_{t+1} = \operatorname{argmax}_\theta \sum_{i=1}^m \sum_{j=1}^k q_{t+1}(z_j|x_i) \log \frac{p_\theta(x_i, z_j)}{q_{t+1}(z_j|x_i)}$.

Example: mixture of Gaussians

- ▶ $x_i \sim \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$. Take $\Sigma_j = I$ for simplicity.
- ▶ That is, $\theta = (\pi_1, \dots, \pi_k, \mu_1, \dots, \mu_k)$.
- ▶ $p_\theta(x|z_j) = \mathcal{N}(\mu_j, \Sigma_j)$; $p_\theta(z_j) = \pi_j$.
- ▶ E step: $q_{t+1}(z|x) = p_{\theta_t}(z|x) = \frac{p_{\theta_t}(x|z)p_{\theta_t}(z)}{p_{\theta_t}(x)}$.
- ▶ M step: $\theta_{t+1} = \operatorname{argmax}_\theta \sum_{i=1}^m \sum_{j=1}^k q_{t+1}(z_j|x_i) \log \frac{p_\theta(x_i, z_j)}{q_{t+1}(z_j|x_i)}$.
- ▶ $\mu_{j,t+1} = \sum_{i=1}^m q_{t+1}(z_j|x_i) x_i$.

Example: mixture of Gaussians

- ▶ $x_i \sim \sum_{j=1}^k \pi_j \mathcal{N}(\mu_j, \Sigma_j)$. Take $\Sigma_j = I$ for simplicity.
- ▶ That is, $\theta = (\pi_1, \dots, \pi_k, \mu_1, \dots, \mu_k)$.
- ▶ $p_\theta(x|z_j) = \mathcal{N}(\mu_j, \Sigma_j)$; $p_\theta(z_j) = \pi_j$.
- ▶ E step: $q_{t+1}(z|x) = p_{\theta_t}(z|x) = \frac{p_{\theta_t}(x|z)p_{\theta_t}(z)}{p_{\theta_t}(x)}$.
- ▶ M step: $\theta_{t+1} = \operatorname{argmax}_\theta \sum_{i=1}^m \sum_{j=1}^k q_{t+1}(z_j|x_i) \log \frac{p_\theta(x_i, z_j)}{q_{t+1}(z_j|x_i)}$.
- ▶ $\mu_{j,t+1} = \sum_{i=1}^m q_{t+1}(z_j|x_i) x_i$.
- ▶ $\pi_{j,t+1} = \frac{1}{m} \sum_{i=1}^m q_{t+1}(z_j|x_i) / Z_{j,t+1}$.

VAE revisited

- ▶ Hard to compute q_t in general. VAE solves this problem differently.
- ▶ $p_\theta(x|z) = \mathcal{N}(f_\theta(z), \sigma^2 I)$. (Decoder: f_θ)
- ▶ $q_\phi(z|x) = \mathcal{N}(\mu_\phi(x), \Sigma_\phi(x))$. (Encoder: μ_ϕ, Σ_ϕ)