Lecture 1: Introduction to dynamical systems theory

Nisha Chandramoorthy

January 8, 2024

Deterministic dynamical systems, i.e., systems that evolve according to a set of rules.

- Deterministic dynamical systems, i.e., systems that evolve according to a set of rules.
- Stochastic dynamical systems, i.e., systems that evolve according to a set of rules with some randomness.

- Deterministic dynamical systems, i.e., systems that evolve according to a set of rules.
- Stochastic dynamical systems, i.e., systems that evolve according to a set of rules with some randomness.
- Questions we will not ask:

- Deterministic dynamical systems, i.e., systems that evolve according to a set of rules.
- Stochastic dynamical systems, i.e., systems that evolve according to a set of rules with some randomness.
- Questions we will not ask:
- How to control a dynamical system?
- How to design/model a dynamical system?

Questions we will ask:

- Questions we will ask:
- How to analyze a dynamical system?

- Questions we will ask:
- How to analyze a dynamical system?
- ▶ How to predict the future of a dynamical system?

- Questions we will ask:
- How to analyze a dynamical system?
- How to predict the future of a dynamical system?
- How to understand the behavior of a dynamical system?

- Questions we will ask:
- How to analyze a dynamical system?
- How to predict the future of a dynamical system?
- How to understand the behavior of a dynamical system?
- How to understand the behavior of a dynamical system in the presence of noise?

Learning outcomes

- ▶ To be able to do research in dynamical systems theory.
- "Take the dynamical systems approach" to a computational math problem.

Learning outcomes

- To be able to do research in dynamical systems theory.
- "Take the dynamical systems approach" to a computational math problem.
- Understand problems at the interface of dynamical systems and machine learning.

▶ 6 homeworks (70% of grade) and 1 final project (30% of grade).

- ▶ 6 homeworks (70% of grade) and 1 final project (30% of grade).
- ► Homeworks will be a mix of theory and computation.

- 6 homeworks (70% of grade) and 1 final project (30% of grade).
- Homeworks will be a mix of theory and computation.
- Final project will be related to your research interests and involve computational dynamical systems.

- 6 homeworks (70% of grade) and 1 final project (30% of grade).
- Homeworks will be a mix of theory and computation.
- Final project will be related to your research interests and involve computational dynamical systems.
- Syllabus, homeworks, and lecture notes will be posted on Canvas and Github.

- 6 homeworks (70% of grade) and 1 final project (30% of grade).
- Homeworks will be a mix of theory and computation.
- Final project will be related to your research interests and involve computational dynamical systems.
- Syllabus, homeworks, and lecture notes will be posted on Canvas and Github.
- See the syllabus section of canvas for the timeline.

- 6 homeworks (70% of grade) and 1 final project (30% of grade).
- Homeworks will be a mix of theory and computation.
- Final project will be related to your research interests and involve computational dynamical systems.
- Syllabus, homeworks, and lecture notes will be posted on Canvas and Github.
- See the syllabus section of canvas for the timeline.
- See syllabus.pdf for details and topics.

- 6 homeworks (70% of grade) and 1 final project (30% of grade).
- Homeworks will be a mix of theory and computation.
- Final project will be related to your research interests and involve computational dynamical systems.
- Syllabus, homeworks, and lecture notes will be posted on Canvas and Github.
- See the syllabus section of canvas for the timeline.
- See syllabus.pdf for details and topics.
- Office hours: Friday 10-11 am, CODA S1323.

► Given an initial point x_0 in phase space, $φ^t(x_0)$ is the point in phase space at time t.

- Given an initial point x_0 in phase space, $\varphi^t(x_0)$ is the point in phase space at time t.
- Flow: φ^t , which satisfies group (composition) action.

- Given an initial point x_0 in phase space, $\varphi^t(x_0)$ is the point in phase space at time t.
- Flow: φ^t , which satisfies group (composition) action.

- Given an initial point x_0 in phase space, $\varphi^t(x_0)$ is the point in phase space at time t.
- Flow: φ^t , which satisfies group (composition) action.
- $ightharpoonup \phi^0(x_0) = \mathrm{Id}(x_0) = x_0.$
- But what is a phase space?

- Given an initial point x_0 in phase space, $\varphi^t(x_0)$ is the point in phase space at time t.
- Flow: φ^t , which satisfies group (composition) action.
- But what is a phase space?
- For us, it will be a compact manifold, denoted M, for deterministic systems.

- Given an initial point x_0 in phase space, $\varphi^t(x_0)$ is the point in phase space at time t.
- Flow: φ^t , which satisfies group (composition) action.
- But what is a phase space?
- For us, it will be a compact manifold, denoted M, for deterministic systems.
- M is a d-dimensional smooth manifold.

A vector field X is a map from the phase space to the tangent space of the phase space.

- A vector field X is a map from the phase space to the tangent space of the phase space.
- $ightharpoonup v(x) \in T_X M.$

- A vector field X is a map from the phase space to the tangent space of the phase space.
- \triangleright $v(x) \in T_xM$.
- ightharpoonup ODE: $\dot{x} = v(x)$.

- A vector field X is a map from the phase space to the tangent space of the phase space.
- \triangleright $v(x) \in T_xM$.
- ightharpoonup ODE: $\dot{x} = v(x)$.

- A vector field X is a map from the phase space to the tangent space of the phase space.
- \triangleright $v(x) \in T_xM$.
- ightharpoonup ODE: $\dot{x} = v(x)$.
- Vector fields can also be interpreted as linear functionals on the space of smooth functions on the manifold.

- A vector field X is a map from the phase space to the tangent space of the phase space.
- \triangleright $v(x) \in T_xM$.
- ightharpoonup ODE: $\dot{x} = v(x)$.
- $\frac{d\varphi^t}{dt}(x) = v(\varphi^t(x)).$
- Vector fields can also be interpreted as linear functionals on the space of smooth functions on the manifold.
- ▶ $X(f) = \langle v, df \rangle$, where df is the differential of $f \in C^{\infty}(M)$.

▶ Given an initial point x_0 in phase space, $F^t(x_0)$ is the point in phase space after t iterations.

- ▶ Given an initial point x_0 in phase space, $F^t(x_0)$ is the point in phase space after t iterations.
- Map: F, which satisfies group (composition) action.

- ▶ Given an initial point x_0 in phase space, $F^t(x_0)$ is the point in phase space after t iterations.
- ▶ Map: *F*, which satisfies group (composition) action.
- $\blacktriangleright \ F^0(x) = \mathrm{Id}(x) = x.$

- ▶ Given an initial point x_0 in phase space, $F^t(x_0)$ is the point in phase space after t iterations.
- ▶ Map: *F*, which satisfies group (composition) action.
- $F^0(x) = Id(x) = x.$
- ► Can be obtained from a flow by taking the time-1 map, e.g., from time-integration of the ODE $F(x) = \varphi^{1}(x)$.

➤ A tube with a flame inside – model of jet engine – modeled using the Navier-Stokes equation.

- A tube with a flame inside model of jet engine modeled using the Navier-Stokes equation.
- ▶ The Lorenz system model of convection in the atmosphere.

- A tube with a flame inside model of jet engine modeled using the Navier-Stokes equation.
- ► The Lorenz system model of convection in the atmosphere.
- Price of a stock:

$$dS/S = \mu dt + \sigma dW$$

.

- A tube with a flame inside model of jet engine modeled using the Navier-Stokes equation.
- ► The Lorenz system model of convection in the atmosphere.
- Price of a stock:

$$dS/S = \mu dt + \sigma dW$$

.

S is the price of the stock, μ is the drift, σ is the volatility, W is the Wiener process.

- A tube with a flame inside model of jet engine modeled using the Navier-Stokes equation.
- The Lorenz system model of convection in the atmosphere.
- Price of a stock:

$$dS/S = \mu dt + \sigma dW$$

- S is the price of the stock, μ is the drift, σ is the volatility, W is the Wiener process.
- ➤ *S* is a stochastic process, dealt with later in the course.
- Delay differential equations, Partial differential equations, etc. also have infinite dimensional phase spaces.

Analysis and basic topology