

2425-MA140 Engineering Calculus

Week 04, Lecture 2 Differentiation Rules

Dr Niall Madden

University of Galway

Wednesday, 08 October, 2025

Calculus	Calculus																																	
Diorthaigh	Derivatives																																	
<table><tr><th>$f(x)$</th><th>$f'(x)$</th></tr><tr><td>x^n</td><td>nx^{n-1}</td></tr><tr><td>$\ln x$</td><td>$\frac{1}{x}$</td></tr><tr><td>e^x</td><td>e^x</td></tr><tr><td>e^{ax}</td><td>ae^{ax}</td></tr><tr><td>a^x</td><td>$a^x \ln a$</td></tr><tr><td>$\cos x$</td><td>$-\sin x$</td></tr><tr><td>$\sin x$</td><td>$\cos x$</td></tr><tr><td>$\tan x$</td><td>$\sec^2 x$</td></tr><tr><td>$\cos^{-1} \frac{x}{a}$</td><td>$-\frac{1}{\sqrt{a^2 - x^2}}$</td></tr><tr><td>$\sin^{-1} \frac{x}{a}$</td><td>$\frac{1}{\sqrt{a^2 - x^2}}$</td></tr><tr><td>$\tan^{-1} \frac{x}{a}$</td><td>$\frac{a}{a^2 + x^2}$</td></tr></table>	$f(x)$	$f'(x)$	x^n	nx^{n-1}	$\ln x$	$\frac{1}{x}$	e^x	e^x	e^{ax}	ae^{ax}	a^x	$a^x \ln a$	$\cos x$	$-\sin x$	$\sin x$	$\cos x$	$\tan x$	$\sec^2 x$	$\cos^{-1} \frac{x}{a}$	$-\frac{1}{\sqrt{a^2 - x^2}}$	$\sin^{-1} \frac{x}{a}$	$\frac{1}{\sqrt{a^2 - x^2}}$	$\tan^{-1} \frac{x}{a}$	$\frac{a}{a^2 + x^2}$	<table><tr><td>Rial an toraidh</td><td>$y = uv$ $\Rightarrow \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$</td><td>Product rule</td></tr><tr><td>Rial an lin</td><td>$y = \frac{u}{v}$ $\Rightarrow \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$</td><td>Quotient rule</td></tr><tr><td>Cuingriail</td><td>$f(x) = u(v(x))$ $\Rightarrow f'(x) = \frac{du}{dv} \frac{dv}{dx}$</td><td>Chain rule</td></tr></table>	Rial an toraidh	$y = uv$ $\Rightarrow \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$	Product rule	Rial an lin	$y = \frac{u}{v}$ $\Rightarrow \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$	Quotient rule	Cuingriail	$f(x) = u(v(x))$ $\Rightarrow f'(x) = \frac{du}{dv} \frac{dv}{dx}$	Chain rule
$f(x)$	$f'(x)$																																	
x^n	nx^{n-1}																																	
$\ln x$	$\frac{1}{x}$																																	
e^x	e^x																																	
e^{ax}	ae^{ax}																																	
a^x	$a^x \ln a$																																	
$\cos x$	$-\sin x$																																	
$\sin x$	$\cos x$																																	
$\tan x$	$\sec^2 x$																																	
$\cos^{-1} \frac{x}{a}$	$-\frac{1}{\sqrt{a^2 - x^2}}$																																	
$\sin^{-1} \frac{x}{a}$	$\frac{1}{\sqrt{a^2 - x^2}}$																																	
$\tan^{-1} \frac{x}{a}$	$\frac{a}{a^2 + x^2}$																																	
Rial an toraidh	$y = uv$ $\Rightarrow \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$	Product rule																																
Rial an lin	$y = \frac{u}{v}$ $\Rightarrow \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$	Quotient rule																																
Cuingriail	$f(x) = u(v(x))$ $\Rightarrow f'(x) = \frac{du}{dv} \frac{dv}{dx}$	Chain rule																																

Let's learn about...

- | | |
|-------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|
| 1 Remember: | 4 The Quotient Rule |
| 2 Differentiation by rule <ul style="list-style-type: none">■ 3. The constant multiple rule■ 4. Sum and Difference Rules | 5 Chain Rule <ul style="list-style-type: none">■ Repeated application |
| 3 The Product Rule | 6 Page 16 of the "log tables" |
| | 7 Exercises |

See also Section 3.3 of **Calculus** by Strang & Herman:
<https://openstax.org/books/calculus-volume-1/pages/3-3-differentiation-rules>

And section 3.6 of the text.

Remember:

Reminders

- ▶ Assignment 2 is open; deadline is 5pm, 13 Oct. You can access it at <https://universityofgalway.instructure.com/courses/46734/assignments/129715>. (Or: go to Canvas, click on Assignments ... Problem Set 2 ... the bottom of the page, click `Load Problem Set 2 in a new window`)
- ▶ This week's **Tutorial Sheet** is available at https://universityofgalway.instructure.com/courses/46734/files/2883465?module_item_id=943734
- ▶ Info on next Tuesday's will follow in an announcement tomorrow morning.

▶ Assignment 3 - posted tomorrow

Differentiation by rule

Yesterday, we saw how to compute derivatives of some functions using the “limit” definition (i.e., **differentiation from first principles**). However, while it is useful to be able to do that for some examples, almost always we use a set of “**rules**” which makes the process much more efficient.

These rules are themselves derived from the “limit” definition – but we don’t have to use that every time.

Differentiation by rule

Yesterday, we looked at two rules:

1. The Constant Rule

If f is a constant function, i.e. $f(x) = c$ for all x , then:

$$\frac{df}{dx} = \frac{d}{dx}(c) = 0.$$

2. The Power Rule

If n is any real number, then

$$\frac{d}{dx}(x^n) = nx^{n-1}.$$

eg

$$f(x) = x^{72}$$
$$f'(x) = 72x^{71}$$

$$f(x) = x^{-3} \Rightarrow f'(x) = (-3)x^{-4}.$$

The constant multiple rule

Let $f(x)$ be any differentiable function, and let k be constant, then

$$\frac{d}{dx}(\underline{kf(x)}) = k \frac{d}{dx}(f(x)).$$

Example: Find the derivative of $f(x) = 5x^4$.

$$\begin{aligned} f(x) &= 5x^4 && (\text{so } k = 5) \\ \frac{df}{dx} = f'(x) &= 5 \underbrace{(x^4)'}_{\substack{\text{ie} \\ \text{derivative of } x^4}} = 5(4x^3) = 20x^3. \end{aligned}$$

u

The Sum and Difference Rules

Let $u(x)$ and $v(x)$ be any differentiable functions. Then

$$\frac{d}{dx}(u(x) + v(x)) = \frac{d}{dx}(u(x)) + \frac{d}{dx}(v(x)).$$

$$\text{Similarly, } \frac{d}{dx}(u(x) - v(x)) = \frac{d}{dx}(u(x)) - \frac{d}{dx}(v(x)).$$

Example: Find the derivative of $f(x) = 1 + x + x^2$.

$$u = (1+x)$$

$$v = x^2$$

$$f(x) = u(x) + v(x)$$

$$f'(x) = u'(x) + v'(x)$$

$$= (1+x)' + (x^2)' = (1)' + (x)' + 2x = 0 + 1 + 2x = 2x + 1$$

Apply rule again!

Ans: $2x + 1$

Actually, the “**Difference Rule**”, which states that

$$\frac{d}{dx}(u(x) - v(x)) = \frac{d}{dx}(u(x)) - \frac{d}{dx}(v(x)).$$

can be combined by combining the **Sum Rule** and the **Constant Multiple Rule**.

$$\begin{aligned} f(x) &= u(x) - v(x) \\ &= u(x) + (-1)v(x) \end{aligned}$$

$$\begin{aligned} \text{Then } f'(x) &= u'(x) + [(-1)v(x)]' && \text{by Sum Rule} \\ &= u'(x) + (-1)v'(x) && \text{by Constant Multiple Rule} \\ &= u'(x) - v'(x) \end{aligned}$$

Example

Suppose that $f(x) = -5x^3 + 3x^2 - 9x + 7$, then find:

- (a) The derivative of $f(x)$;
- (b) The slope of the tangent line at $x = 2$; , i.e. $f'(2)$
- (c) The equation of the tangent at $x = 2$.

(a) $f'(x) = -15x^2 + 6x - 9$

(b) The slope of the tangent line at $x = 2$ is $f'(2)$:

$$f'(2) = -15(2)^2 + 6(2) - 9 = -15(4) + 12 - 9 = -60 + 12 - 9 = -57.$$

- (c) The equation of the line with slope M and passing through a point (x_1, y_1) is

$$y - y_1 = M(x - x_1)$$

$$x_1 = 2 \quad y_1 = f(x_2)$$

The y coordinate at $x = 2$ is

$$\begin{aligned} f(2) &= -5(2)^3 + 3(2)^2 - 9(2) + 7 \\ &= -5(8) + 3(4) - 18 + 7 \\ &= -40 + 12 - 18 + 7 \\ &= -39. \end{aligned}$$

So the tangent line passes through the point $(2, -39)$ and the slope of the line is -57 .

Therefore, the equation of this line is $y + 39 = -57(x - 2)$

Ans: The equation of the tangent line is $x = 2$ is $y = 75 - 57x$.

The Product Rule

We now consider some rules which are a little more complicated and, I think, less obvious.

The first concerns the derivative of the **product** of two functions.

The Product Rule

Let $u(x)$ and $v(x)$ be any differentiable functions. Then

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}.$$

Let's check that this gives the current answer when evaluating the derivative of $f(x) = x^3$ when we set $u(x) = x^2$ and $v(x) = x$:

$$x^3 = (x^2)(x) \quad \text{so}$$

$$u'(x) = 2x$$

$$v'(x) = 1$$

$$\begin{aligned} \frac{d}{dx}(f) &= \frac{d}{dx} \left(\underbrace{(x^2)}_u \underbrace{(x)}_v \right) = (x^2)(1) + x(2x) \\ &= x^2 + 2x^2 \\ &= 3x^2 \quad \checkmark \end{aligned}$$

The Product Rule

The Product Rule

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}.$$

Example Use the **product rule** to find the derivative of

$f(x) = x^3(x^2 + 1)$. We'll do this 2 ways:

1. Without Product Rule: $f(x) = x^3(x^2 + 1)$
 $= x^5 + x^3$
 $f'(x) = 5x^4 + 3x^2$

2. With Product Rule:

$$u(x) = x^3 \quad v(x) = x^2 + 1$$

$$u'(x) = 3x^2 \quad v'(x) = 2x$$

$$\begin{aligned} f'(x) &= u v'(x) + v u'(x) = (x^3)(2x) + (x^2 + 1)(3x^2) \\ &= 2x^4 + 3x^2 + 3x^2 = 5x^4 + 3x^2 \end{aligned}$$

The Product Rule

Example: use the product rule to show that, if $f(x) = x \sin(x)$, then $f'(x) = \underline{x \cos(x) + \sin(x)}$.

$$u(x) = x$$

$$v(x) = \sin(x)$$

$$u'(x) = 1$$

$$v'(x) = \cos(x)$$

$$\begin{aligned} f'(x) &= u(x) v'(x) + v(x) u'(x) \\ &= x \cos(x) + \sin(x)(1) \end{aligned}$$



The Quotient Rule

The Quotient Rule

If u and v are differentiable at x and if $v(x) \neq 0$, then $f(x) = \frac{u(x)}{v(x)}$ is differentiable at x , and

$$\frac{d}{dx} \left(\frac{u}{v} \right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}.$$

Example: Use this rule to find the derivative of $f(x) = \frac{x+1}{x^2}$

$$u(x) = x+1$$

$$v(x) = x^2$$

$$u'(x) = 1$$

$$v'(x) = 2x.$$

$$\begin{aligned} \frac{df}{dx} &= \frac{v(x) u'(x) - u(x) v'(x)}{(v(x))^2} = \frac{(x^2)(1) - (x+1)(2x)}{(x^2)^2} \\ &= \frac{x^2 - 2x^2 - 2x}{x^4} = \frac{-x^2 - 2x}{x^4} = -x^{-2} - 2x^{-3}. \end{aligned}$$

The Quotient Rule

Example

We know that

- ▶ $\tan(x) = \frac{\sin(x)}{\cos(x)}$,
- ▶ $\sin^2(x) + \cos^2(x) = 1$ ✓
- ▶ $\sin'(x) = \cos(x)$ and $\cos'(x) = -\sin(x)$.

Use these facts, and the Quotient Rule to show that

$$\frac{d}{dx}(\tan(x)) = \left(\frac{1}{\cos(x)}\right)^2.$$

The Quotient Rule

$$f(x) = \tan(x) = \frac{\sin(x)}{\cos(x)}.$$

$$u(x) = \sin(x)$$

$$v(x) = \cos(x)$$

$$u'(x) = \cos(x)$$

$$v'(x) = -\sin(x).$$

Apply the Quotient Rule:

$$\frac{df}{dx} = \frac{v u'(x) - u v'(x)}{(v(x))^2} = \frac{\cos(x)\cos(x) - \sin(x)(-\sin(x))}{\cos^2(x)}$$

$$= \frac{\cos^2(x) + \sin^2(x)}{\cos^2(x)} = \frac{1}{\cos^2(x)}.$$

$$=: \sec^2(x)$$

Chain Rule

Of all the differentiation rules, the **chain rule** is the most important: most other rules are actually just special cases of it. It applies to a “function of a function”

The Chain Rule

If $u(x)$ and $v(x)$ are differentiable, and f is the composite function $f(x) = u(v(x))$, then

$$\frac{df}{dx} = \frac{du}{dv} \frac{dv}{dx}.$$

$u(v(x))$ is not $u(x)v(x)$.

Eg, if $f(x) = (2x+1)^3 = u(v(x)) (= u \circ v(x))$
then $u(v) = v^3$ $v(x) = 2x+1$.

Chain Rule

The Chain Rule

If $f(x) = u(v(x))$, then

$$\frac{df}{dx} = \frac{du}{dv} \frac{dv}{dx}.$$

Example: What is the derivative of $f(x) = \cos(x^2)$?

$$f(x) = \cos(\underline{x^2})$$

$$u(v) = \cos(v)$$

$$v(x) = x^2$$

$$\frac{du}{dv} = -\sin(v)$$

$$\frac{dv}{dx} = 2x.$$

$$\begin{aligned} \text{So } \frac{df}{dx} &= \frac{du}{dv} \cdot \frac{dv}{dx} = -\sin(v) \cdot 2x. \\ &= -\sin(x^2) \cdot 2x = -2x \sin(x^2) \end{aligned}$$

Chain Rule

Example

Find $\frac{dy}{dx}$ if $y = (x^3 + 4x^4 + 7)^{12}$.

Example: Let $u(v) = v^{12}$ and $v(x) = x^3 + 4x^4 + 7$, then y is $y = u(v(x))$.

Note that

$$\frac{du}{dv} = 12v^{11} \quad \text{and} \quad \frac{dv}{dx} = 3x^2 + 16x^3.$$

By the Chain Rule we have

$$\frac{dy}{dx} = \frac{du}{dv} \frac{dv}{dx} = 12v^{11}(3x^2 + 16x^3),$$

and therefore

$$\frac{dy}{dx} = 12(x^3 + 4x^4 + 7)^{11}(3x^2 + 16x^3).$$

Example (Not done in detail in class)

Find $\frac{dy}{dx}$ if $y = \frac{1}{(x^4 + 2x^2 + 8)^{40}}$.

We have $y = (x^4 + 2x^2 + 8)^{-40}$. We can write y as $y(x) = u(v(x))$ with

► $u(v) = \underline{v^{-40}}$ and so $\frac{du}{dv} = -40v^{-41}$; and

► $v(x) = \underline{x^4 + 2x^2 + 8}$, so $\frac{dv}{dx} = 4x^3 + 4x$.

Applying the Chain Rule: $\frac{dy}{dx} = \frac{du}{dv} \frac{dv}{dx}$, we get

$$\frac{dy}{dx} = -40v^{-41}(4x^3 + 4x) = \frac{-40(4x^3 + 4x)}{(x^4 + 2x^2 + 8)^{41}}$$

Often we apply the **Chain Rule** to “functions of functions of functions”: if $y(x) = t(u(v(x)))$, then

$$\frac{dy}{dx} = \frac{dt}{du} \frac{du}{dv} \frac{dv}{dx}$$

Example

Find $\frac{dy}{dx}$ when $y = \sin^4(x^5 + 7)$.

Example

Show that the derivative of $y = \cos^2(1/x)$ is

$$\frac{dy}{dx} = 2 \frac{\sin(1/x) \cos(1/x)}{x^2}$$

Calcalas

Diorthaigh

$f(x)$	$f'(x)$
x^n	nx^{n-1}
$\ln x$	$\frac{1}{x}$
e^x	e^x
e^{ax}	ae^{ax}
a^x	$a^x \ln a$
$\cos x$	$-\sin x$
$\sin x$	$\cos x$
$\tan x$	$\sec^2 x$
$\cos^{-1} \frac{x}{a}$	$-\frac{1}{\sqrt{a^2 - x^2}}$
$\sin^{-1} \frac{x}{a}$	$\frac{1}{\sqrt{a^2 - x^2}}$
$\tan^{-1} \frac{x}{a}$	$\frac{a}{a^2 + x^2}$

Calculus

Derivatives

Riail an toraidh	$y = uv$ $\Rightarrow \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$	Product rule
Riail an lín	$y = \frac{u}{v}$ $\Rightarrow \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$	Quotient rule
Cuingriail	$f(x) = u(v(x))$ $\Rightarrow f'(x) = \frac{du}{dv} \frac{dv}{dx}$	Chain rule

Exercises

Exercises 4.2.1 (Based on Q2(a), 2023/2024)

Find the derivative of $f(x) = \frac{\sin(x)}{\sqrt{x}}$.

Exercise 4.2.2 (Based on Q2(b), 2019/2020)

Find the derivative of $f(x) = \frac{x^2 + x - 2}{x^3 + 6}$.

Exercise 4.2.3

Find the derivative of

1. $f(x) = x^3 \cos(x^2)$
2. $f(x) = \tan^3(\sin^2(x^4))$