


Week 2, Lecture 1 Partial Fractions

Dr Niall Madden

School of Mathematical and Statistical Sciences, University of Galway

24 September, 2024



This version of the slides are by Niall Madden, but are adapted from original notes by Dr Kirsten Pfeiffer.

Outline

1 News!

- Tutorials
- Assignments

2 Partial Fractions

- Case 1
- Case 2
- Case 3
- Case 4
- Exercises

For more, see Section 2.5 (Rational Functions) of *Modern Engineering Mathematics*:

https://search.library.nuigalway.ie/permalink/f/3b1kce/TN_cdi_askewsholts_vlebooks_9780273742517

Tutorials start **this** week. The schedule is:

- ▶ Teams 1+2: Tuesday 15:00 ENG-**2003**
- ▶ Teams 3+4: Tuesday 15:00 ENG-**2034**
- ▶ Teams 9+10: Thursday 11:00 ENG-**2002**
- ▶ Teams 11+12: Thursday 11:00 ENG-**3035**
- ▶ Teams 5+6: Friday 13:00 Eng-**2002**
- ▶ Teams 7+8: Friday 13:00 Eng-**2035**

If you are interested to taking a tutorial through Irish, please complete this survey: <http://tinyurl.com/suairbhe1>

- ▶ There is currently a “practice” assignment open. See <https://universityofgalway.instructure.com/courses/35693/assignments/94873>
- ▶ During tutorials, the tutor will solve some similar questions. You can access the tutorial sheet at https://universityofgalway.instructure.com/courses/35693/files/2023552?module_item_id=650912
- ▶ A new assignment will open by tomorrow...

.....

In addition, each week I'll post a set of exercises related to the material covered. You don't have to submit your work for these, but you should try them: they are similar in style and standard to exam questions.

Partial Fractions

Rational Functions have the general form $f(x) = \frac{p(x)}{q(x)}$, where $p(x)$ and $q(x)$ are polynomials.

An (proper) rational function can often be written as a sum of simpler ones: **partial fractions**.

For example

$$\frac{8x - 12}{x^2 - 2x - 3}$$

can be written as

$$\frac{3}{x - 3} + \frac{5}{x + 1}$$

In order to do this, we try to **factorize** the denominator.

Partial Fractions

Note: Any polynomial (with real coefficients) can be factorised fully into the product of

- ▶ linear
- ▶ and irreducible quadratic factors.

We get different combinations of factors in the denominator. Let's look at **four cases**, and how to find the partial fractions in each case.

Partial Fractions

The four cases

1. Linear factors to the power of 1 in the denominator.
2. Linear factors to the power greater than 1 in the denominator, (i.e repeated linear factors).
3. Irreducible quadratic factors.
4. Irreducible quadratic factors to power greater than 1.

(1) Linear factors to the power of 1 in the denominator.

Example

$$\frac{3x}{(x-1)(x+2)}$$

There are **two methods** for finding A and B .

Method 1: Comparing coefficients

Method 2: Substituting specific values for x .

Example

Write $\frac{8x - 12}{x^2 - 2x - 3}$ as sum of partial fractions.

(2) Linear factors to the power greater than 1 in the denominator, (i.e repeated linear factors).

If $(x - \alpha)^k$ appears in the denominator, it will give rise to the following terms:

$$\frac{A_1}{x - \alpha} + \frac{A_2}{(x - \alpha)^2} + \dots + \frac{A_k}{(x - \alpha)^k}$$

Example

Find A , B and C such that

$$\frac{3x + 1}{(x - 1)^2(x + 2)} = \frac{A}{x - 1} + \frac{B}{(x - 1)^2} + \frac{C}{x + 2}$$

(Note: we'll find that $A = 5/9$, $B = 4/3$ and $C = -5/9$).

(3) Irreducible quadratic factors.

Irreducible quadratic factors can not be factorised using real numbers, e.g. $x^2 + x + 1$.

An irreducible quadratic factor $ax^2 + bx + c$ gives rise to partial fractions of the form

$$\frac{Ax + B}{ax^2 + bx + c}.$$

Example 2.34 from textbook

If one writes

$$\frac{5x}{(x^2 + x + 1)(x - 2)} = \frac{Ax + B}{x^2 + x + 1} + \frac{C}{x - 2}$$

then we find $A = 10/7$, $B = 5/7$ and $C = 10/7$.

(4) Irreducible quadratic factors to power greater than 1.

Each repeated irreducible quadratic factor $(ax^2 + bx + c)^k$ in the denominator will give rise to

$$\frac{A_1x + B_1}{ax^2 + bx + c} + \frac{A_2x + B_2}{(ax^2 + bx + c)^2} + \cdots + \frac{A_kx + B_k}{(ax^2 + bx + c)^k}.$$

These can be done in a similar way to the previous case. But the calculations are pretty messy, so we won't even try!

Exercise 2.1.1

Find the constants A , B and C , so that

$$\frac{2x+1}{(x-2)(x+1)(x-3)} = \frac{A}{x-2} + \frac{B}{x+1} + \frac{C}{x-3}$$

Exercise 2.1.2

Express the following as partial fractions.

1. $\frac{6}{x^2 - x - 2}$

2. $\frac{2x-1}{x^2 - x - 2}$

3. $\frac{x-1}{(x+1)(x^2 - x - 2)}$

4. $\frac{x}{x^2 + 2x + 1}$