


MA211

Lecture 3: Limits

Dr Niall Madden

Monday 15 September 2008



Outline

Problem Solving Sessions

Problem Solving sessions (tutorials) will start next week. There will be three per week. Attend whichever one you like

- Tuesday, 3pm, AC202
- Wednesday, 5pm, QA003 (Physiology lecture room)
- ????? (Thursday, 6pm, IT207??)

A a **function** f from a set X to a set Y is a rule of correspondences that associates every element of X with some (single) element of Y . We write:

$$f : X \rightarrow Y.$$

- X is called the **Domain** of f , and
- Y is called the *Codomain*.
- the *Range* of f the subset of Y that contains all the elements of Y that are the image under f of some element of X .

A function from X to Y is *One-to-One* if no two elements of X are mapped to the same element of Y . (In some books this is called *injective*).

Definition (One-to-one)

The function $f : X \rightarrow Y$ is one-to-one if whenever $f(x_1) = f(x_2)$ then $x_1 = x_2$.

Example

Is the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = x^2$ is one-to-one? Why?

Find other sets $X \subseteq \mathbb{R}$ and $Y \subseteq \mathbb{R}$ such that $f : X \rightarrow Y$ is one-to-one.

A function from X to Y is *onto* if every element of Y is the image of some element of X . (In some books this is called *surjective*).

Definition (Onto)

The function $f : X \rightarrow Y$ is onto if for each $y \in B$ there exists $x \in A$ such that y is the image of x . That is

$$\forall y \in B, \exists x \in A \text{ such that } f(x) = y.$$

A third way of expressing this is by saying the *range* of f is equal to its *codomain*.

Example

Show that the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = x^2$ is not **onto**.

Find other sets $X \subseteq \mathbb{R}$ and $Y \subseteq \mathbb{R}$ such that $f : X \rightarrow Y$ is onto.

Exercise (3.1)

Give an example of a function:

- (i) $f : \mathbb{Z} \rightarrow \mathbb{N}$ that is onto but *not* one-to-one.
- (ii) $f : \mathbb{N} \rightarrow \mathbb{N}$ that is one-to-one, but not onto.

Inverse functions

When a function is both **one-to-one** and **onto** it has an **Inverse**.

Definition (Inverse)

If the function g is the **inverse** of f then

when $f(x) = y$, we get that $g(y) = x$.

Usually we write $g = f^{-1}$.

Examples:

$$1 \quad y = f(x) \iff x = f^{-1}(y)$$

2 The domain of f^{-1} is the range of f

3 The range of f^{-1} is the domain of f

$$4 \quad f^{-1}(f(x)) = x$$

$$5 \quad f(f^{-1}(x)) = x$$

$$6 \quad (f^{-1})^{-1} = f.$$

7 The graph of f^{-1} is the reflection of the graph of f in the line $x = y$.

The trigonometric functions


You'll remember the definition of \cos and \sin in terms of a right-angle triangle.

Here is another, equivalent definition.

- Take the usual coordinate axes centred on $(0, 0)$ and draw the unit circle,
- Mark the point $(1, 0)$.
- Trace an arc of length t *anti-clockwise* from $(1, 0)$ along the circle. Call the point at the end of that arc P .
- The x -coordinate of P is $\cos(t)$ and the y -coordinate is $\sin(t)$.

The trigonometric functions

The cos (top) and sin (bottom) functions



The trigonometric functions

Exercise (3.2)

Find subsets X and Y of the real numbers such that the functions $f : X \rightarrow Y$ are *invertible* (i.e., both *one-to-one* and *onto*) for

- (i) $f(x) = \sin(x)$
- (ii) $f(x) = \cos(x^2)$

Limits

When we write

$$\lim_{x \rightarrow c} f(x) = L$$

or say “*The limit of f as x approaches c is L* ” we mean that we can make f as close to L as we would like by taking x as close to c as is needed.

Definition (Limit)

If for any $\varepsilon > 0$, no matter how small, we can find $\delta > 0$ such that

$$|f(x) - L| < \varepsilon \quad \text{when} \quad |x - c| < \delta.$$

then we can say

$$\lim_{x \rightarrow c} f(x) = L.$$

Example

Show that

$$\lim_{x \rightarrow 3} (2x + 1) = 7.$$

Exercise (3.3)

Show *carefully* that

(i) $\lim_{x \rightarrow 4} 3x - 7 = 5.$

(ii) $\lim_{x \rightarrow 2} \left(\frac{x}{2} + 3 \right)$ is 4

Calculating the limit of a given function as it approaches a certain point is a fairly standard task.

Example

Find the limit of

$$f(x) = \frac{2x^2 - 3x - 2}{x - 2}$$

as x approaches 2.

Let n be an integer, k a constant real number, and f and g be functions that have a limit at c . Then

$$1 \quad \lim_{x \rightarrow c} k = k;$$

$$2 \quad \lim_{x \rightarrow c} x = c;$$

$$3 \quad \lim_{x \rightarrow c} kf(x) = k \lim_{x \rightarrow c} f(x);$$

$$4 \quad \lim_{x \rightarrow c} (f(x) + g(x)) = \lim_{x \rightarrow c} f(x) + \lim_{x \rightarrow c} g(x).$$

$$5 \quad \lim_{x \rightarrow c} (f(x) - g(x)) = \lim_{x \rightarrow c} f(x) - \lim_{x \rightarrow c} g(x).$$

$$7 \quad \lim_{x \rightarrow c} (f(x) \cdot g(x)) = \lim_{x \rightarrow c} f(x) \cdot \lim_{x \rightarrow c} g(x).$$

$$8 \quad \lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow c} f(x)}{\lim_{x \rightarrow c} g(x)} \quad \text{providing that} \quad \lim_{x \rightarrow c} g(x) \neq 0.$$

$$9 \quad \lim_{x \rightarrow c} \left(f(x) \right)^n = \left(\lim_{x \rightarrow c} f(x) \right)^n.$$

$$10 \quad \lim_{x \rightarrow c} \left(f(x) \right)^{(1/n)} = \left(\lim_{x \rightarrow c} f(x) \right)^{(1/n)}.$$

There are many neat tricks for computing limits, particularly those of the form

$$\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow c} f(x)}{\lim_{x \rightarrow c} g(x)}, \quad \text{where} \quad \lim_{x \rightarrow c} g(x) = 0$$

One of these is *The Squeeze Theorem*, but even better is l'Hopital's Rule

We'll cover these on Wednesday.