


Week 03, Lecture 2 Vertical Asymptotes and Continuity

Dr Niall Madden

University of Galway

Wednesday, 1 October, 2025



This slides are by Niall Madden, with some content based on notes by Dr Kirsten Pfeiffer, and from Strang & Herman's "Calculus".

Outline

- 1 Reminders...
- 2 Recall: One-sided Limits
 - Notation
 - Existence of a limit
- 3 Vertical Asympotes
 - Horizontal Asymptotes
- 4 Continuity
- 5 Types of discontinuity
- 6 Exercises

For more, see Chapter 2 (Limits) in **Calculus** by Strang & Herman. See openstax.org/books/calculus-volume-1/pages/2-introduction

In particular §2.2: One-sided limits (§2.2.4+§2.2.5) and vertical asymptotes (§2.2.7).

Reminders

- ▶ **Assignment 1** due 5pm, Monday 6 October. You may access it multiple times, by clicking on **Assignments ... Problem Set 1 ...** and then, at the bottom of the page:

Load Problem Set 1 in a new window

- ▶ The **Tutorial Sheet** is available at <https://universityofgalway.instructure.com/courses/46734/files/2883465?wrap=1>
- ▶ Assignment 2 is also open; deadline is 5pm, 13 Oct.
- ▶ The first (of two) class tests will take place Tuesday, 14th October.
- ▶ If you wish to avail of Reasonable Accommodations for it tests, please complete this form: <https://forms.office.com/e/HaAsrzaE3D> by **10am Thursday 2nd Oct.**

Yesterday we met the concept of **one-sided limits**:

$\lim_{x \rightarrow a^-} f(x)$ is: **limit of f as x approaches a from the left**

$\lim_{x \rightarrow a^+} f(x)$ is: **limit of f as x approaches a from the right**

These mean that

- ▶ if $\lim_{x \rightarrow a^-} f(x) = L$, then we can make $f(x)$ as close to L as we would like by taking $x < a$ as close to a as needed.
- ▶ If $\lim_{x \rightarrow a^+} f(x) = L$, then we can make $f(x)$ as close to L as we would like by taking $x > a$ as close to a as needed.

Existence of a limit

$\lim_{x \rightarrow a} f(x)$ **exists** if and only if

$$\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x)$$

So if $\lim_{x \rightarrow a} f(x) = L$ exists, we have

$$\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a^+} f(x) = L$$

though it is not necessary that $f(a) = L$

Example

Sketch the function


$$f(x) = \begin{cases} 3 - x, & x < 2 \\ 4, & x = 2 \\ \frac{x}{2}, & x > 2 \end{cases}$$

Determine if $\lim_{x \rightarrow 2} f(x)$ exists.

Vertical Asymptotes

Let's revisit the following example from yesterday:

$$f(x) = \frac{1}{(x-1)^2}$$



Note that the points on the graph having x -coordinates very near to 1 are very close to the vertical line $x = 1$. That is, as x approaches 1, the points on the graph of $f(x)$ are closer to the line $x = 1$.

We call the line $x = 1$ a **vertical asymptote** of the graph.

Vertical Asymptotes

Definition: Vertical Asymptote

The vertical line $x = a$ is a **vertical asymptote** of $f(x)$ if any of $\lim_{x \rightarrow a^-} f(x)$, $\lim_{x \rightarrow a^+} f(x)$, or $\lim_{x \rightarrow a} f(x)$ are ∞ or $-\infty$.

To find a vertical asymptote of a function $f(x) = \frac{p(x)}{q(x)}$, we find a value, a for which $p(a) \neq 0$ but $q(a) = 0$.

Vertical Asymptotes


Example

Find any vertical asymptotes of

$$f(x) = \frac{x^2 - x - 6}{x + 1}$$

Vertical Asymptotes

$$f(x) = \frac{x^2 - x - 6}{x + 1}$$



Vertical Asymptotes


Example

Find all vertical asymptotes of the graph of

$$g(x) = -\frac{8}{x^2 - 4}.$$

Vertical Asymptotes

$$f(x) = -\frac{8}{x^2 - 4}$$



There is a related concept of a **horizontal asymptote**, but we'll save that for later, when we cover “limits at infinity”.

Continuity

Many functions have the property that you can trace their graphs with pen and paper, without lifting the pen from the page. Such functions are called **continuous**.

Some other functions have points where you have to lift the pen occasionally. We say they have a **discontinuity** at such points.

Intuitively, a function is continuous at a particular point if there is no **break** (or “**jump**”) in its graph at that point.

More formally, we define continuity in terms of **limits**

Continuity

Definition

A function f is **continuous at** $x = a$ if

1. $f(a)$ is defined, i.e., a is in the domain of f ,
2. $\lim_{x \rightarrow a} f(x)$ exists.
3. $\lim_{x \rightarrow a} f(x) = f(a)$.

If $f(x)$ is not continuous at $x = a$ we say it is **discontinuous** at $x = a$.

If f is continuous **at every point** in its domain, we say f **is continuous**.

Many functions are continuous, e.g. all polynomial functions, **most** trigonometric functions (not **tan**), $|x|$, and so on.

Continuity

Example 1

Determine if $f(x) = \frac{x^2 - 4}{x - 2}$ is continuous at $x = 2$.

Example 2

Determine if $f(x) = \begin{cases} 1 - x & x \leq 0 \\ 2 + x & x > 0 \end{cases}$ is continuous at $x = 0$.

Example 3

Determine if $f(x) = \begin{cases} \frac{\sin x}{x} & x \neq 0 \\ 1 & x = 0 \end{cases}$ is continuous at $x = 0$.

Example

Consider the function

$$f(x) = \begin{cases} x + 1, & x < 2 \\ bx^2, & x \geq 2 \end{cases}$$

For what value of b is f continuous at $x = 2$?

Continuity

Example

For what values of x is $f(x) = \frac{2x + 1}{2x - 2}$ continuous?

Types of discontinuity

We have encountered three types of discontinuity.

- ▶ **Removable discontinuity:** $\lim_{x \rightarrow a} f(x)$ exists but

$$\lim_{x \rightarrow a} f(x) \neq f(a)$$

- ▶ **Jump discontinuity:** $\lim_{x \rightarrow a^-} f(x)$ and $\lim_{x \rightarrow a^+} f(x)$ both exist (and are finite), but $\lim_{x \rightarrow a^-} f(x) \neq \lim_{x \rightarrow a^+} f(x)$
- ▶ **Infinite discontinuity:** At least one of the one-sided limits does not exist.

Types of discontinuity

Example

Each of the following functions has a discontinuity at $x = 2$.
Classify it.

1. $f(x) = \frac{x^2 - 4}{x - 2}$

2. $g(x) = \frac{x^2}{x - 2}$

3. $h(x) = \begin{cases} x/2 & x < 2 \\ -2 & x = 2 \\ x^2 - 3 & x > 2. \end{cases}$

4. $h(x) = \begin{cases} x/2 & x < 2 \\ x^2 - 2 & x > 2. \end{cases}$

Exercises

Exercise 3.2.1

Find all the vertical asymptotes of $f(x) = \frac{x+2}{x^2+2x-8}$.

Exercises 3.2.2 (Based on Q1(a), 23/24)

$$\text{Let } g(x) = \begin{cases} 3 & x \leq 0 \\ 2x+1 & 0 < x < 1 \\ x^2 & x \geq 1. \end{cases}$$

- (i) Sketch the graph of $g(x)$ on the interval $[-3, 4]$, making use of the empty and full circle notation.
- (ii) Compute $\lim_{x \rightarrow 1^-} g(x)$ and $\lim_{x \rightarrow 1^+} g(x)$. Is g continuous at $x = 1$. If not, classify the type of discontinuity.

Exercises

Exercise 3.2.3

For what values of b and c is $f(x) = \begin{cases} x^2 + 1 & x \leq -1 \\ x + b & -1 < x < 1 \\ cx^2 & x \geq 1. \end{cases}$
continuous at $x = -1$ and $x = 1$?