


Week 1, Lecture 3: Polynomials and Partial Fractions

Dr Niall Madden

School of Mathematical and Statistical Sciences, University of Galway

Thursday, 19 September, 2024



This version of the slides are by Niall Madden, but are adapted from original notes by Dr Kirsten Pfeiffer.

Outline

1 News!

- Tutorials
- Exercise sheet

2 Functions (again)

- Recall...

3 Polynomials

- Sketching polynomials

4 Rational Functions

- Long division

5 Partial Fractions

For more, see Sections 2.4 (Polynomials) 2.5 (Rational Functions) of *Modern Engineering Mathematics*:

https://search.library.nuigalway.ie/permalink/f/3b1kce/TN_cdi_askewsholts_vlebooks_9780273742517

Tutorials start next week. Here is the schedule:

- ▶ Teams 1+2: Tuesday 15:00 ENG-**2003**
- ▶ Teams 3+4: Tuesday 15:00 ENG-**2034**
- ▶ Teams 9+10: Thursday 11:00 ENG-**2002**
- ▶ Teams 11+12: Thursday 11:00 ENG-**3035**
- ▶ Teams 5+6: Friday 13:00 Eng-**2002**
- ▶ Teams 7+8: Friday 13:00 Eng-**2035**

Note: I think the schedule is correct, but the venues are not confirmed... An announcement will be posted to Canvas on Monday confirming.

Would you be interested to taking a tutorial through Irish? If so, please complete this survey: <https://tinyurl.com/suirbhel>

You don't have to complete a graded assignment next week. However, this is a “practice” one available. See <https://universityofgalway.instructure.com/courses/35693/assignments/94873>

During tutorials, the tutor will solve some similar questions. You can access the tutorial sheet at https://universityofgalway.instructure.com/courses/35693/files/2023552?module_item_id=650912

Yesterday, we learned that

- ▶ A **function** is a rule for mapping from elements of one set (the domain) to elements of another (the codomain).
- ▶ When we write $y = f(x)$, we say “ x ” is the **argument** of the function.
- ▶ When $y = f(x)$ for some $x \in X$, y is said to be the **image** of x under f .
- ▶ The set of all images $y = f(x), x \in X$, is called the **range** of f .

Polynomials

A **polynomial function** (or just **polynomial**) is a function of the form

$$y = f(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x^1 + a_0, \quad x \in \mathbb{R},$$

where a_0, a_1, \dots, a_n are real numbers called the **coefficients** of the polynomial.

The number n is called the **degree** of the polynomial.

Polynomials

Example: linear

$y = x$ is a **linear** polynomial with degree $n = 1$.

Example: quadratic

$x^2 - 2x - 3$ is a **quadratic** polynomial with degree $n = 2$.

There are many occasions when we want to **factorise** such quadratics, meaning we write them as the product of a pair of linear polynomials.

For example, we can **factorise** $x^2 - 2x - 3$ as

$$x^2 - 4x + 3 = (x - 3)(x + 1)$$

Polynomials


It is important to note that not all quadratic polynomials can be factorised as two linear polynomials. Such quadratics are called **irreducible**.

For example, $x^2 + 1$ is irreducible.

Polynomials

Example

$y = x^3 - 4x^2 + x + 6$ is a **cubic** function with degree $n = 3$.



Fact: A polynomial function of grade n has **up to** $n - 1$ turning points (“bends”).

Examples:

Break Time

During the break, think and talk about what you might do to sketch the graph of

$$y = -x^3 + x^2 + 2x$$

- ▶ To sketch the graph, first find the **intercepts**:
 - ▶ The **y-intercepts** can be found by letting $x = 0$.
 - ▶ The **x-intercepts** are called the **roots** (or **zeros**).
To find the roots, set y equal to zero and solve for x .
- ▶ You don't have to use the same scale on the x - and on the y -axis.
- ▶ Do not use graph paper.


Example

Sketch the graph of

$$y = -x^3 + x^2 + 2x$$

How to sketch $y = -x^3 + x^2 + 2x$

Actual plot of $y = -x^3 + x^2 + 2x$



Exercise 1.3.1

Sketch the graphs of

(i) $y = 5x^2 - 7$

(ii) $y = x^2 - 4x + 3$

(iii) $y = x^3 - 6x^2 - 11x - 6$

Rational Functions

Rational Functions have the general form

$$f(x) = \frac{p(x)}{q(x)},$$

where $p(x)$ and $q(x)$ are polynomials.

- ▶ If degree of $p(x) <$ degree of $q(x)$,
 $f(x)$ is called a **strictly proper rational function**.
- ▶ If degree of $p(x) =$ degree of $q(x)$,
 $f(x)$ is called a **proper rational function**.
- ▶ If degree of $p(x) >$ degree of $q(x)$,
 $f(x)$ is called an **improper rational function**.

Rational Functions

An improper or proper rational function can always be expressed as a polynomial plus a strictly proper rational function, for example by algebraic division.

Example

$$\frac{4x^3 + 4x^2 + 4}{x^2 - 3} = 4x + 4 + \frac{12x + 16}{x^2 - 3}$$

For the previous example, we can work this out ourselves using **Long Division** to divide numerator by denominator:

Example 2.30 from text book

Use long division to show that

$$\frac{3x^4 + 2x^3 - 5x^2 + 6x - 7}{x^2 - 2x + 3} = 3x^2 + 8x + 2 - \frac{14x + 13}{x^2 - 2x + 3}$$

Partial Fractions

A (proper) rational function can often be written as a sum of simpler ones: **partial fractions**.

For example

$$\frac{8x - 12}{x^2 - 2x - 3}$$

can be written as

$$\frac{3}{x - 3} + \frac{5}{x + 1}$$

We verified this in class. Next week, we see **how to compute partial fractions?**