
2526-MA140: Week 03, Lecture 1 (L04)

Partial Fractions Dr Niall Madden

University of Galway

Tuesday, 23 September, 2025



Outline

- 1 News!
 - Tutorials
 - Assignments
 - Class tests
- 2 Partial Fractions
 - 4 cases
 - Case 1
 - Case 2
 - Case 3
 - Case 4

- 3 Towards Limits
- 4 Exercises



News! Tutorials

Tutorials start this week. The schedule is:

Teams	Time	Venue	Leader
1, 2	Tuesday 15:00	ENG- 2003	ST
3, 4	Tuesday 15:00	ENG- 2034	JM
9, 10	Thursday 11:00	ENG- 2002	ST
11, 12	Thursday 11:00	ENG- 3035	JM
5, 6	Friday 13:00	Eng- 2002	ST
7, 8	Friday 13:00	Eng- 2035	JM

Rang teagaisc trí Ghaeilge (Irish tutorial): Dé Máirt (Tuesday) 15:00, Áras na Gaeilge 221.

- ► There is currently a "practice" assignment open. See https://universityofgalway.instructure.com/courses/46734/assignments/128373
- During tutorials, the tutor will solve some similar questions. You can access the **tutorial sheet** at https://universityofgalway.instructure.com/courses/46734/files/2842617?module_item_id=925893. You can also access this through the Canvas page: Modules... Tutorial Sheets.
- ► **Assignment 1** will be due 5pm, Monday 5 October.

Also: try the exercises at the end of each set of lecture slides: they are similar in style and standard to exam questions.

News! Class tests

There are two class test planned for this module:

- MCQ format;
- ▶ both worth 10% of the final grade;
- ► Test 1: **Tuesday, 14 October** (Week 5)
- ► Test 2: Tuesday, 18 November (Week 10)
- Contact Niall if you have any concerns, or wish to avail of alternative arrangements, as provided by LENS reports.

Partial Fractions

Rational Functions have the general form $f(x) = \frac{p(x)}{q(x)}$, where p(x) and q(x) are polynomials.

An (proper) rational function can often be written as a sum of simpler ones: partial fractions.

For example

$$\frac{8x-12}{x^2-2x-3}$$

can be written as

$$\frac{3}{x-3} + \frac{5}{x+1}$$

Check: (next slide)

Partial Fractions

Partial Fractions

Note: Any polynomial (with real coefficients) can be factorised fully into the product of

- ▶ linear
- ▶ and irreducible quadratic factors.

Examples:

We get different combinations of factors in the denominator. Let's look at **four cases**, and how to find the partial fractions in each case.

The four cases

- 1. Denominator has linear factors to the power of 1
- 2. Denominator has factors to the power greater than 1 (i.e repeated linear factors).
- 3. Denominator has irreducible quadratic factors.
- 4. Denominator has irreducible quadratic factors to power greater than 1.

Case 1: Linear factors to the power of 1 in the denominator.

We have **two methods** to find A and B.

Method 1: Comparing coefficients

Example

$$\overline{(x-1)(x+2)}$$

Method 2: Substituting specific values for x.

Example

Write
$$\frac{8x-12}{x^2-2x-3}$$
 as sum of partial fractions.

(2) Linear factors to the power greater than 1 in the denominator, (i.e repeated linear factors).

If $(x - \alpha)^k$ appears in the denominator, it will give rise to the following terms:

$$\frac{A_1}{x-\alpha} + \frac{A_2}{(x-\alpha)^2} + \dots + \frac{A_k}{(x-\alpha)^k}$$

Example

Find A, B and C such that

$$\frac{3x+1}{(x-1)^2(x+2)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+2}$$

(Note: we'll find that A = 5/9, B = 4/3 and C = -5/9).

(3) Irreducible quadratic factors.

Irreducible quadratic factors can not be factorised using real numbers, e.g. $x^2 + x + 1$.

An irreducible quadratic factor $ax^2 + bx + c$ gives rise to partial fractions of the form

$$\frac{Ax+B}{ax^2+bx+c}.$$

Example 2.34 from textbook

If one writes

$$\frac{5x}{(x^2+x+1)(x-2)} = \frac{Ax+B}{x^2+x+1} + \frac{C}{x-2}$$

then we find A - 10/7, B = 5/7 and C = 10/7.

(4) Irreducible quadratic factors to power greater than 1.

Each repeated irreducible quadratic factor $(ax^2 + bx + c)^k$ in the denominator will give rise to

$$\frac{A_1x + B_1}{ax^2 + bx + c} + \frac{A_2x + B_2}{(ax^2 + bx + c)^2} + \dots + \frac{A_kx + B_k}{(ax^2 + bx + c)^k}.$$

These can be done in a similar way to the previous case. But the calculations are pretty messy, so we won't even try!

Towards Limits

When we were considering the domain of a function, we looked at those *x*-values for which the function was not defined.

Example

$$f(x) = \frac{x^2 - 2}{x - 1}$$

$$g(x) = \frac{x^2 - 1}{x - 1}$$

Neither f nor g are defined at x = 1.

But what happens if x gets very closed to 1?

X	0.900	0.990	0.999	1	1.001	1.010	1.100
f(x)							
g(x)							


Let's look at the graphs of f and g.

Towards Limits

Example

$$f(x) = \frac{x^2 - 2}{x - 1}$$

$$g(x) = \frac{x^2 - 1}{x - 1}$$



Towards Limits

In the previous example, we saw that, although neither f nor g was defined at x = 1, they behaved very differently as x approaches 1.

To discuss this we'll need the concept of a **limit** which, roughly, relates to the value of function as it **approaches** a point (but not actually at that point).

$$\lim_{x \to a} f(x) = L$$

The concept of a limit is a prerequisite for a proper understanding of calculus and numerical methods.

Exercises

Exercise 2.1.1

Find the constants A, B and C, so that

$$\frac{2x+1}{(x-2)(x+1)(x-3)} = \frac{A}{x-2} + \frac{B}{x+1} + \frac{C}{x-3}$$

Exercises

Exercise 2.1.2

Express the following as partial fractions.

1.
$$\frac{6}{x^2 - x - 2}$$

2.
$$\frac{2x-1}{x^2-x-2}$$

3.
$$\frac{x-1}{(x+1)(x^2-x-2)}$$

4.
$$\frac{x}{x^2 + 2x + 1}$$

5.
$$\frac{1}{\sqrt{3}-1}$$

Exercises

Exercise 2.1.3 (MA140 Exam, 24/25)

Express $\frac{3x+1}{x^2-x-2}$ as partial fractions.