

Interior layers in singularly perturbed time dependent problems

E. O’Riordan*

* *School of Mathematical Sciences, Dublin City University, address.
E-mail: eugene.oriordan@dcu.ie,*

The length of the following short abstract might vary from 1 to a few paragraphs.

Interior layers can appear in linear singularly perturbed parabolic problems when the data

KEY WORDS: time dependent, interior layers.

REFERENCES

1. R. K. Dunne and E. O’ Riordan, Interior layers arising in linear singularly perturbed differential equations with discontinuous coefficients, Proceedings of the Fourth International Conference on Finite Difference Methods: Theory and Applications, Lozenetz, Bulgaria, August 26–29, 2006 (I. Farago, P. Vabishchevich and L. Vulkov eds.), Rouse University, Bulgaria, 2007, 29–38. (extended version appeared as DCU preprint 2006, MS-06-09).
2. P. W. Hemker and G.I. Shishkin, Approximation of parabolic PDEs with a discontinuous initial condition, *East-West J. Numer. Math.*, **1**(4), 287–302, 1993.
3. E. O’ Riordan and G. I. Shishkin Singularly perturbed parabolic problems with non-smooth data, *Journal Computational and Applied Mathematics*, 2004, v. 166, n. 1, 233-245.