


2425-MA140 Engineering Calculus

Week 2, Lectures 3 The Squeeze Theorem

Dr Niall Madden

School of Mathematical and Statistical Sciences, University of Galway

Thursday, 26 September, 2024



This version of the slides are by Niall Madden, but are adapted from original notes by Dr Kirsten Pfeiffer.

Outline

1 News!

- Assignments, Tutorials and SUMS

2 Recall... Limits

3 Limits of rational functions

4 More limits

- Exercises

5 The Squeeze Theorem

- $\sin(\theta)/\theta$
- Other examples

For more, see Section 7.8.1 (Limit of a function of a real number) in *Modern Engineering Mathematics*:

https://search.library.nuigalway.ie/permalink/f/3b1kce/TN_cdi_askewsholts_vlebooks_9780273742517

Assignment 1

- ▶ **Assignment 1** has started! You can access it on Canvas... 2425-MA140... Assignments.
- ▶ Deadline: 5pm, Friday 4 Oct 2024. (Note: that's just the deadline, you can actually start before then!)
- ▶ The **Tutorial Sheet** is available at https://universityofgalway.instructure.com/files/2040359/download?download_frd=1

.....

Tutorials started **this** week. The schedule is on the Canvas “Course Information” page: <https://universityofgalway.instructure.com/courses/35693/pages/2425-ma140-information>

.....

Support is also available at **SUMS**...

Recall... Limits


Yesterday, we learned that

$$\epsilon \quad \lim_{x \rightarrow a} f(x) = L, \quad \delta$$

means that we can make $f(x)$ as close to L as we like, by taking x as close to a as needed.

Crucially, we are usually interested in finding the limit of $f(x)$ as $x \rightarrow a$, when a is not in the domain of f .

A typical example of this is when we evaluate a rational function:


$$\lim_{x \rightarrow a} \frac{p(x)}{q(x)}$$

where **both** $p(a) = 0$ and $q(a) = 0$. **Idea:** Since we care about the value of p and q **near** $x = a$, but not actually at $x = a$, it is safe to factor out and $(x - a)$ term from both.

Limits of rational functions

Example

Evaluate Consider

$$\lim_{x \rightarrow 1} \frac{x^2 + x - 2}{x^2 - x} = \lim_{x \rightarrow 1} \frac{p(x)}{q(x)}$$

$$p(x) = x^2 + x - 2$$

$$q(x) = x^2 - x$$

$$\text{Check } p(1) = 1^2 + 1 - 2 = 0 \quad q(1) = 1 - 1 = 0.$$

$$\text{So we factorise } p(x) = x^2 + x - 2 = (x+2)(x-1).$$

$$q(x) = x^2 - x = x(x-1).$$

$$\text{So } \frac{x^2 + x - 2}{x^2 - x} = \frac{(x+2)(x-1)}{x(x-1)} = \frac{x+2}{x} \quad \text{f.o. } x \neq 1$$

$$\lim_{x \rightarrow 1} \frac{p(x)}{q(x)} = \lim_{x \rightarrow 1} \frac{x+2}{x} = \frac{3}{1} = 3 \quad \checkmark$$

Limits of rational functions

In that last example, we found that

$$\lim_{x \rightarrow 1} \frac{x^2 + x - 2}{x^2 - x} = \lim_{x \rightarrow 1} \frac{x + 2}{x}$$

But these are different functions:

$x=1$ is not in the domain of $\frac{x^2 + x - 2}{x^2 - x}$ but is in the domain of $\frac{x+2}{x}$.

Limits of rational functions

Evaluate the limit

$$\lim_{x \rightarrow 2} \left(\frac{\frac{1}{2} - \frac{1}{x}}{x - 2} \right) = \lim_{x \rightarrow 2} \frac{p(x)}{q(x)} \quad \text{51}$$

where $p(x) = \frac{1}{2} - \frac{1}{x}$

$$q(x) = x - 2$$

Note $p(2) = \frac{1}{2} - \frac{1}{2} = 0$

$$q(2) = 2 - 2 = 0$$

Note, however, $\frac{1}{2} - \frac{1}{x} = \frac{x}{2x} - \frac{2}{2x} = \frac{x-2}{2x}$.

Then $\left(\frac{1}{2} - \frac{1}{x} \right) \cdot \frac{1}{x-2} = \left(\frac{x-2}{2x} \right) \left(\frac{1}{x-2} \right)$

If $x \neq 2$, this is $\frac{1}{2x}$.

$$\text{So } \lim_{x \rightarrow 2} \left(\frac{\frac{1}{2} - \frac{1}{x}}{x - 2} \right) = \lim_{x \rightarrow 2} \frac{1}{2x} = \frac{1}{4}.$$

More limits

Very often, we'll evaluate limits of the form:

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)}$$

where f and g are not polynomials. Some of the same ideas still apply.

Example

Evaluate

$$\lim_{x \rightarrow 0} \frac{\sqrt{1+x^2} - 1}{x^2}$$

Idea: simplify by completing the square in the numerator.

More limits

$$\lim_{x \rightarrow 0} \frac{\sqrt{1+x^2} - 1}{x^2} \quad (\text{Note if}$$

$$f(x) = \frac{\sqrt{1+x^2} - 1}{x^2}, \text{ then } f(0) = \frac{0}{0} \text{ . Not defined!}$$

However

$$\frac{\sqrt{1+x^2} - 1}{x^2} = \frac{(\sqrt{1+x^2} - 1)(\sqrt{1+x^2} + 1)}{x^2 (\sqrt{1+x^2} + 1)}.$$

$$= \frac{(\sqrt{1+x^2})^2 - 1}{x^2 (\sqrt{1+x^2} + 1)} = \frac{1+x^2 - 1}{x^2 (\sqrt{1+x^2} + 1)}$$

$$= \frac{x^2}{x^2 (\sqrt{1+x^2} + 1)} = \frac{1}{\sqrt{1+x^2} + 1} \quad \text{for } x \neq 0.$$

$$\text{So } \lim_{x \rightarrow 0} \frac{\sqrt{1+x^2} - 1}{x^2} = \lim_{x \rightarrow 0} \frac{1}{\sqrt{1+x^2} + 1} = \frac{1}{2}$$

Exercise 2.4

(From 2023/2024 MA140 exam, Q1(a)) Evaluate the limit

$$\lim_{x \rightarrow 4} \frac{x - 4}{(\sqrt{x} - 2)(x + 9)}$$

The Squeeze Theorem

There are various approaches to evaluating limits. One significant one is...

The Squeeze Theorem (a.k.a. Sandwich Theorem)


Suppose that for functions f , g and h in a given interval I :

$$g(x) \leq f(x) \leq h(x) \quad \text{and} \quad \lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} h(x) = L.$$

Then

$$\lim_{x \rightarrow c} f(x) = L.$$

So, if $f(x)$ is between g & h
and g, h have some limit,
so too does f .



The Squeeze Theorem

Example

Suppose $f(x)$ is a function such that

$$1 - \frac{x^2}{4} \leq f(x) \leq 1 + \frac{x^2}{2}, \quad \forall x \neq 0$$

Find $\lim_{x \rightarrow 0} f(x)$.

Since $\lim_{x \rightarrow 0} g(x) = \lim_{x \rightarrow 0} \left(1 - \frac{x^2}{4}\right) = 1$.

Similarly $\lim_{x \rightarrow 0} h(x) = 1$. So

$$\lim_{x \rightarrow 0} f(x) = 1.$$


The Squeeze Theorem

$$\sin(\theta)/\theta$$

We use the Squeeze Theorem to explain **an important limit**:

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$$

Before we show this is true, let's convince ourselves:



Note $\sin(0) = 0$.

So $\frac{\sin(\theta)}{\theta} = \frac{0}{0}$ at $\theta = 0$.

Finish here

Thursday

Before we use the Squeeze Theorem, we need a few facts about trigonometric functions.

- ▶ **In this module, we only ever use radians** (never, ever degrees).
- ▶ Given the triangle drawn below, $\sin \theta = \frac{b}{h}$, $\cos \theta = \frac{a}{h}$,
$$\tan \theta = \frac{b}{a} = \frac{\sin \theta}{\cos \theta}$$
- ▶ Area of a sector of a circle is $\frac{1}{2}r^2\theta$ where r is the radius of the circle, and θ is the angle subtended by the sector.

Various other facts are summarised in the State Examination Commission's Tables:

Triantánacht


Trigonometry

$$\tan A = \frac{\sin A}{\cos A}$$

$$\cot A = \frac{\cos A}{\sin A}$$

$$\sec A = \frac{1}{\cos A}$$

$$\operatorname{cosec} A = \frac{1}{\sin A}$$



$$\cos^2 A + \sin^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\cos(-A) = \cos A$$

$$\sin(-A) = -\sin A$$

$$\tan(-A) = -\tan A$$

Nóta: Bíonn $\tan A$ agus $\sec A$ gan sainiú nuair $\cos A = 0$.

Bíonn $\cot A$ agus $\operatorname{cosec} A$ gan sainiú nuair $\sin A = 0$.

Note: $\tan A$ and $\sec A$ are not defined when $\cos A = 0$.

$\cot A$ and $\operatorname{cosec} A$ are not defined when $\sin A = 0$.

| A (céimeanna) | 0° | 90° | 180° | 270° | 30° | 45° | 60° | A (degrees) |
|-----------------|-----------|-----------------|-------------|------------------|----------------------|----------------------|----------------------|---------------|
| A (raidiaín) | 0 | $\frac{\pi}{2}$ | π | $\frac{3\pi}{2}$ | $\frac{\pi}{6}$ | $\frac{\pi}{4}$ | $\frac{\pi}{3}$ | A (radians) |
| $\cos A$ | 1 | 0 | -1 | 0 | $\frac{\sqrt{3}}{2}$ | $\frac{1}{\sqrt{2}}$ | $\frac{1}{2}$ | $\cos A$ |
| $\sin A$ | 0 | 1 | 0 | -1 | $\frac{1}{2}$ | $\frac{1}{\sqrt{2}}$ | $\frac{\sqrt{3}}{2}$ | $\sin A$ |
| $\tan A$ | 0 | - | 0 | - | $\frac{1}{\sqrt{3}}$ | 1 | $\sqrt{3}$ | $\tan A$ |

$$1 \text{ rad.} \approx 57.296^\circ$$

$$1^\circ \approx 0.01745 \text{ rad.}$$

Foirmlí uillinneacha comhshuite

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

Compound angle formulae

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

Foirmlí uillinneacha dúbailte

$$\cos 2A = \cos^2 A - \sin^2 A$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos^2 A = \frac{1}{2}(1 + \cos 2A)$$

$$\sin^2 A = \frac{1}{2}(1 - \cos 2A)$$

Double angle formulae

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

$$\cos 2A = \frac{1 - \tan^2 A}{1 + \tan^2 A}$$

$$\sin 2A = \frac{2 \tan A}{1 + \tan^2 A}$$

Iolraigh a thiontú ina suimeanna agus ina ndifríochtaí**Products to sums and differences**

$$2 \cos A \cos B = \cos(A + B) + \cos(A - B)$$

$$2 \sin A \cos B = \sin(A + B) + \sin(A - B)$$

$$2 \sin A \sin B = \cos(A - B) - \cos(A + B)$$

$$2 \cos A \sin B = \sin(A + B) - \sin(A - B)$$

Suimeanna agus difríochtaí a thiontú ina n-iolraigh**Sums and differences to products**

$$\cos A + \cos B = 2 \cos \frac{A+B}{2} \cos \frac{A-B}{2}$$

$$\cos A - \cos B = -2 \sin \frac{A+B}{2} \sin \frac{A-B}{2}$$


$$\sin A + \sin B = 2 \sin \frac{A+B}{2} \cos \frac{A-B}{2}$$

$$\sin A - \sin B = 2 \cos \frac{A+B}{2} \sin \frac{A-B}{2}$$

The Squeeze Theorem


$$\sin(\theta)/\theta$$

Here are plots of $\sin \theta$ (**red**) and $\cos \theta$ (**blue**).



$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$$

Now let's reason more carefully:



Example

Evaluate $\lim_{x \rightarrow 0} \frac{\tan 3x}{\sin 2x}$

Example

Evaluate $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2}$