


Lecture 2: Functions

Dr Niall Madden

School of Mathematical and Statistical Sciences, University of Galway

18 September, 2024



This version of the slides are by Niall Madden, but are adapted from original notes by Dr Kirsten Pfeiffer.

Outline

1 Functions

- Notation and terminology
- 4 Ways to Represent a Function

2 Graphical Representation

- Domain Convention

3 A Catalog of Functions

- Linear functions

4 Polynomials

- Sketching polynomials

For more, see Chapter 2 of *Modern Engineering Mathematics*:

https://search.library.nuigalway.ie/permalink/f/3b1kce/TN_cdi_askewsholts_vlebooks_9780273742517

MA313 Lecture 2: Functions

Start of ...

Section 1: Functions

*The single most important concept in MA140 is that of a **function**.
For more, see Chapter 2 of “Modern Engineering Calculus (James).*

We represent a function symbolically in two ways, either

"is mapped"
or "is sent to".

$f: x \rightarrow y$

"f is a function mapping x to y"

$y = f(x)$

Here x is in the set of X (or $x \in X$), and y is in the set of Y (or $y \in Y$).

$y = f(x)$ "y is f of x"

Later we'll also have $z = f(x, y)$.

More over $y = f(x)$ defines a point (x, y)
 $= (x, f(x))$

If f is a function from X to Y ...

- ▶ The set X is called the domain of the function.
- ▶ The set Y is called the codomain.
- ▶ When we write $y = f(x)$, we say “ x ” is the argument of the function. *ie = input.*
- ▶ When $y = f(x)$ for some $x \in X$, y is said to be the image of x under f . *image = output.*
- ▶ The set of all images $y = f(x), x \in X$, is called the range (or image set) of f . *Range \subseteq Codomain.*

f is a function from DOMAIN to
the CODOMAIN

$\uparrow X$


$\uparrow Y$

- ▶ While we could have functions between any pair of sets (e.g., a function from students in this class to their ID numbers), usually X and Y are *sets of numbers*.
- ▶ It is not necessary for all elements y of the codomain Y to be images under f .
- ▶ One element $y \in Y$ can serve as value $f(x)$ for several $x \in X$.

← uniqueness

A function can be represented in different ways:

1. **verbally** (by a description in *words*);
2. **numerically** (as a *table of values*);
3. **visually** (as a *graph*);
4. **algebraically** (by an explicit *formula*).



Often it is possible, and useful, to go from one way to another.


x	$f(x)$
-1	2
0	4
1	6
2	8
3	10.

eg $f(x) = 2x + 4$

Graphical Representation

Graph \rightarrow Table

A common way to *visualize* a function $f: X \rightarrow \mathbb{R}$ is its *graph* in the x, y -plane. In this example, $f(x) = x^2 - x + 4$.



$$f(-4) = (-4)^2 - (-4) + 4 \\ = 16 + 4 + 4 = 24$$

x	$f(x)$
-4	24 ✓
-2	10 ✓
0	4
2	6
4	16

Often, the domain of a function is not explicitly stated.
In such a case the following **Domain Convention** applies.

The **domain** of a function f is the set of all numbers x for which $f(x)$ makes sense and gives a *real-number output*.

Example

1. Find the domain D of $f(x) = \frac{1}{x^2 - x}$.

eg, if $x = 2$ $f(x) = \frac{1}{4-2} = \frac{1}{2}$

$x = 1$ $f(x) = \frac{1}{1-1} = \frac{1}{0}$ undefined.

$x = 0$ $f(x) = \frac{1}{0-0} = \frac{1}{0}$ undefined.

All other points are OK !!

Often, the domain of a function is not explicitly stated.
In such a case the following **Domain Convention** applies.

The **domain** of a function f is the set of all numbers x for which $f(x)$ makes sense and gives a *real-number* output.

Example

1. Find the domain D of $f(x) = \frac{1}{x^2 - x}$.

Note .

- $\{a, b\}$ means the set with two points a and b .
- $[a, b]$ means all point x with $a \leq x \leq b$
- $(a, b]$ means $a < x \leq b$
- (a, b) means $a < x < b$

\cup = "union"

Often, the domain of a function is not explicitly stated.
In such a case the following **Domain Convention** applies.

The **domain** of a function f is the set of all numbers x for which $f(x)$ makes sense and gives a *real-number* output.

Example

1. Find the domain D of $f(x) = \frac{1}{x^2 - x}$.

Answer: the domain is
 $(-\infty, 0) \cup (0, 1) \cup (1, \infty)$
 \uparrow union \nearrow
 $= \mathbb{R} / \{0, 1\}$

Example

Find the domain of the function $f(x) = \sqrt{x+2}$.

Since we want $f(x)$ to be real, we
can't have $f(x) = \sqrt{\text{something negative}}$

So the domain is when $x+2 \geq 0$

That is $x \geq -2$

Equivalent Domain is $[-2, \infty)$

Example

Identify the domain, codomain and range of $f_1(x) = 3x^2 + 1$

Domain is $\mathbb{R} = (-\infty, \infty)$

Codomain is also \mathbb{R} .

Note that, for all x , $x^2 \geq 0$

$$\text{So } 3x^2 \geq 0$$

$$\text{And } 3x^2 + 1 \geq 1$$

So range is $y \geq 1 \Leftrightarrow [1, \infty)$

Example

Identify the domain, codomain and range of $f_2(x) = \sqrt{(x+4)(3-x)}$ $= \mathbb{R}$.

Take a 60 second break
& think about this!

Idea ?? Need $(x+4)(3-x) \geq 0$.

So

- Both $x+4 \geq 0, 3-x \geq 0$
- Both $x+4 \leq 0$ and $3-x \leq 0$.

Domain

Exer: finish.

Example


Identify the domain, codomain and range of $f_2(x) = \sqrt{(x+4)(3-x)}$

[These notes were added after class]

The domain of this function is $[-4, 3]$.

Outside of that range the function is not real valued because $(x+4)(3-x)$ would be negative.

Here is a sketch:



The range is $[0, 7/2]$ but we'll learn in a few weeks how to prove that.

Example

Identify the domain, codomain and range of $f_3(x) = \frac{1}{x}$.
 $= \mathbb{R}$

Domain: All values for which $\frac{1}{x}$ makes sense. So any x except $x=0$.

Ans: $\mathbb{R}/\{0\}$ or $(-\infty, 0) \cup (0, +\infty)$.

Range: $\mathbb{R}/\{0\}$.

A Catalog of Functions

There are many *different types of functions* that can be used to *model relationships* between objects in the *real world*.


The most common types of functions (in MA140) are:

- ▶ Linear Functions, $3x - 2$
 - ▶ Polynomial Functions, $4x^4 - 5x^3 + 1$
 - ▶ Power Functions, 5^x
 - ▶ Rational Functions, $\frac{3x - 2}{4x + 1}$
 - ▶ Algebraic Functions, $\left\{ \begin{array}{l} \text{Rational Functions} \\ \text{Algebraic Functions} \end{array} \right.$
 - ▶ Trigonometric Functions, $\left\{ \begin{array}{l} \text{Algebraic Functions} \\ \text{Trigonometric Functions} \end{array} \right.$
 - ▶ Exponential Functions, $\left\{ \begin{array}{l} \text{Trigonometric Functions} \\ \text{Exponential Functions} \end{array} \right.$
 - ▶ Logarithms, $\left\{ \begin{array}{l} \text{Exponential Functions} \\ \text{Logarithms} \end{array} \right.$
- See tomorrow
...

Linear functions have formulae such as $f(x) = mx + c$, where m and c are some given numbers.

It is often represented graphically as a straight line of slope m through the point $(0, c)$.

$$f(x) = 2x + 1$$



A **polynomial function** (or just **polynomial**) is a function of the form

$$y = f(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x^1 + a_0, \quad x \in \mathbb{R},$$

where a_0, a_1, \dots, a_n are real numbers called the **coefficients** of the polynomial.

The number n is called the **degree** of the polynomial.

n is a finite integer.

Eg $n=0$: constant functions $f(x) = a_0$

$n=1$: Linear functions

$n=2$ Quadratic eg $x^2 + 2x - 3$.

Example: linear

$y = x$ is a **linear** polynomial with degree $n = 1$.


Example: quadratic

$y = x^2 - 4x + 3$ is a **quadratic** polynomial with degree $n = 2$.

Polynomials

Example

$y = x^3 - 4x^2 + x + 6$ is a **cubic** function with degree $n = 3$.



Fact: A polynomial function of grade n has **up to** $n - 1$ bends,

Examples:

Break Time

During the break, think and talk about what you might do to sketch the graph of

$$y = -x^3 + x^2 + 2x$$

- ▶ To sketch the graph, first find the **intercepts**:
 - ▶ The **y-intercepts** can be found by letting $x = 0$.
 - ▶ The **x-intercepts** are called the **roots** (or **zeros**).
To find the roots, set y equal to zero and solve for x .
- ▶ You don't have to use the same scale on the x - and on the y -axis.
- ▶ Do not use graph paper.


Example

Sketch the graph of

$$y = -x^3 + x^2 + 2x$$

How to sketch $y = -x^3 + x^2 + 2x$

Actual plot of $y = -x^3 + x^2 + 2x$



Exercises

Sketch the graphs of

(i) $y = 5x^2 - 7$

(ii) $y = x^2 - 4x + 3$

(iii) $y = x^3 - 6x^2 - 11x - 6$

