

MA385: Tutorial Sheet 2

These exercises are for Tutorial 2 (Week 4). You do not have to submit solutions to these questions. However, similar questions may appear on the assignment, class test, or final exam.

.....

In the following questions, take

$$f(x) = x - \cos(x) \tag{1}$$

It may be useful to know that, if $f(\tau) = 0$ then $\tau \approx 0.7390851332$.

- Q1. Show that there is a solution to $f(x) = 0$ in the interval $[0, 1]$.
- Q2. Is it possible to use Theorem 1.5.2 to determine that Newton's method, applied to f in (1) will converge for any choice of $x_0 \in [0, 1]$? Can it be used to determine that Newton's method will converge for *any* $x_0 \in \mathbb{R}$?
- Q3. Take $x_0 = 1$. Use Newton's method x_1 , x_2 , and x_3 as estimates for solutions to $x - \cos(x) = 0$. If possible, evaluate the corresponding values of $f(x_k)$, and $|\tau - x_k|$, to convince yourself that the method is converging. Use a computer program to do this, if you like.
- Q4. Let's suppose we don't know τ , only that it is located between $x = 0$ and $x = 1$. Taking $x_0 = 1$, give upper bounds for $|\tau - x_1|$, $|\tau - x_2|$, and $|\tau - x_3|$ using the *Newton Error Formula*. How does this compare with the corresponding bounds for the Bisection Method, and with the actual errors?