


MA378 Chapter 4: Finite Element Methods

§4.1 Boundary value problems

Dr Niall Madden (Niall.Madden@UniversityOfGalway.ie)

Start: 20 March 2026 (W10.2)



(The images don't have much to do with this lecture. But I like them. They are solutions to a certain PDE computed using a Python-based finite element system called [FEniCS](#).)

1.0 Today:

1 Introduction

- Initial V Boundary Value Problems
- Methods
- FEMs

2 Boundary value problems

3 Where the solution lives

4 The variational form

5 Exercises

1.1 Introduction

In this final section of MA378 we consider numerical schemes for particular ordinary differential equations called

Boundary Value Problems (BVPs).

BVPs are types of **differential equations**, meaning that

- ▶ They are equations (i.e., feature an “=” symbol) that we try to solve for some unknown.
- ▶ The equation features derivatives of the unknown.

BVPs V IVPs

The main differences BVPs and the initial value problems (IVPs) of MA385 are:

- ▶ The equation gives the solution to the DE **two** (boundary) points, not just one (initial) point.
- ▶ The second derivative of the solution **always** appears in the equation. The IVPs we studied usually only had first derivatives.
- ▶ We usually think of the independent variable, x , as representing **space** rather than time.

For some BVPs it is possible to write down the exact solution (an example is given below). Usually, however, this is not easy, or, indeed, possible. So numerical methods are needed to give *approximate* solutions. Two of the most popular numerical methods are

- ▶ **Finite Difference Methods** (FDMs), where one approximates the differential equation, based on Taylor series expansions.
- ▶ **Finite Element Methods** (FEMs), where the solution space is approximated, using ideas related to interpolation.

We'll study **FEMs** because they are in keeping with the philosophy of this course, **and because they are the most important family of algorithms in modern computational mathematics.**

More details are in Chapter 14 of the Süli and Mayers.


The practical applications of FEMs are too many to mention.

The figure opposite, produced by Sean Tobin, your tutor, relates to modelling fluid flow in a human aorta.

It is joint work with Niamh Hynes (google her!) from Biomedical Engineering/Surgery.

We coded the simulation in a Python-based system called **Firedrake**.

If you are interested in learning more (maths and programming), talk to be about a Summer Research Project.



1.2 Boundary value problems

To formulate a model BVP we first define a differential operator

$$L(u) := -u''(x) + r(x)u(x). \quad (1)$$

The practice of using L for such an operator comes from the fact that it is **linear**, meaning that $L(u + v) = L(u) + L(v)$.

This operator can be applied to any function with a defined second derivative.

1.2 Boundary value problems

The general form of a BVP is:

Find a function u , defined on the interval $[a, b]$, such that

$$L(u) = f(x) \quad \text{for } a < x < b, \quad \text{with } u(a) = 0, u(b) = 0. \quad (2)$$

We make the assumptions that r and f are continuous and have as many continuous derivatives as we would like. **Furthermore we always have that $r(x) > 0$ for all $x \in [a, b]$.**

Some BVPs are easy to solve by hand. That is, we can write down a solution in terms of elementary functions. That is certainly the case if the functions r and f are constant.

1.2 Boundary value problems

Example 1.1

Verify that, for any constants c_0 and c_1 ,
 $u(x) = c_0 e^{2x} + c_1 e^{-2x} + x/4$ is a solution to

$$-u''(x) + 4u(x) = x \quad \text{for } 0 \leq x \leq 3, \quad (3)$$

(If we have two boundary conditions, we can uniquely determine c_0 and c_1 . See Exercise 1.2).

$$u(x) = c_0 e^{2x} + c_1 e^{-2x} + \frac{1}{4}x$$

$$u'(x) = 2c_0 e^{2x} - 2c_1 e^{-2x} + \frac{1}{4}$$

$$u''(x) = 4c_0 e^{2x} + 4c_1 e^{-2x}$$

$$\begin{aligned} \Rightarrow -u''(x) + 4u(x) &= -(\cancel{4c_0 e^{2x}} + \cancel{4c_1 e^{-2x}}) \\ &\quad + \cancel{4c_0 e^{2x}} + \cancel{4c_1 e^{-2x}} + x = x \end{aligned}$$

1.2 Boundary value problems

Unlike the example above, most boundary value problems are difficult or impossible to solve analytically and so, in general, we need approximations. However, it can often be possible to obtain **qualitative** information about the solutions.

One crucial tool for this is called a maximum principle.

1.2 Boundary value problems

Lemma 1.2 (Maximum Principle)

Let L be the differential operator $L(u) := -u''(x) + r(x)u(x)$, where $r(x) > 0$ for all $x \in [a, b]$, and u is a function such that $L(u) \geq 0$ on (a, b) . If $u(a) \geq 0$, $u(b) \geq 0$. Then $u \geq 0$ for all $x \in [a, b]$.

Proof: Let's suppose there is a point $x^* \in [a, b]$

such that $u(x^*) = \min_{a \leq x \leq b} u(x)$ & $u(x^*) < 0$.

Clearly $x^* \neq a$, $x^* \neq b$, so $x^* \in (a, b)$.

That is x^* is a local minimum of u .

So $u'(x^*) = 0$ and, moreover $u''(x^*) \geq 0$.

But then

$$\underbrace{-u''(x^*)}_{\leq 0} + \underbrace{r(x^*)}_{> 0} \underbrace{u(x^*)}_{< 0} < 0, \text{ which is not possible}$$

1.2 Boundary value problems

This lemma is as useful as it is simple (see also Exercise 1.4).

Lemma 1.3

There is at most one solution to the BVP

$$L(u) = f(x) \quad \text{for } a < x < b, \quad \text{with } u(a) = 0, u(b) = 0.$$

Suppose there are two solutions to this, u & v .

So $L(u) = f(x)$ & $L(v) = f(x)$.

Let $w = u - v$ then $L(u) - L(v) = f(x) - f(x)$

giving $L(w) = 0$. Also $w(a) = w(b) = 0$.

So we can apply lemma 1.2 to show $w(x) \geq 0$.
However, we also get that

$L(-w) = L(v) - L(u) = 0$. So $-w(x) \geq 0$ too. So $w(x) = 0$.

1.3 Where the solution lives

The boundary value problem

$$-u''(x) + r(x)u(x) = f(x) \quad a < x < b,$$

and

$$u(a) = 0, u(b) = 0,$$

equates the terms in u and u'' with a function f that is continuous on (a, b) , so we must have that u , u' and u'' are all continuous on the interval (a, b) . That is, u belongs to the space of functions $C^2(a, b)$.

So we can state the problem more precisely: *find $u \in C^2(a, b)$ such that*

$$-u''(x) + r(x)u(x) = f(x) \quad \text{for all } x \text{ in } (a, b),$$

$$\text{and } u(a) = u(b) = 0.$$

1.4 The variational form

Before we give a numerical method for computing an approximate solution to a boundary value problem, we will rewrite it as an integral equation.

Definition 1.4 ($H_0^1(a, b)$)

Define (again) the inner product: $(u, v) := \int_a^b u(x)v(x)dx$. With this we can define the norm: $\|u\|_2 := (u, u)^{1/2}$. Then $H_0^1(a, b)$ is defined to be the space of (absolutely) continuous functions on $[a, b]$ such that if $w \in H_0^1(a, b)$ then

$$w(a) = w(b) = 0 \quad \text{and} \quad \|w'\|_2 < \infty.$$

(We met similar space before in the section on cubic splines.)

ie $\|u\|_2 = \sqrt{\int_a^b (u(x))^2 dx}$

1.4 The variational form

Consider the Boundary Value Problem: *find* $u \in C^2(a, b)$ *such that*

$$-u''(x) + r(x)u(x) = f(x) \quad \text{on } (a, b); \text{ and } u(a) = u(b) = 0. \quad (4)$$

Suppose that we have a solution u to this equation. Then the **variational form** of (4) is derived as on the following slide.

1.4 The variational form

$$-u''(x) + r(x)u(x) = f(x) \text{ on } (a, b); \quad \text{and } u(a) = u(b) = 0.$$

Suppose u solves this. So $-u''(x) + r(x)u(x) = f(x)$

Then, for any function v :

$$(-u''(x) + r(x)u(x))v(x) = f(x)v(x) \quad \text{for all } x \in (a, b)$$

So we can integrate

$$\int_a^b -u''(x)v(x) + r(x)u(x)v(x) dx = \int_a^b f(x)v(x) dx$$

Using integration by parts:

$$-\int_a^b u''(x)v(x) dx = -u'(x)v(x) \Big|_a^b + \int_a^b u'(x)v'(x) dx.$$

This holds for all v , including those with $v(a) = v(b) = 0$

1.4 The variational form

$$-u''(x) + r(x)u(x) = f(x) \text{ on } (a, b); \quad \text{and } u(a) = u(b) = 0.$$

In summary, we have that

$$\begin{aligned} \int_a^b u'(x) v'(x) dx + \int_a^b r(x) u(x) v(x) dx \\ = \int_a^b f(x) v(x) dx \end{aligned}$$

for all v with $v(a) = v(b) = 0$.

ie $(u', v') + (ru, v) = (f, v).$

□

Finished here Friday (W10.2)

1.4 The variational form

Definition 1.5 (Variational formulation)

The variational/weak formulation of (4) is: *Find $u \in H_0^1(a, b)$ such that*

$$\mathcal{A}(u, v) = (f, v) \quad \text{for all } v \in H_0^1(a, b). \quad (5)$$

where $\mathcal{A}(\cdot, \cdot)$ is the symmetric bilinear functional

$$\mathcal{A}(u, v) := (u', v') + (ru, v).$$

1.4 The variational form

This $\mathcal{A}$ has several important properties. One of these is that, for any function w ,

$$\mathcal{A}(w, w) \geq 0,$$

and

$$\mathcal{A}(w, w) = 0 \Leftrightarrow w = 0.$$

1.4 The variational form

The following lemma is an immediate consequence of these properties.

Lemma 1.6

The variational problem (5) has at most one solution.

1.5 Exercises

Exercise 1.1

In class we considered the differential operator

$$L(u) := -u''(x) + r(x)u(x).$$

where $r(x) > 0$ for all x . Suppose, instead have the more general operator

$$L_q(u) := -u''(x) + q(x)u'(x) + r(x)u(x),$$

where again r is a positive function. Does this L_q also satisfy a maximum principle? If so, provide a proof. If not, give a counter example.

Exercise 1.2

Verify that $u(x) = \frac{x}{4} + \frac{3e^6(e^{-2x} - e^{2x})}{4(e^{12} - 1)}$ is the exact solution to the differential equation $-u''(x) + 4u(x) = x$ for $x \in (0, 3)$, with the boundary conditions $u(0) = 0$, $u(3) = 0$,

1.5 Exercises

Exercise 1.3

In this section of the course, we'll always assume homogeneous boundary conditions. That is, that $u(x) = 0$ at the boundaries. Suppose the problem we wish to solve is

$$-u''(x) + r(x)u(x) = f(x) \quad u(0) = \alpha, u(1) = \beta.$$

Show how to find a problem which has the same left-hand side as this one, homogeneous boundary conditions, and with a solution that differs from this one only by a known linear function.

Exercise 1.4

Suppose that u solves $-u''(x) + r(x)u(x) = f(x)$ on $(0, 1)$, and $u(0) = u(1) = 0$. Let ϱ be such $r(x) \geq \varrho > 0$, and define

$$C = \max_{0 \leq x \leq 1} |f(x)|/\varrho.$$

Prove that $u(x) \leq C$. (Hint: Consider $L(C - u)$).

1.5 Exercises

Exercise 1.5

Consider the differential equation:

$$-u''(x) = \exp(x + 1), \text{ on } (0, 2), \text{ and } u(0) = u(2) = 0.$$

- (i) State the variational formulation of this differential equation.
- (ii) Show that the solution to the variational problem is unique.