

MA385 Part 1: Solving nonlinear equations

1.3: The Secant Method

Dr Niall Madden

September 2026

0. Outline

- 1 Motivation
- 2 Order of Convergence
- 3 Side quest: The Mean Value Theorem
- 4 Analysis of the Secant Method
- 5 Exercises

For more details, see Section 1.5 (The secant method) of [Süli and Mayers, *An Introduction to Numerical Analysis*](#)



1. Motivation

We'll start with considering, heuristically, how we could improve upon bisection:

1. Motivation

Idea:

- ▶ Choose two points, x_0 and x_1 .
- ▶ Take x_2 to be the zero of the line joining $(x_0, f(x_0))$ to $(x_1, f(x_1))$.
- ▶ Take x_3 to be the zero of the line joining $(x_1, f(x_1))$ to $(x_2, f(x_2))$.
- ▶ Etc.



1. Motivation

The Secant Method

Choose x_0 and x_1 so that there is a solution in $[x_0, x_1]$. Then define

$$x_{k+1} = x_k - f(x_k) \frac{x_k - x_{k-1}}{f(x_k) - f(x_{k-1})}. \quad (1)$$

1. Motivation

Example 1.3.1

Use the Secant Method to solve $x^2 - 2 = 0$ in $[0, 2]$. Results are shown below. We see that, not only does the method appear to converge to the true solution, it seem to do so *much* more efficiently than Bisection. We'll return to why this is later.

	Secant		Bisection	
k	x_k	$ x_k - \tau $	x_k	$ x_k - \tau $
0	0.000000	1.41	0.000000	1.41
1	2.000000	5.86e-01	2.000000	5.86e-01
2	1.000000	4.14e-01	1.000000	4.14e-01
3	1.333333	8.09e-02	1.500000	8.58e-02
4	1.428571	1.44e-02	1.250000	1.64e-01
5	1.413793	4.20e-04	1.375000	3.92e-02
6	1.414211	2.12e-06	1.437500	2.33e-02
7	1.414214	2.16e-10	1.406250	7.06e-02

2. Order of Convergence

To compare different methods, we need the following concept.

Definition 1.3.1 (Linear Convergence)

Suppose that $\tau = \lim_{k \rightarrow \infty} x_k$. We say that the sequence $\{x_k\}_{k=0}^{\infty}$ converges to τ **at least linearly** if there is a sequence of positive numbers $\{\varepsilon_k\}_{k=0}^{\infty}$, and $\mu \in (0, 1)$, such that

$$\lim_{k \rightarrow \infty} \varepsilon_k = 0, \quad (2a)$$

and

$$|\tau - x_k| \leq \varepsilon_k \quad \text{for } k = 0, 1, 2, \dots \quad (2b)$$

and

$$\lim_{k \rightarrow \infty} \frac{\varepsilon_{k+1}}{\varepsilon_k} = \mu. \quad (2c)$$

2. Order of Convergence

2. Order of Convergence

For Example 6, the bisection method converges at least linearly. As we have seen, the Secant Method converge more quickly than bisection. Now we'll give a more precise description of what “*more quickly*” means.

Definition 1.3.2 (Order of Convergence)

Let $\tau = \lim_{k \rightarrow \infty} x_k$. Suppose there exists $\mu > 0$ and a sequence of positive numbers $\{\varepsilon_k\}_{k=0}^{\infty}$ such that (2a) and (2b) both hold. Then we say that the sequence $\{x_k\}_{k=0}^{\infty}$ converges with at least order q if

$$\lim_{k \rightarrow \infty} \frac{\varepsilon_{k+1}}{(\varepsilon_k)^q} = \mu.$$

2. Order of Convergence

$$\lim_{k \rightarrow \infty} \frac{\varepsilon_{k+1}}{(\varepsilon_k)^q} = \mu.$$

Two particular values of q are important to us:

- (i) If $q = 1$, and we have that $0 < \mu < 1$, then the rate is **linear**.
- (ii) If $q = 2$, the rate is **quadratic** for any $\mu > 0$.

3. Side quest: The Mean Value Theorem

The **Mean Value Theorem** (aka “MVT”) is simple, and incredibly useful. The modern version is due to Cauchy (1823), but there is a version called Rolle’s Theorem which applies to polynomials only from 1691, and one for the sin function to Parameshvara (1380–1460) of Kerala, SW India.

Theorem 1.3.1 (Mean Value Theorem (MVT))

If $f : [a, b] \rightarrow R$ is continuous and differentiable, then there is a point $c \in (a, b)$, such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}.$$

Informally: *there is a point between a and b where the tangent to f is parallel to the line joining $(a, f(a))$ at $(b, f(b))$.*

3. Side quest: The Mean Value Theorem

A visual proof:

3. Side quest: The Mean Value Theorem

From a numerical analysis point of view, the MVT tells us “how different $f(b)$ might be from $f(a)$ ”:

$$f(b) = f(a) + f'(c)(b - a)$$

gives that

$$|f(b) - f(a)| \leq \max_{a \leq x \leq b} |f'(x)|(b - a).$$

That is, in a computation, if we replace $f(a)$ with $f(b)$, the error we introduce depends on $b - a$ and also the magnitude of f' .

4. Analysis of the Secant Method

Theorem 1.3.2

Suppose that f and f' are real-valued functions, continuous and defined in an interval $I = [\tau - h, \tau + h]$ for some $h > 0$. If $f(\tau) = 0$ and $f'(\tau) \neq 0$, then the sequence (1) converges at least linearly to τ .

4. Analysis of the Secant Method

- ▶ We wish to show that $|\tau - x_{k+1}| < |\tau - x_k|$.
- ▶ From the MVT, there is a point $w_k \in [x_{k-1}, x_k]$ such that

$$\frac{f(x_k) - f(x_{k-1})}{x_k - x_{k-1}} = f'(w_k). \quad (3)$$

- ▶ Also by the MVT, there is a point $z_k \in [x_k, \tau]$ such that

$$\frac{f(x_k) - f(\tau)}{x_k - \tau} = \frac{f(x_k)}{x_k - \tau} = f'(z_k). \quad (4)$$

Therefore $f(x_k) = (x_k - \tau)f'(z_k)$.

4. Analysis of the Secant Method

► Using (3) and (4), we can show that

$$\tau - x_{k+1} = (\tau - x_k) \left(1 - f'(z_k)/f'(w_k) \right).$$

Therefore

$$\frac{|\tau - x_{k+1}|}{|\tau - x_k|} = \left| 1 - \frac{f'(z_k)}{f'(w_k)} \right|.$$

4. Analysis of the Secant Method

- Suppose that $f'(\tau) > 0$. (If $f'(\tau) < 0$ just tweak the arguments accordingly). Saying that f' is *continuous in the region* $[\tau - h, \tau + h]$ means that, for any $\varepsilon > 0$ there is a $\delta > 0$ such that

$$|f'(x) - f'(\tau)| < \varepsilon \text{ for any } x \in [\tau - \delta, \tau + \delta].$$

Take $\varepsilon = f'(\tau)/4$. Then $|f'(x) - f'(\tau)| < f'(\tau)/4$. Thus

$$\frac{3}{4}f'(\tau) \leq f'(x) \leq \frac{5}{4}f'(\tau) \quad \text{for any } x \in [\tau - \delta, \tau + \delta].$$

Then, so long as w_k and z_k are both in $[\tau - \delta, \tau + \delta]$

$$\frac{f'(z_k)}{f'(w_k)} \leq \frac{5}{3}.$$

4. Analysis of the Secant Method

Given enough time and effort we *could* show that the Secant Method converges faster than linearly. In particular, that the order of convergence is

$$q = (1 + \sqrt{5})/2 \approx 1.618.$$

This number arises as the only positive root of $q^2 - q - 1$. It is called the **Golden Mean**, and arises in many areas of Mathematics, including finding an explicit expression for the Fibonacci Sequence:

$$f_0 = 1,$$

$$f_1 = 1,$$

$$f_{k+1} = f_k + f_{k-1} \text{ for } k = 2, 3, \dots$$

That gives, $f_0 = 1$, $f_1 = 1$, $f_2 = 2$, $f_3 = 3$, $f_4 = 5$, $f_5 = 8$, $f_6 = 13$,
...

4. Analysis of the Secant Method

The connection here is that it turns out that $\varepsilon_{k+1} \leq C\varepsilon_k\varepsilon_{k-1}$. Repeatedly using this we get:

- ▶ Let $r = |x_1 - x_0|$ so that $\varepsilon_0 \leq r$ and $\varepsilon_1 \leq r$,
- ▶ Then $\varepsilon_2 \leq C\varepsilon_1\varepsilon_0 \leq Cr^2$
- ▶ Then $\varepsilon_3 \leq C\varepsilon_2\varepsilon_1 \leq C(Cr^2)r = C^2r^3$.
- ▶ Then $\varepsilon_4 \leq C\varepsilon_3\varepsilon_2 \leq C(C^2r^3)(Cr^2) = C^4r^5$.
- ▶ Then $\varepsilon_5 \leq C\varepsilon_4\varepsilon_3 \leq C(C^4r^5)(C^2r^3) = C^7r^8$.
- ▶ And in general, $\varepsilon_k = C^{f_k-1}r^{f_k}$.

5. Exercises

Exercise 1.3.1

Suppose we define the Secant Method as follows.

Choose any two points x_0 and x_1 .

For $k = 1, 2, \dots$, set x_{k+1} to be the point where the line through $(x_{k-1}, f(x_{k-1}))$ and $(x_k, f(x_k))$ that intersects the x -axis.

Show how to derive the formula for the secant method.

5. Exercises

Exercise 1.3.2

Another interpretation of the Secant Method, is that it computes x_{k+1} as a weighted average:

$$x_{k+1} = (1 - \sigma_k)x_k + \sigma_k x_{k-1},$$

where σ_k chosen to obtain fast convergence to the true solution. (This is something called a “relaxation method”).

Derive a formula for σ_k in terms of $f(x_{k-1})$ and $f(x_k)$. Can you give a physical interpretation to σ_k and $1 - \sigma_k$ that might justify the claim that the method should be more efficient than bisection?

5. Exercises

Exercise 1.3.3

- (i) Is it possible to construct a problem for which the bisection method will work, but the secant method will fail? If so, give an example.
- (ii) Is it possible to construct a problem for which the secant method will work, but bisection will fail? If so, give an example.

5. Exercises

Exercise 1.3.4

In the proof of Theorem 1.3.2, we exploited the “ $\varepsilon - \delta$ ” definition of continuity to assume we could choose $\varepsilon \leq f'(\tau)/4$. Here the choice of 4 as the divisor is somewhat arbitrary. What other choices would work?