

2627-MA140 Engineering Calculus

Week 04, Lecture 2 Differentiation Rules

Dr Niall Madden

University of Galway

Wednesday, 30 September, 2026

Calculus	Calculus																																	
Diorthaigh	Derivatives																																	
<table><tr><th>$f(x)$</th><th>$f'(x)$</th></tr><tr><td>x^n</td><td>nx^{n-1}</td></tr><tr><td>$\ln x$</td><td>$\frac{1}{x}$</td></tr><tr><td>e^x</td><td>e^x</td></tr><tr><td>e^{ax}</td><td>ae^{ax}</td></tr><tr><td>a^x</td><td>$a^x \ln a$</td></tr><tr><td>$\cos x$</td><td>$-\sin x$</td></tr><tr><td>$\sin x$</td><td>$\cos x$</td></tr><tr><td>$\tan x$</td><td>$\sec^2 x$</td></tr><tr><td>$\cos^{-1} \frac{x}{a}$</td><td>$-\frac{1}{\sqrt{a^2 - x^2}}$</td></tr><tr><td>$\sin^{-1} \frac{x}{a}$</td><td>$\frac{1}{\sqrt{a^2 - x^2}}$</td></tr><tr><td>$\tan^{-1} \frac{x}{a}$</td><td>$\frac{a}{a^2 + x^2}$</td></tr></table>	$f(x)$	$f'(x)$	x^n	nx^{n-1}	$\ln x$	$\frac{1}{x}$	e^x	e^x	e^{ax}	ae^{ax}	a^x	$a^x \ln a$	$\cos x$	$-\sin x$	$\sin x$	$\cos x$	$\tan x$	$\sec^2 x$	$\cos^{-1} \frac{x}{a}$	$-\frac{1}{\sqrt{a^2 - x^2}}$	$\sin^{-1} \frac{x}{a}$	$\frac{1}{\sqrt{a^2 - x^2}}$	$\tan^{-1} \frac{x}{a}$	$\frac{a}{a^2 + x^2}$	<table><tr><td>Riall an toraidh</td><td>$y = uv$ $\Rightarrow \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$</td><td>Product rule</td></tr><tr><td>Riall an lin</td><td>$y = \frac{u}{v}$ $\Rightarrow \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$</td><td>Quotient rule</td></tr><tr><td>Cuingrial</td><td>$f(x) = u(v(x))$ $\Rightarrow f'(x) = \frac{du}{dv} \frac{dv}{dx}$</td><td>Chain rule</td></tr></table>	Riall an toraidh	$y = uv$ $\Rightarrow \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$	Product rule	Riall an lin	$y = \frac{u}{v}$ $\Rightarrow \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$	Quotient rule	Cuingrial	$f(x) = u(v(x))$ $\Rightarrow f'(x) = \frac{du}{dv} \frac{dv}{dx}$	Chain rule
$f(x)$	$f'(x)$																																	
x^n	nx^{n-1}																																	
$\ln x$	$\frac{1}{x}$																																	
e^x	e^x																																	
e^{ax}	ae^{ax}																																	
a^x	$a^x \ln a$																																	
$\cos x$	$-\sin x$																																	
$\sin x$	$\cos x$																																	
$\tan x$	$\sec^2 x$																																	
$\cos^{-1} \frac{x}{a}$	$-\frac{1}{\sqrt{a^2 - x^2}}$																																	
$\sin^{-1} \frac{x}{a}$	$\frac{1}{\sqrt{a^2 - x^2}}$																																	
$\tan^{-1} \frac{x}{a}$	$\frac{a}{a^2 + x^2}$																																	
Riall an toraidh	$y = uv$ $\Rightarrow \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$	Product rule																																
Riall an lin	$y = \frac{u}{v}$ $\Rightarrow \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$	Quotient rule																																
Cuingrial	$f(x) = u(v(x))$ $\Rightarrow f'(x) = \frac{du}{dv} \frac{dv}{dx}$	Chain rule																																

Let's learn about...

- | | |
|------------------------------------|-------------------------------|
| 1 Recall: Derivative as a function | ■ 4. Sum and Difference Rules |
| 2 Differentiation by rule | 3 The Product Rule |
| ■ 1. The Constant Rule | 4 The Quotient Rule |
| ■ 2. The Power Rule | 5 Chain Rule |
| ■ 3. The constant multiple rule | 6 Page 16 of the "log tables" |
| | 7 Exercises |

See also Section 3.3 of **Calculus** by Strang & Herman:

<https://openstax.org/books/calculus-volume-1/pages/3-3-differentiation-rules>

Recall: Derivative as a function

Yesterday, we learned that...

Definition: derivative as a function

If $f(x)$ is a be a function, its derivative (function), denoted $f'(x)$, is given by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}.$$

wherever that limit exists. The process of finding f' is called **differentiation**.

We then used that definition to find the derivative of $f(x) = x^2$, $f(x) = \sqrt{x}$, and $f(x) = \sin(x)$.

Using this “limit” definition to find f' is called “*differentiation from first principles*”. However, we usually use some rules (which all derive from the above definition) in order to find f' .

The Constant Rule

If f is a constant function, i.e. $f(x) = c$ for all x , then $f'(x) = 0$.

Why:

We've already deduced that

- ▶ The derivative of $f(x) = x^2$ is $f'(x) = 2x$
- ▶ The derivative of $f(x) = x^{1/2}$ is $f'(x) = \frac{1}{2}x^{-1/2}$

These are particular examples of the **Power Rules**

The Power Rule

If n is any real number, then

$$\frac{d}{dx}(x^n) = nx^{n-1}.$$

Examples Calculate the derivatives of the following functions

1. $f(x) = x^6$

2. $f(x) = \sqrt[3]{x}$

The constant multiple rule

Let $f(x)$ be any differentiable function, and let k be constant, then

$$\frac{d}{dx}(kf(x)) = k \frac{d}{dx}(f(x)).$$

Example: Find the derivative of $f(x) = 5x^4$.

The Sum and Difference Rules

Let $u(x)$ and $v(x)$ be any differentiable functions. Then

$$\frac{d}{dx}(u(x) + v(x)) = \frac{d}{dx}(u(x)) + \frac{d}{dx}(v(x)).$$

Similarly,
$$\frac{d}{dx}(u(x) - v(x)) = \frac{d}{dx}(u(x)) - \frac{d}{dx}(v(x)).$$

Example: Find the derivative of $f(x) = 1 + x + x^2$.

Actually, the “**Difference Rule**”, which states that

$$\frac{d}{dx}(u(x) - v(x)) = \frac{d}{dx}(u(x)) - \frac{d}{dx}(v(x)).$$

can be combined by combining the **Sum Rule** and the **Constant Multiple Rule**.

Example

Suppose that $f(x) = -5x^3 + 3x^2 - 9x + 7$, then find:

- (a) The derivative of $f(x)$;
- (b) The slope of the tangent line at $x = 2$;
- (c) The equation of the tangent at $x = 2$.

(a) $f'(x) = -15x^2 + 6x - 9$

(b) The slope of the tangent line at $x = 2$ is $f'(2)$:

$$f'(2) = -15(2)^2 + 6(2) - 9 = -15(4) + 12 - 9 = -60 + 12 - 9 = -57.$$

- (c) The equation of the line with slope M and passing through a point (x_1, y_1) is

$$y - y_1 = M(x - x_1)$$

The y coordinate at $x = 2$ is

$$\begin{aligned} f(2) &= -5(2)^3 + 3(2)^2 - 9(2) + 7 \\ &= -5(8) + 3(4) - 18 + 7 \\ &= -40 + 12 - 18 + 7 \\ &= -39. \end{aligned}$$

So the tangent line passes through the point $(2, -39)$ and the slope of the line is -57 .

Therefore, the equation of this line is $y + 39 = -57(x - 2)$

Ans: The equation of the tangent line is $x = 2$ is $y = 75 - 57x$.

The Product Rule

We now consider some rules which are a little more complicated and, perhaps, less obvious.

The first concerns the derivative of the **product** of two functions.

The Product Rule

Let $u(x)$ and $v(x)$ be any differentiable functions. Then

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}.$$

Let's check that this gives the current answer when evaluating the derivative of $f(x) = x^3$ when we set $u(x) = x^2$ and $v(x) = x$:

The Product Rule

The Product Rule

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}.$$

Example Use the **product rule** to find the derivative of $f(x) = x^3(x^2 + 1)$.

The Product Rule

Example: use the product rule to show that, if $f(x) = x \sin(x)$, then $f'(x) = x \cos(x) + \sin(x)$.

The Quotient Rule

The Quotient Rule

If u and v are differentiable at x and if $v(x) \neq 0$, then $f(x) = \frac{u(x)}{v(x)}$ is differentiable at x , and

$$\frac{d}{dx} \left(\frac{u}{v} \right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}.$$

Example: Use this rule to find the derivative of $f(x) = \frac{x+1}{x^2}$

The Quotient Rule

Example

We know that

- ▶ $\tan(x) = \frac{\sin(x)}{\cos(x)},$
- ▶ $\sin^2(x) + \cos^2(x) = 1$
- ▶ $\sin'(x) = \cos(x)$ and $\cos'(x) = -\sin(x).$

Use these facts, and the Quotient Rule to show that

$$\frac{d}{dx}(\tan(x)) = \left(\frac{1}{\cos(x)}\right)^2.$$

The Quotient Rule

Chain Rule

Of all the differentiation rules, the **chain rule** is the most important: most other rules are actually just special cases of it. It applies to a “function of a function”

The Chain Rule

If $u(x)$ and $v(x)$ are differentiable, and f is the composite function $f(x) = u(v(x))$, then

$$\frac{df}{dx} = \frac{du}{dv} \frac{dv}{dx}.$$

Chain Rule

The Chain Rule

If $f(x) = u(v(x))$, then

$$\frac{df}{dx} = \frac{du}{dv} \frac{dv}{dx}.$$

Example: What is the derivative of $f(x) = \cos(x^2)$?

Chain Rule

Example

Find $\frac{dy}{dx}$ if $y = (x^3 + 4x^4 + 7)^{12}$.

Example: Let $u(v) = v^{12}$ and $v(x) = x^3 + 4x^4 + 7$, then y is $y = u(v(x))$.

Note that

$$\frac{du}{dv} = 12v^{11} \quad \text{and} \quad \frac{dv}{dx} = 3x^2 + 16x^3.$$

By the Chain Rule we have

$$\frac{dy}{dx} = \frac{du}{dv} \frac{dv}{dx} = 12v^{11}(3x^2 + 16x^3),$$

and therefore

$$\frac{dy}{dx} = 12(x^3 + 4x^4 + 7)^{11}(3x^2 + 16x^3).$$

Chain Rule

Example (Not done in detail in class)

Find $\frac{dy}{dx}$ if $y = \frac{1}{(x^4 + 2x^2 + 8)^{40}}$.

We have $y = (x^4 + 2x^2 + 8)^{-40}$. We can write y as $y(x) = u(v(x))$ with

► $u(v) = v^{-40}$ and so $\frac{du}{dv} = -40v^{-41}$; and

► $v(x) = x^4 + 2x^2 + 8$, so $\frac{dv}{dx} = 4x^3 + 4x$.

Applying the Chain Rule: $\frac{dy}{dx} = \frac{du}{dv} \frac{dv}{dx}$, we get

$$\frac{dy}{dx} = -40v^{-41}(4x^3 + 4x) = \frac{-40(4x^3 + 4x)}{(x^4 + 2x^2 + 8)^{41}}$$

Calcalas

Calculus

Diorthaigh

Derivatives

$f(x)$	$f'(x)$
x^n	nx^{n-1}
$\ln x$	$\frac{1}{x}$
e^x	e^x
e^{ax}	ae^{ax}
a^x	$a^x \ln a$
$\cos x$	$-\sin x$
$\sin x$	$\cos x$
$\tan x$	$\sec^2 x$
$\cos^{-1} \frac{x}{a}$	$-\frac{1}{\sqrt{a^2 - x^2}}$
$\sin^{-1} \frac{x}{a}$	$\frac{1}{\sqrt{a^2 - x^2}}$
$\tan^{-1} \frac{x}{a}$	$\frac{a}{a^2 + x^2}$

Riail an toraidh	$y = uv$ $\Rightarrow \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$	Product rule
Riail an lín	$y = \frac{u}{v}$ $\Rightarrow \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$	Quotient rule
Cuingriail	$f(x) = u(v(x))$ $\Rightarrow f'(x) = \frac{du}{dv} \frac{dv}{dx}$	Chain rule

Exercises

Exercises 4.2.1 (Based on Q2(a), 2023/2024)

Find the derivative of $f(x) = \frac{\sin(x)}{\sqrt{x}}$.

Exercise 4.2.2 (Based on Q2(b), 2019/2020)

Find the derivative of $f(x) = \frac{x^2 + x - 2}{x^3 + 6}$.

Exercise 4.2.3

Find the derivative of

1. $f(x) = x^3 \cos(x^2)$
2. $f(x) = \tan^3(\sin^2(x^4))$