

Slow entropy of some skew products

Nicanor Carrasco-Vargas

Jagiellonian University

<https://arxiv.org/abs/2506.17932>

nicanor.vargas@uj.edu.pl

Beamer ∈ [nicanorcarrascovargas.github.io](https://github.com/nicanorcarrascovargas)

Please interrupt me.

Disclaimer

Please interrupt me.

In this talk all systems have discrete time \mathbb{Z} or \mathbb{N} (no flows or group actions)

Measurable dynamical system

Measure preserving transformation $T \curvearrowright (X, \mathcal{X}, \mu)$ on a probability space.

Topological dynamical system

Continuous transformation $T \curvearrowright (X, d)$ on a compact metric space.

Disclaimer

Please interrupt me.

In this talk all systems have discrete time \mathbb{Z} or \mathbb{N} (no flows or group actions)

Measurable dynamical system

Measure preserving transformation $T \curvearrowright (X, \mathcal{X}, \mu)$ on a probability space.

Topological dynamical system

Continuous transformation $T \curvearrowright (X, d)$ on a compact metric space.

Often they will be invertible, but not always

What is this talk about?

- 1 It is about a class of skew products $S \rtimes_{\tau} T$ (defined soon).

What is this talk about?

- 1 It is about a class of skew products $S \rtimes_{\tau} T$ (defined soon).
- 2 We are interested in the complexity of these systems: entropy and related concepts

What is this talk about?

- 1 It is about a class of skew products $S \rtimes_{\tau} T$ (defined soon).
- 2 We are interested in the complexity of these systems: entropy and related concepts
- 3 Natural problem: distinguish $S \rtimes_{\tau} T_1$ and $S \rtimes_{\tau} T_2$

What is this talk about?

- 1 It is about a class of skew products $S \rtimes_{\tau} T$ (defined soon).
- 2 We are interested in the complexity of these systems: entropy and related concepts
- 3 Natural problem: distinguish $S \rtimes_{\tau} T_1$ and $S \rtimes_{\tau} T_2$
- 4 This problem is well-studied for measure preserving systems; we study it for topological systems.

What is this talk about?

- 1 It is about a class of skew products $S \rtimes_{\tau} T$ (defined soon).
- 2 We are interested in the complexity of these systems: entropy and related concepts
- 3 Natural problem: distinguish $S \rtimes_{\tau} T_1$ and $S \rtimes_{\tau} T_2$
- 4 This problem is well-studied for measure preserving systems; we study it for topological systems.
- 5 We propose a solution based on slow entropy, an invariant introduced by Katok and Thouvenot.


The $[T, T^{-1}]$ system (informally)

Take your favorite measure preserving invertible transformation T on a probability space. Flip a coin, and apply either T or its inverse T^{-1} depending on the coin.

The $[T, T^{-1}]$ system (informally)

Take your favorite measure preserving invertible transformation T on a probability space. Flip a coin, and apply either T or its inverse T^{-1} depending on the coin.


For instance:



The $[T, T^{-1}]$ system (informally)

Take your favorite measure preserving invertible transformation T on a probability space. Flip a coin, and apply either T or its inverse T^{-1} depending on the coin.


For instance:



The $[T, T^{-1}]$ system (informally)

Take your favorite measure preserving invertible transformation T on a probability space. Flip a coin, and apply either T or its inverse T^{-1} depending on the coin.


For instance:



The $[T, T^{-1}]$ system (informally)

Take your favorite measure preserving invertible transformation T on a probability space. Flip a coin, and apply either T or its inverse T^{-1} depending on the coin.


For instance:



The $[T, T^{-1}]$ system (informally)

Take your favorite measure preserving invertible transformation T on a probability space. Flip a coin, and apply either T or its inverse T^{-1} depending on the coin.


For instance:



The $[T, T^{-1}]$ system (informally)

Take your favorite measure preserving invertible transformation T on a probability space. Flip a coin, and apply either T or its inverse T^{-1} depending on the coin.

For instance:



Take the space

$$\{-1, 1\}^{\mathbb{Z}} \times \mathbb{T}$$

and the transformation

$$(y, x) \mapsto (\sigma(y), T^{y(0)}(x))$$

The $[T, T^{-1}]$ system

The $[T, T^{-1}]$ system is the skew product

- Base: $\sigma \curvearrowright (\{-1, 1\}^{\mathbb{Z}}, \mu_{\frac{1}{2}, \frac{1}{2}})$
- Fiber: $T \curvearrowright (X, \mathcal{X}, \mu)$

$$\begin{aligned}\sigma \rtimes T &\curvearrowright \{-1, 1\}^{\mathbb{Z}} \times X \\ (y, x) &\mapsto (\sigma(y), T^{y(0)}(x))\end{aligned}$$

The $[T, T^{-1}]$ system

The $[T, T^{-1}]$ system is the skew product

- Base: $\sigma \curvearrowright (\{-1, 1\}^{\mathbb{Z}}, \mu_{\frac{1}{2}, \frac{1}{2}})$
- Fiber: $T \curvearrowright (X, \mathcal{X}, \mu)$

$$\begin{aligned}\sigma \rtimes T &\curvearrowright \{-1, 1\}^{\mathbb{Z}} \times X \\ (y, x) &\mapsto (\sigma(y), T^{y(0)}(x))\end{aligned}$$

In 1982 Kalikow proved that if T has positive entropy, then $\sigma \rtimes T$ is K and not Bernoulli (first “natural” known examples).

The $[T, T^{-1}]$ system

The $[T, T^{-1}]$ system is the skew product

- Base: $\sigma \curvearrowright (\{-1, 1\}^{\mathbb{Z}}, \mu_{\frac{1}{2}, \frac{1}{2}})$
- Fiber: $T \curvearrowright (X, \mathcal{X}, \mu)$

$$\begin{aligned}\sigma \rtimes T &\curvearrowright \{-1, 1\}^{\mathbb{Z}} \times X \\ (y, x) &\mapsto (\sigma(y), T^{y(0)}(x))\end{aligned}$$

In 1982 Kalikow proved that if T has positive entropy, then $\sigma \rtimes T$ is K and not Bernoulli (first “natural” known examples).

Question: are they all isomorphic copies of the same system, or are they many different examples?

The entropy of the fiber disappears

How can we distinguish these systems?

The entropy of the fiber disappears

How can we distinguish these systems?

One can prove that all of these systems have the same entropy

$$h_{\mu_{\frac{1}{2}, \frac{1}{2}} \times \mu}(\sigma \rtimes T) = h_{\mu_{\frac{1}{2}, \frac{1}{2}}}(\sigma) = \log 2$$

The entropy of the fiber disappears

How can we distinguish these systems?

One can prove that all of these systems have the same entropy

$$h_{\mu_{\frac{1}{2}, \frac{1}{2}} \times \mu}(\sigma \rtimes T) = h_{\mu_{\frac{1}{2}, \frac{1}{2}}}(\sigma) = \log 2$$

Is the entropy of the fiber an isomorphism invariant within this class?

The entropy of the fiber disappears

How can we distinguish these systems?

One can prove that all of these systems have the same entropy

$$h_{\mu_{\frac{1}{2}, \frac{1}{2}} \times \mu}(\sigma \rtimes T) = h_{\mu_{\frac{1}{2}, \frac{1}{2}}}(\sigma) = \log 2$$

Is the entropy of the fiber an isomorphism invariant within this class?:

$\sigma \rtimes T_1$ measurably isomorphic to $\sigma \rtimes T_2 \Rightarrow T_1$ and T_2 have the same entropy.

The entropy of the fiber disappears

How can we distinguish these systems?

One can prove that all of these systems have the same entropy

$$h_{\mu_{\frac{1}{2}, \frac{1}{2}} \times \mu}(\sigma \rtimes T) = h_{\mu_{\frac{1}{2}, \frac{1}{2}}}(\sigma) = \log 2$$

Is the entropy of the fiber an isomorphism invariant within this class?:

$\sigma \rtimes T_1$ measurably isomorphic to $\sigma \rtimes T_2 \Rightarrow T_1$ and T_2 have the same entropy.

Theorem (Austin 2015)

Yes. There is an invariant I with $I_{\mu_{\frac{1}{2}, \frac{1}{2}} \times \mu}(\sigma \rtimes T) = h_{\mu}(T)$.

Generalized $[T, T^{-1}]$ systems

Generalized $[T, T^{-1}]$ systems

Generalized $[T, T^{-1}]$ systems are skew products:

Generalized $[T, T^{-1}]$ systems

Generalized $[T, T^{-1}]$ systems are skew products:

- 1 Base: $S \curvearrowright (Y, \mathcal{Y}, \nu)$

Generalized $[T, T^{-1}]$ systems

Generalized $[T, T^{-1}]$ systems are skew products:

- 1 Base: $S \curvearrowright (Y, \mathcal{Y}, \nu)$
- 2 Cocycle: $\tau: Y \rightarrow \mathbb{Z}$ measurable

Generalized $[T, T^{-1}]$ systems

Generalized $[T, T^{-1}]$ systems are skew products:

- ① Base: $S \curvearrowright (Y, \mathcal{Y}, \nu)$
- ② Cocycle: $\tau: Y \rightarrow \mathbb{Z}$ measurable
- ③ Fiber: $T \curvearrowright (X, \mathcal{X}, \mu)$ invertible

Generalized $[T, T^{-1}]$ systems

Generalized $[T, T^{-1}]$ systems are skew products:

- ① Base: $S \curvearrowright (Y, \mathcal{Y}, \nu)$
- ② Cocycle: $\tau: Y \rightarrow \mathbb{Z}$ measurable
- ③ Fiber: $T \curvearrowright (X, \mathcal{X}, \mu)$ invertible

The new transformation is

$$S \rtimes_{\tau} T \curvearrowright (Y, \mathcal{Y}, \nu) \times (X, \mathcal{X}, \mu)$$

$$(y, x) \mapsto (S(y), T^{\tau(y)}(x))$$

What happens in the general case?

For which choices of $S \curvearrowright (Y, \mathcal{Y}, \nu)$ and $\tau: Y \rightarrow \mathbb{Z}$ is the entropy of the fiber an isomorphism invariant in the corresponding class of skew products?

What happens in the general case?

For which choices of $S \curvearrowright (Y, \mathcal{Y}, \nu)$ and $\tau: Y \rightarrow \mathbb{Z}$ is the entropy of the fiber an isomorphism invariant in the corresponding class of skew products?

- 1 The one-sided Bernoulli shift $S \curvearrowright (\{-1, 1\}^{\mathbb{N}}, \mu_{\frac{1}{2}, \frac{1}{2}})$ and $\tau(y) = y(0)$ (Heicklen, Hoffman, Rudolf 2000; Ball 2003)

What happens in the general case?

For which choices of $S \curvearrowright (Y, \mathcal{Y}, \nu)$ and $\tau: Y \rightarrow \mathbb{Z}$ is the entropy of the fiber an isomorphism invariant in the corresponding class of skew products?

- 1 The one-sided Bernoulli shift $S \curvearrowright (\{-1, 1\}^{\mathbb{N}}, \mu_{\frac{1}{2}, \frac{1}{2}})$ and $\tau(y) = y(0)$ (Heicklen, Hoffman, Rudolf 2000; Ball 2003)
- 2 The two-sided Bernoulli shift $S \curvearrowright (\{-1, 1\}^{\mathbb{Z}}, \mu_{\frac{1}{2}, \frac{1}{2}})$ and $\tau(y) = y(0)$ (Austin 2015)

What happens in the general case?

For which choices of $S \curvearrowright (Y, \mathcal{Y}, \nu)$ and $\tau: Y \rightarrow \mathbb{Z}$ is the entropy of the fiber an isomorphism invariant in the corresponding class of skew products?

- 1 The one-sided Bernoulli shift $S \curvearrowright (\{-1, 1\}^{\mathbb{N}}, \mu_{\frac{1}{2}, \frac{1}{2}})$ and $\tau(y) = y(0)$ (Heicklen, Hoffman, Rudolf 2000; Ball 2003)
- 2 The two-sided Bernoulli shift $S \curvearrowright (\{-1, 1\}^{\mathbb{Z}}, \mu_{\frac{1}{2}, \frac{1}{2}})$ and $\tau(y) = y(0)$ (Austin 2015)
- 3 A mixing SFT $S \curvearrowright (Y, \mathcal{Y}, \nu)$, ν a Gibbs measure, and $\tau: Y \rightarrow \mathbb{Z}$ satisfying a technical condition (Austin 2015)

What happens in the general case?

For which choices of $S \curvearrowright (Y, \mathcal{Y}, \nu)$ and $\tau: Y \rightarrow \mathbb{Z}$ is it true that the fiber entropy is an isomorphism invariant in this corresponding class of skew products?

- No known results with zero entropy systems in the base.

What happens in the general case?

For which choices of $S \curvearrowright (Y, \mathcal{Y}, \nu)$ and $\tau: Y \rightarrow \mathbb{Z}$ is it true that the fiber entropy is an isomorphism invariant in this corresponding class of skew products?

- No known results with zero entropy systems in the base.
- What happens for topological systems?

What happens in the general case?

For which choices of $S \curvearrowright (Y, \mathcal{Y}, \nu)$ and $\tau: Y \rightarrow \mathbb{Z}$ is it true that the fiber entropy is an isomorphism invariant in this corresponding class of skew products?

- No known results with zero entropy systems in the base.
- What happens for topological systems?
- We propose a solution, it works for a diverse class of S and τ .

Topological skew products

We consider skew products given by:

Topological skew products

We consider skew products given by:

- 1 Base: a subshift (Y, S)

Topological skew products

We consider skew products given by:

- 1 Base: a subshift (Y, S)
- 2 Cocycle: $\tau: Y \rightarrow \mathbb{Z}$ continuous

Topological skew products

We consider skew products given by:

- 1 Base: a subshift (Y, S)
- 2 Cocycle: $\tau: Y \rightarrow \mathbb{Z}$ continuous
- 3 Fiber: $T \curvearrowright (X, d)$ invertible

Topological skew products

We consider skew products given by:

- 1 Base: a subshift (Y, S)
- 2 Cocycle: $\tau: Y \rightarrow \mathbb{Z}$ continuous
- 3 Fiber: $T \curvearrowright (X, d)$ invertible

The new transformation is

$$S \rtimes_{\tau} T \curvearrowright Y \times X$$
$$(y, x) \mapsto (S(y), T^{\tau(y)}(x))$$

The problem

The class of systems

For fixed (Y, S, τ) , we consider the family of skew products $S \rtimes_{\tau} T$, where (X, T) is an arbitrary invertible topological dynamical system.

The problem

The class of systems

For fixed (Y, S, τ) , we consider the family of skew products $S \rtimes_{\tau} T$, where (X, T) is an arbitrary invertible topological dynamical system.

Problem

Find an invariant that captures the topological entropy of T .

The problem

The class of systems

For fixed (Y, S, τ) , we consider the family of skew products $S \times_{\tau} T$, where (X, T) is an arbitrary invertible topological dynamical system.

Problem

Find an invariant that captures ~~topological entropy~~ *entropy-type invariants* of T .

Intuition

The problem is interesting when the entropy of the fiber “disappears”. Informally,

$\# \text{ of } (n, \epsilon)\text{-Bowen balls} \approx e^{n \cdot h_{\text{top}}(S) + f(n)h_{\text{top}}(T)}$
needed to cover the space

for some sublinear $f(n)$.

Intuition

The problem is interesting when the entropy of the fiber “disappears”. Informally,

$\# \text{ of } (n, \epsilon)\text{-Bowen balls} \approx e^{n \cdot h_{\text{top}}(S) + f(n)h_{\text{top}}(T)}$
needed to cover the space

for some sublinear $f(n)$.

We want to capture the contribution of $h_{\text{top}}(T)$ in this relation.

of (n, ϵ) -Bowen balls $:= \text{spa}(T, n, \epsilon)$ $n \in \mathbb{N}, \epsilon > 0$
needed to cover the space

Topological slow entropy

Definition (Katok and Thouvenot '97)

Let $\mathbf{a} = \{a_n(t)\}_{n \in \mathbb{N}, t > 0}$ be a family of functions $(0, \infty) \rightarrow (0, \infty)$ with

- for each n , $t \rightarrow a_n(t)$ is monotone,
- for each t , $\lim_{n \rightarrow \infty} a_n(t) = \infty$

Topological slow entropy

Definition (Katok and Thouvenot '97)

Let $\mathbf{a} = \{a_n(t)\}_{n \in \mathbb{N}, t > 0}$ be a family of functions $(0, \infty) \rightarrow (0, \infty)$ with

- for each n , $t \rightarrow a_n(t)$ is monotone,
- for each t , $\lim_{n \rightarrow \infty} a_n(t) = \infty$

The upper slow entropy of (X, T) with scale \mathbf{a} is

$$\overline{\text{ent}}_{\mathbf{a}}(T) = \lim_{\epsilon \rightarrow 0} \overline{\text{ent}}_{\mathbf{a}}(T, \epsilon)$$

$$\overline{\text{ent}}_{\mathbf{a}}(T, \epsilon) = \sup(\{0\} \cup \{t > 0 : \limsup_{n \rightarrow \infty} \frac{\text{spa}(T, n, \epsilon)}{a_n(t)} > 0\})$$

$$\overline{\text{ent}}_{\mathbf{a}}(T, \epsilon) = \sup(\{0\} \cup \{t > 0 : \limsup_{n \rightarrow \infty} \frac{\text{spa}(T, n, \epsilon)}{a_n(t)} > 0\})$$

- One defines the lower slow entropy $\underline{\text{ent}}_{\mathbf{a}}(T)$ by taking \liminf instead of \limsup .

$$\overline{\text{ent}}_{\mathbf{a}}(T, \epsilon) = \sup(\{0\} \cup \{t > 0 : \limsup_{n \rightarrow \infty} \frac{\text{spa}(T, n, \epsilon)}{a_n(t)} > 0\})$$

- One defines the lower slow entropy $\underline{\text{ent}}_{\mathbf{a}}(T)$ by taking \liminf instead of \limsup .
- The upper and lower slow entropy may not coincide (i.e. when $n \rightarrow \text{spa}(T, n, \epsilon)$ behave differently along different subsequences)

$$\overline{\text{ent}}_{\mathbf{a}}(T, \epsilon) = \sup(\{0\} \cup \{t > 0 : \limsup_{n \rightarrow \infty} \frac{\text{spa}(T, n, \epsilon)}{a_n(t)} > 0\})$$

- One defines the lower slow entropy $\underline{\text{ent}}_{\mathbf{a}}(T)$ by taking \liminf instead of \limsup .
- The upper and lower slow entropy may not coincide (i.e. when $n \rightarrow \text{spa}(T, n, \epsilon)$ behave differently along different subsequences)
- With scale $\{e^{nt}\}_{n \in \mathbb{N}, t > 0}$ we recover topological entropy.

$$\overline{\text{ent}}_{\mathbf{a}}(T, \epsilon) = \sup(\{0\} \cup \{t > 0 : \limsup_{n \rightarrow \infty} \frac{\text{spa}(T, n, \epsilon)}{a_n(t)} > 0\})$$

- One defines the lower slow entropy $\underline{\text{ent}}_{\mathbf{a}}(T)$ by taking \liminf instead of \limsup .
- The upper and lower slow entropy may not coincide (i.e. when $n \rightarrow \text{spa}(T, n, \epsilon)$ behave differently along different subsequences)
- With scale $\{e^{nt}\}_{n \in \mathbb{N}, t > 0}$ we recover topological entropy.
- With scale $\{n^t\}_{n \in \mathbb{N}, t > 0}$ we obtain the so called polynomial complexity or polynomial entropy of the system.

The main result

The main result

Theorem (C, arXiv:2506.17932)

Let (Y, S) be a subshift and let $\tau: Y \rightarrow \mathbb{Z}$ be continuous. Then there is a scale \mathbf{a} depending only on (Y, S, τ) such that for every invertible topological dynamical system (X, T) we have

$$\overline{\text{ent}}_{\mathbf{a}}(S \rtimes_{\tau} T) = \underline{\text{ent}}_{\mathbf{a}}(S \rtimes_{\tau} T) = h_{\text{top}}(T),$$

provided that τ satisfies the condition of being λ -unbounded for some $\lambda > 0$ (defined soon).

The main result

Theorem (C, arXiv:2506.17932)

Let (Y, S) be a subshift and let $\tau: Y \rightarrow \mathbb{Z}$ be continuous. Then there is a scale \mathbf{a} depending only on (Y, S, τ) such that for every invertible topological dynamical system (X, T) we have

$$\overline{\text{ent}}_{\mathbf{a}}(S \rtimes_{\tau} T) = \underline{\text{ent}}_{\mathbf{a}}(S \rtimes_{\tau} T) = h_{\text{top}}(T),$$

provided that τ satisfies the condition of being λ -unbounded for some $\lambda > 0$ (defined soon).

A relative version of the main result

Theorem (C, arXiv:2506.17932)

Let (Y, S) be a subshift and let $\tau: Y \rightarrow \{-1, 0, 1\}$ be continuous. For every scale \mathbf{b} we can find a scale \mathbf{c} such that for every invertible topological dynamical system (X, T) we have

$$\underline{\text{ent}}_{\mathbf{b}}(T) \leq \overline{\text{ent}}_{\mathbf{c}}(S \rtimes_{\tau} T) \leq \overline{\text{ent}}_{\mathbf{b}}(T)$$

provided that τ satisfies the condition of being λ -unbounded for some $\lambda > 0$ (defined soon).

Thus we can recover the slow entropy of T with scale b provided we restrict to systems with the property $\overline{\text{ent}}_{\mathbf{b}}(T) = \underline{\text{ent}}_{\mathbf{b}}(T)$.

The property of being λ -unbounded

- For simplicity we will assume that Y is a subshift on symbols $\{-1, 1\}$ and $\tau(y) = y(0)$.

The property of being λ -unbounded

- For simplicity we will assume that Y is a subshift on symbols $\{-1, 1\}$ and $\tau(y) = y(0)$.
- For $y \in Y$ we interpret $N \rightarrow \sum_{n=0}^{N-1} \tau(S^n(y)) = y_0 + \dots + y_{N-1}$ as a walk on \mathbb{Z} , and we define the *range* $R_N(y)$ as the set of places in \mathbb{Z} it visits in N steps:

$$R_N(y) = \left\{ \sum_{n=0}^{i-1} \tau(S^n(y)) : 0 \leq i < N \right\}$$

The property of being λ -unbounded

- For simplicity we will assume that Y is a subshift on symbols $\{-1, 1\}$ and $\tau(y) = y(0)$.
- For $y \in Y$ we interpret $N \rightarrow \sum_{n=0}^{N-1} \tau(S^n(y)) = y_0 + \cdots + y_{N-1}$ as a walk on \mathbb{Z} , and we define the *range* $R_N(y)$ as the set of places in \mathbb{Z} it visits in N steps:

$$R_N(y) = \left\{ \sum_{n=0}^{i-1} \tau(S^n(y)) : 0 \leq i < N \right\}$$

- The condition says that $R_N(y)$ is large with positive “probability” λ , this refers to the proportion of words in the language of Y having some property.

The property of being λ -unbounded

- For simplicity we will assume that Y is a subshift on symbols $\{-1, 1\}$ and $\tau(y) = y(0)$.
- For $y \in Y$ we interpret $N \rightarrow \sum_{n=0}^{N-1} \tau(S^n(y)) = y_0 + \dots + y_{N-1}$ as a walk on \mathbb{Z} , and we define the *range* $R_N(y)$ as the set of places in \mathbb{Z} it visits in N steps:

$$R_N(y) = \left\{ \sum_{n=0}^{i-1} \tau(S^n(y)) : 0 \leq i < N \right\}$$

- The condition says that $R_N(y)$ is large with positive “probability” λ , this refers to the proportion of words in the language of Y having some property.
- $R_N(y)$ depends only on $y_0 \dots y_{N-1}$, so we can also write $R_N(w)$ for a word w of length N in symbols $\{-1, 1\}$.

The property of being λ -unbounded

- For simplicity we will assume that Y is a subshift on symbols $\{-1, 1\}$ and $\tau(y) = y(0)$.
- For $y \in Y$ we interpret $N \rightarrow \sum_{n=0}^{N-1} \tau(S^n(y)) = y_0 + \dots + y_{N-1}$ as a walk on \mathbb{Z} , and we define the *range* $R_N(y)$ as the set of places in \mathbb{Z} it visits in N steps:

$$R_N(y) = \left\{ \sum_{n=0}^{i-1} \tau(S^n(y)) : 0 \leq i < N \right\}$$

- The condition says that $R_N(y)$ is large with positive “probability” λ , this refers to the proportion of words in the language of Y having some property.
- $R_N(y)$ depends only on $y_0 \dots y_{N-1}$, so we can also write $R_N(w)$ for a word w of length N in symbols $\{-1, 1\}$.

Definition

Given (Y, S, τ) as before, we say that τ is λ -unbounded ($\lambda > 0$) if for all $C \in \mathbb{N}$

$$\liminf_{n \rightarrow \infty} \frac{|\{w \in L_n(Y) : |R_n(w)| \geq C\}|}{|L_n(Y)|} \geq \lambda$$

Definition

Given (Y, S, τ) as before, we say that τ is λ -unbounded ($\lambda > 0$) if for all $C \in \mathbb{N}$

$$\liminf_{n \rightarrow \infty} \frac{|\{w \in L_n(Y) : |R_n(w)| \geq C\}|}{|L_n(Y)|} \geq \lambda$$

- If $n \rightarrow \sum_{i=0}^{n-1} \tau(S^i y)$ unbounded for all y , then this holds.

Definition

Given (Y, S, τ) as before, we say that τ is λ -unbounded ($\lambda > 0$) if for all $C \in \mathbb{N}$

$$\liminf_{n \rightarrow \infty} \frac{|\{w \in L_n(Y) : |R_n(w)| \geq C\}|}{|L_n(Y)|} \geq \lambda$$

- If $n \rightarrow \sum_{i=0}^{n-1} \tau(S^i y)$ unbounded for all y , then this holds.
- Also true for $Y = \{-1, 1\}^{\mathbb{Z}}$.

Examples

Lets review some choices of (Y, S) and τ that have this property.

Examples

Lets review some choices of (Y, S) and τ that have this property.

Example

(Y, S) is a minimal subshift and $\tau: Y \rightarrow \mathbb{Z}$ is not a coboundary (i.e. $\tau = g - g \circ S$ for some continuous $g: Y \rightarrow \mathbb{R}$).

Example

Y is a Sturmian subshift on symbols $\{0, 1\}$ and $\tau(y) = y(0)$.

Example

Y is a Sturmian subshift on symbols $\{0, 1\}$ and $\tau(y) = y(0)$.

- For any T we have $h_{top}(S \rtimes_{\tau} T) = h_{top}(S) + \alpha h_{top}(T)$, so applying the main theorem is an overkill.


Example

Y is a Sturmian subshift on symbols $\{0, 1\}$ and $\tau(y) = y(0)$.

- For any T we have $h_{top}(S \rtimes_{\tau} T) = h_{top}(S) + \alpha h_{top}(T)$, so applying the main theorem is an overkill.
- The conclusion of the “relative main theorem” is nontrivial: if T_1 and T_2 have zero entropy but different polynomial complexity ($\underline{\text{ent}}_{nt}(T_1) < \overline{\text{ent}}_{nt}(T_2)$), one obtains a scale for slow entropy that distinguishes $S \rtimes_{\tau} T_1$ and $S \rtimes_{\tau} T_2$.

Example

The deterministic random walk (the subshift copy).



Example


The deterministic random walk (the subshift copy).

- Choose an irrational α , and take the function $\rho: [0, 1) \rightarrow \{-1, 1\}$ with value 1 on $[0, 1/2)$ and value -1 over $[1/2, 1)$.


Example

The deterministic random walk (the subshift copy).

- Choose an irrational α , and take the function $\rho: [0, 1) \rightarrow \{-1, 1\}$ with value 1 on $[0, 1/2)$ and value -1 over $[1/2, 1)$.
- Let (Y, S) be the smallest subshift on symbols $\{-1, 1\}$ containing the sequence $(\rho(x + n\alpha))_{n \in \mathbb{Z}}$.




Example



The deterministic random walk (the subshift copy).


- Choose an irrational α , and take the function $\rho: [0, 1) \rightarrow \{-1, 1\}$ with value 1 on $[0, 1/2)$ and value -1 over $[1/2, 1)$.
- Let (Y, S) be the smallest subshift on symbols $\{-1, 1\}$ containing the sequence $(\rho(x + n\alpha))_{n \in \mathbb{Z}}$.
- Take the cocycle $\tau: Y \rightarrow \mathbb{Z}$, $\tau(y) = y(0)$.



Example

The deterministic random walk (the subshift copy).


- Choose an irrational α , and take the function $\rho: [0, 1) \rightarrow \{-1, 1\}$ with value 1 on $[0, 1/2)$ and value -1 over $[1/2, 1)$.
- Let (Y, S) be the smallest subshift on symbols $\{-1, 1\}$ containing the sequence $(\rho(x + n\alpha))_{n \in \mathbb{Z}}$.
- Take the cocycle $\tau: Y \rightarrow \mathbb{Z}$, $\tau(y) = y(0)$.
- Here $h_{\text{top}}(S \rtimes_{\tau} T) = 0$ for all choices of T .



Example

The deterministic random walk (the subshift copy).

- Choose an irrational α , and take the function $\rho: [0, 1) \rightarrow \{-1, 1\}$ with value 1 on $[0, 1/2)$ and value -1 over $[1/2, 1)$.
- Let (Y, S) be the smallest subshift on symbols $\{-1, 1\}$ containing the sequence $(\rho(x + n\alpha))_{n \in \mathbb{Z}}$.
- Take the cocycle $\tau: Y \rightarrow \mathbb{Z}$, $\tau(y) = y(0)$.
- Here $h_{\text{top}}(S \rtimes_{\tau} T) = 0$ for all choices of T .
- The main theorem shows that the entropy of T equals the slow entropy of $S \rtimes_{\tau} T$ at some scale.



Example

The deterministic random walk (the subshift copy).

- Choose an irrational α , and take the function $\rho: [0, 1) \rightarrow \{-1, 1\}$ with value 1 on $[0, 1/2)$ and value -1 over $[1/2, 1)$.
- Let (Y, S) be the smallest subshift on symbols $\{-1, 1\}$ containing the sequence $(\rho(x + n\alpha))_{n \in \mathbb{Z}}$.
- Take the cocycle $\tau: Y \rightarrow \mathbb{Z}$, $\tau(y) = y(0)$.
- Here $h_{\text{top}}(S \rtimes_{\tau} T) = 0$ for all choices of T .
- The main theorem shows that the entropy of T equals the slow entropy of $S \rtimes_{\tau} T$ at some scale.
- The same is true for the slow entropy of T .

Examples

Example

The full shift $\{-1, 1\}^{\mathbb{Z}}$ and $\tau(y) = y(0)$

Examples

Example

The full shift $\{-1, 1\}^{\mathbb{Z}}$ and $\tau(y) = y(0)$

- Here $h_{top}(S \rtimes_{\tau} T) = \log(2) + f(h_{top}(T))$, for some $f: [0, \infty) \rightarrow [0, \infty)$ strictly increasing.

Examples

Example

The full shift $\{-1, 1\}^{\mathbb{Z}}$ and $\tau(y) = y(0)$

- Here $h_{top}(S \rtimes_{\tau} T) = \log(2) + f(h_{top}(T))$, for some $f: [0, \infty) \rightarrow [0, \infty)$ strictly increasing.
- Applying the main theorem is an overkill.

Examples

Example

The full shift $\{-1, 1\}^{\mathbb{Z}}$ and $\tau(y) = y(0)$

- Here $h_{\text{top}}(S \rtimes_{\tau} T) = \log(2) + f(h_{\text{top}}(T))$, for some $f: [0, \infty) \rightarrow [0, \infty)$ strictly increasing.
- Applying the main theorem is an overkill.
- The conclusion of the “relative main theorem” is nontrivial: if T_1 and T_2 have zero entropy but different polynomial complexity ($\underline{\text{ent}}_{nt}(T_1) < \overline{\text{ent}}_{nt}(T_2)$), one obtains a scale for slow entropy that distinguishes $S \rtimes_{\tau} T_1$ and $S \rtimes_{\tau} T_2$.

Proof idea

Proof idea

Notation

Take an invertible system $T \curvearrowright (X, d)$.

Given $F \subset \mathbb{Z}$ finite, we write

$$d_F(x, y) = \max\{d(T^i(x), T^i(y)) : i \in F\}$$

Notation

Take an invertible system $T \curvearrowright (X, d)$.

Given $F \subset \mathbb{Z}$ finite, we write

$$d_F(x, y) = \max\{d(T^i(x), T^i(y)) : i \in F\}$$

and

$$\text{spa}(T, F, \epsilon)$$

equals the minimal number of Bowen d_F, ϵ -balls needed to cover X .

Proof idea

For the systems we consider $S \rtimes_{\tau} T$,

$$\text{spa}(S \rtimes_{\tau} T, \{0, \dots, N-1\}, \epsilon) \approx$$

Proof idea

For the systems we consider $S \rtimes_{\tau} T$,

$$\text{spa}(S \rtimes_{\tau} T, \{0, \dots, N-1\}, \epsilon) \approx \sum_{w \in L_N(Y)} \text{spa}(T, R_N(w), \epsilon)$$

Proof idea

For the systems we consider $S \rtimes_{\tau} T$,

$$\text{spa}(S \rtimes_{\tau} T, \{0, \dots, N-1\}, \epsilon) \approx \sum_{w \in L_N(Y)} \text{spa}(T, R_N(w), \epsilon)$$

If $\tau(y) \in \{-1, 1\}$ for all y then the sets $R_N(w)$ are intervals

Proof idea

For the systems we consider $S \rtimes_{\tau} T$,

$$\text{spa}(S \rtimes_{\tau} T, \{0, \dots, N-1\}, \epsilon) \approx \sum_{w \in L_N(Y)} \text{spa}(T, R_N(w), \epsilon)$$

If $\tau(y) \in \{-1, 1\}$ for all y then the sets $R_N(w)$ are intervals

From the definition of topological entropy, for an interval $F \subset \mathbb{Z}$ we have $\text{spa}(T, F, \epsilon) \approx e^{h_{\text{top}}(T)|F|}$.

Proof idea

For the systems we consider $S \rtimes_{\tau} T$,

$$\text{spa}(S \rtimes_{\tau} T, \{0, \dots, N-1\}, \epsilon) \approx \sum_{w \in L_N(Y)} \text{spa}(T, R_N(w), \epsilon)$$

If $\tau(y) \in \{-1, 1\}$ for all y then the sets $R_N(w)$ are intervals

From the definition of topological entropy, for an interval $F \subset \mathbb{Z}$ we have $\text{spa}(T, F, \epsilon) \approx e^{h_{\text{top}}(T)|F|}$.

$$\approx \sum_{w \in L_N(Y)} e^{|R_N(w)|h_{\text{top}}(T)}$$

Proof idea

We concluded

$$\text{spa}(S \rtimes_{\tau} T, \{0, \dots, N-1\}, \epsilon)$$

Proof idea

We concluded

$$\mathrm{spa}(S \rtimes_{\tau} T, \{0, \dots, N-1\}, \epsilon) \approx \sum_{w \in L_N(Y)} e^{|R_N(w)| h_{\mathrm{top}}(T)}$$

Proof idea

We concluded

$$\mathrm{spa}(S \rtimes_{\tau} T, \{0, \dots, N-1\}, \epsilon) \approx \sum_{w \in L_N(Y)} e^{|R_N(w)| h_{\mathrm{top}}(T)}$$

The scale for the first main theorem is

$$a_N(t) = \sum_{w \in L_N(Y)} e^{|R_N(w)| t}$$

Proof idea

We concluded

$$\text{spa}(S \rtimes_{\tau} T, \{0, \dots, N-1\}, \epsilon) \approx \sum_{w \in L_N(Y)} e^{|R_N(w)| h_{\text{top}}(T)}$$

The scale for the first main theorem is

$$a_N(t) = \sum_{w \in L_N(Y)} e^{|R_N(w)| t}$$

We have

$$\limsup_{N \rightarrow \infty} \frac{\text{spa}(S \rtimes_{\tau} T, \{0, \dots, N-1\}, \epsilon)}{a_N(t)}$$

is 0 for $t > h_{\text{top}}(T)$, and ∞ for $t < h_{\text{top}}(T)$. Thus
 $\overline{\text{ent}}_{\mathbf{a}}(S \rtimes_{\tau} T) = h_{\text{top}}(T)$.

General case

If τ takes values outside $\{-1, 0, 1\}$, the sets $R_N(w)$ may have holes, and this causes extra growth of $\text{spa}(T, R_N(w), \epsilon)$.

For instance, if $R_N(w) = \{2, 4, \dots, 2N\}$,

$$\text{spa}(T, R_N(w), \epsilon) \approx e^{2 \cdot h_{\text{top}}(T) |R_N(w)|}$$

It is possible to quantify this extra contribution and include it to the scale, so that they “cancels out”.

Comments and questions

Comments and questions

Question: is it possible to prove analogous result for measure preserving systems using measure-theoretic slow entropy?

Comments and questions

Question: is it possible to prove analogous result for measure preserving systems using measure-theoretic slow entropy?

The scales constructed here don't work for these purposes (in general).

Thanks

Thanks

Thanks

