

CS 511, Fall 2024, Lecture Slides 28

Sequential Compactness

Assaf Kfoury

16 October 2024

A Full Binary Tree $\mathcal{T}_{\text{full}}$

Given arbitrary $a, b \in \mathbb{Q}$ with $a \leq b$, we use two kinds of intervals in this presentation:

closed $[a, b] \stackrel{\text{def}}{=} \{ r \mid a \leq r \leq b \}$

left-closed $[a, b) \stackrel{\text{def}}{=} \{ r \mid a \leq r < b \}$


A Full Binary Tree $\mathcal{T}_{\text{full}}$

Given arbitrary $a, b \in \mathbb{Q}$ with $a \leq b$, we use two kinds of intervals in this presentation:

closed $[a, b] \stackrel{\text{def}}{=} \{ r \mid a \leq r \leq b \}$


left-closed $[a, b) \stackrel{\text{def}}{=} \{ r \mid a \leq r < b \}$

Starting with the interval $[0, 1]$ at the top, as the root node, we repeatedly divide intervals into two disjoint halves, producing an infinite full binary tree $\mathcal{T}_{\text{full}}$:




A Full Binary Tree $\mathcal{T}_{\text{full}}$

More systematically, in order to exhibit a common pattern for all the intervals in $\mathcal{T}_{\text{full}}$:



A Full Binary Tree $\mathcal{T}_{\text{full}}$

More systematically, in order to exhibit a common pattern for all the intervals in $\mathcal{T}_{\text{full}}$:



Important observation:

A path from the root of $\mathcal{T}_{\text{full}}$ produces a strict nested sequence of intervals, e.g.:

$$[0, 1] \supsetneq [1 \cdot 2^{-1}, 2 \cdot 2^{-1}] \supsetneq [2 \cdot 2^{-2}, 3 \cdot 2^{-2}) \supsetneq \dots$$

This is a sequence of increasingly narrower intervals converging to a single number.

A Full Binary Tree $\mathcal{T}_{\text{full}}$


The nodes at the k -th level of $\mathcal{T}_{\text{full}}$ are, from left to right:

$$\underbrace{[0 \cdot 2^{-k}, 1 \cdot 2^{-k}) \quad [1 \cdot 2^{-k}, 2 \cdot 2^{-k}) \quad [2 \cdot 2^{-k}, 3 \cdot 2^{-k}) \quad \dots \quad [(2^k - 2) \cdot 2^{-k}, (2^k - 1) \cdot 2^{-k})}_{\text{left-closed}} \quad \underbrace{[(2^k - 1) \cdot 2^{-k}, 2^k \cdot 2^{-k}]}_{\text{closed}}$$

More succinctly, at level $k \geq 1$, there are:

- ▶ $(2^k - 1)$ left-closed intervals, each of the form $[(r-1) \cdot 2^{-k}, r \cdot 2^{-k})$ where $0 \leq r \leq 2^k - 1$,
- ▶ and one closed interval, the rightmost, $[(2^k - 1) \cdot 2^{-k}, 2^k \cdot 2^{-k}]$.

A Full Binary Tree $\mathcal{T}_{\text{full}}$ – once more




How to follow a path in $\mathcal{T}_{\text{full}}$ and prune it in the process

How to follow a path in $\mathcal{T}_{\text{full}}$ and prune it in the process

1. Let $A \subseteq [0, 1]$ be arbitrary, finite or infinite, which we use to follow a path in $\mathcal{T}_{\text{full}}$.


How to follow a path in $\mathcal{T}_{\text{full}}$ and prune it in the process

1. Let $A \subseteq [0, 1]$ be arbitrary, finite or infinite, which we use to follow a path in $\mathcal{T}_{\text{full}}$.
2. At every node I_n of $\mathcal{T}_{\text{full}}$:




How to follow a path in $\mathcal{T}_{\text{full}}$ and prune it in the process

1. Let $A \subseteq [0, 1]$ be arbitrary, finite or infinite, which we use to follow a path in $\mathcal{T}_{\text{full}}$.
2. At every node I_n of $\mathcal{T}_{\text{full}}$:




if $|A \cap I_{2n}| \geq |A \cap I_{2n+1}|$, replace I_{2n+1} by \bullet and prune the subtree of $\mathcal{T}_{\text{full}}$ rooted at I_{2n+1} :




How to follow a path in $\mathcal{T}_{\text{full}}$ and prune it in the process


1. Let $A \subseteq [0, 1]$ be arbitrary, finite or infinite, which we use to follow a path in $\mathcal{T}_{\text{full}}$.
2. At every node I_n of $\mathcal{T}_{\text{full}}$:



if $|A \cap I_{2n}| \geq |A \cap I_{2n+1}|$, replace I_{2n+1} by \bullet and prune the subtree of $\mathcal{T}_{\text{full}}$ rooted at I_{2n+1} :




if $|A \cap I_{2n}| < |A \cap I_{2n+1}|$, replace I_{2n} by \bullet and prune the subtree of $\mathcal{T}_{\text{full}}$ rooted at I_{2n} :




How to follow a path in $\mathcal{T}_{\text{full}}$ and prune it in the process


1. Let $A \subseteq [0, 1]$ be arbitrary, finite or infinite, which we use to follow a path in $\mathcal{T}_{\text{full}}$.
2. At every node I_n of $\mathcal{T}_{\text{full}}$:



if $|A \cap I_{2n}| \geq |A \cap I_{2n+1}|$, replace I_{2n+1} by \bullet and prune the subtree of $\mathcal{T}_{\text{full}}$ rooted at I_{2n+1} :




if $|A \cap I_{2n}| < |A \cap I_{2n+1}|$, replace I_{2n} by \bullet and prune the subtree of $\mathcal{T}_{\text{full}}$ rooted at I_{2n} :




3. Let $I_1 = I_{k_1} \supsetneq I_{k_2} \supsetneq I_{k_3} \supsetneq \cdots$ be the nested chain of intervals in the path followed using A , which is an infinite path by WKL (even if A is finite) converging to a single number b .

How to follow a path in $\mathcal{T}_{\text{full}}$ and prune it in the process


1. Let $A \subseteq [0, 1]$ be arbitrary, finite or infinite, which we use to follow a path in $\mathcal{T}_{\text{full}}$.
2. At every node I_n of $\mathcal{T}_{\text{full}}$:



if $|A \cap I_{2n}| \geq |A \cap I_{2n+1}|$, replace I_{2n+1} by \bullet and prune the subtree of $\mathcal{T}_{\text{full}}$ rooted at I_{2n+1} :



if $|A \cap I_{2n}| < |A \cap I_{2n+1}|$, replace I_{2n} by \bullet and prune the subtree of $\mathcal{T}_{\text{full}}$ rooted at I_{2n} :



3. Let $I_1 = I_{k_1} \supsetneq I_{k_2} \supsetneq I_{k_3} \supsetneq \dots$ be the nested chain of intervals in the path followed using A , which is an infinite path by WKL (even if A is finite) converging to a single number b .
4. The nested sequence $(A \cap I_{k_1}) \supsetneq (A \cap I_{k_2}) \supsetneq (A \cap I_{k_3}) \supsetneq \dots$ consists of increasingly narrower non-empty subsets of A converging to the same single point b .

(THIS PAGE INTENTIONALLY LEFT BLANK)