

# CS 511, Fall 2024, Lecture Slides 21

## Extended Example in First-Order Logic

Assaf Kfoury

7 October 2024

# several structures over the domain $\mathbb{N}$ (assume “ $\approx$ ” is available)

structures over the domain of natural numbers	vocabulary/signature	
	predicate symbols	function symbols
$\mathcal{N} \stackrel{\text{def}}{=} (\mathbb{N}, 0, S)$	$\mathcal{P} = \emptyset$	$\mathcal{F} = \{0, S\}$
$\mathcal{N}_1 \stackrel{\text{def}}{=} (\mathbb{N}, 0, S, <)$	$\mathcal{P} = \{<\}$	$\mathcal{F} = \{0, S\}$
$\mathcal{N}_2 \stackrel{\text{def}}{=} (\mathbb{N}, 0, S, <, +)$	$\mathcal{P} = \{<\}$	$\mathcal{F} = \{0, S, +\}$
$\mathcal{N}_3 \stackrel{\text{def}}{=} (\mathbb{N}, 0, S, <, +, \cdot)$	$\mathcal{P} = \{<\}$	$\mathcal{F} = \{0, S, +, \cdot\}$
$\mathcal{N}_4 \stackrel{\text{def}}{=} (\mathbb{N}, 0, S, <, +, \cdot, \text{pr})$ $\text{pr}(x) \stackrel{\text{def}}{=} \text{true iff } x \text{ is prime}$	$\mathcal{P} = \{<, \text{pr}\}$	$\mathcal{F} = \{0, S, +, \cdot\}$
$\mathcal{N}_5 \stackrel{\text{def}}{=} (\mathbb{N}, 0, S, <, +, \cdot, \text{pr}, \uparrow)$ $x \uparrow y \stackrel{\text{def}}{=} x^y$	$\mathcal{P} = \{<, \text{pr}\}$	$\mathcal{F} = \{0, S, +, \cdot, \uparrow\}$
$\mathcal{N}_6 \stackrel{\text{def}}{=} \dots$		

Question: Is a new predicate (function) definable from earlier ones?

# first-order definability over $\mathbb{N}$

- ▶ every number  $n$  is definable from 0 and  $S$  :

$$1 \stackrel{\text{def}}{=} S(0)$$

$$2 \stackrel{\text{def}}{=} S(S(0))$$

$$3 \stackrel{\text{def}}{=} S(S(S(0)))$$

$\dots$

$$n \stackrel{\text{def}}{=} S(\underbrace{\dots S(0) \dots}_n)$$

# first-order definability over $\mathbb{N}$

- ▶ every number  $n$  is definable from 0 and  $S$  :

$$1 \stackrel{\text{def}}{=} S(0)$$

$$2 \stackrel{\text{def}}{=} S(S(0))$$

$$3 \stackrel{\text{def}}{=} S(S(S(0)))$$

$$\begin{array}{c} \dots \\ n \stackrel{\text{def}}{=} S(\underbrace{\dots S(0) \dots}_n) \end{array}$$

- ▶ “ $S$ ” is definable from “ $+$ ” : for all  $m, n \in \mathbb{N}$ , we have  $S(m) = n$  iff  $m + 1 = n$

# first-order definability over $\mathbb{N}$

- ▶ every number  $n$  is definable from 0 and  $S$  :

$$1 \stackrel{\text{def}}{=} S(0)$$

$$2 \stackrel{\text{def}}{=} S(S(0))$$

$$3 \stackrel{\text{def}}{=} S(S(S(0)))$$

$$\begin{array}{c} \dots \\ n \stackrel{\text{def}}{=} S(\underbrace{\dots S(0) \dots}_n) \end{array}$$

- ▶ “ $S$ ” is definable from “ $+$ ” : for all  $m, n \in \mathbb{N}$ , we have  $S(m) = n$  iff  $m + 1 = n$   
formally: the sentence  $\forall x \forall y ( S(x) \approx y \leftrightarrow x + 1 \approx y )$  is true in  $\mathcal{N}_2$ ,  
which implies the graph of  $S^{\mathcal{N}_2}$  is defined by  $\varphi_S(x, y) = (x + 1 \approx y)$ .

# first-order definability over $\mathbb{N}$

- ▶ every number  $n$  is definable from 0 and  $S$  :

$$1 \stackrel{\text{def}}{=} S(0)$$

$$2 \stackrel{\text{def}}{=} S(S(0))$$

$$3 \stackrel{\text{def}}{=} S(S(S(0)))$$

$$\begin{array}{c} \dots \\ n \stackrel{\text{def}}{=} S(\underbrace{\dots S(0) \dots}_n) \end{array}$$

- ▶ “ $S$ ” is definable from “ $+$ ” : for all  $m, n \in \mathbb{N}$ , we have  $S(m) = n$  iff  $m + 1 = n$   
formally: the sentence  $\forall x \forall y ( S(x) \approx y \leftrightarrow x + 1 \approx y )$  is true in  $\mathcal{N}_2$ ,  
which implies the graph of  $S^{\mathcal{N}_2}$  is defined by  $\varphi_S(x, y) = (x + 1 \approx y)$ .
- ▶ is “ $+$ ” definable from “ $S$ ”? perhaps . . .

# first-order definability over $\mathbb{N}$

- ▶ every number  $n$  is definable from 0 and  $S$  :

$$1 \stackrel{\text{def}}{=} S(0)$$

$$2 \stackrel{\text{def}}{=} S(S(0))$$

$$3 \stackrel{\text{def}}{=} S(S(S(0)))$$

$$\dots$$
$$n \stackrel{\text{def}}{=} S(\underbrace{\dots S(0) \dots}_n)$$

- ▶ “ $S$ ” is definable from “ $+$ ” : for all  $m, n \in \mathbb{N}$ , we have  $S(m) = n$  iff  $m + 1 = n$   
formally: the sentence  $\forall x \forall y ( S(x) \approx y \leftrightarrow x + 1 \approx y )$  is true in  $\mathcal{N}_2$ ,  
which implies the graph of  $S^{\mathcal{N}_2}$  is defined by  $\varphi_S(x, y) = (x + 1 \approx y)$ .
- ▶ is “ $+$ ” definable from “ $S$ ”? perhaps . . .

for all  $m, n, p \in \mathbb{N}$ , we have  $m + n = p$  iff  $S(\underbrace{\dots S(m) \dots}_n) = p$

# first-order definability over $\mathbb{N}$

- ▶ every number  $n$  is definable from 0 and  $S$  :

$$1 \stackrel{\text{def}}{=} S(0)$$

$$2 \stackrel{\text{def}}{=} S(S(0))$$

$$3 \stackrel{\text{def}}{=} S(S(S(0)))$$

$$\vdots$$
$$n \stackrel{\text{def}}{=} S(\underbrace{\cdots S(0) \cdots}_n)$$

- ▶ “ $S$ ” is definable from “ $+$ ” : for all  $m, n \in \mathbb{N}$ , we have  $S(m) = n$  iff  $m + 1 = n$   
formally: the sentence  $\forall x \forall y (S(x) \approx y \leftrightarrow x + 1 \approx y)$  is true in  $\mathcal{N}_2$ ,  
which implies the graph of  $S^{\mathcal{N}_2}$  is defined by  $\varphi_S(x, y) = (x + 1 \approx y)$ .
- ▶ is “ $+$ ” definable from “ $S$ ”? perhaps . . .

for all  $m, n, p \in \mathbb{N}$ , we have  $m + n = p$  iff  $S(\underbrace{\cdots S(m) \cdots}_n) = p$

“formally”:  $\forall x \forall y \forall z [ \underbrace{S(\cdots S(x) \cdots)}_y \approx z \leftrightarrow x + y \approx z ]$

so perhaps  $\varphi_+(x, y, z) \stackrel{\text{def}}{=} ( \underbrace{S(\cdots S(x) \cdots)}_y \approx z \dots )$       Not quite!



# first-order definability over $\mathbb{N}$

## 1. FACT

“+” is **NOT** (first-order) definable from “0” and “S” (*difficult!*)

## first-order definability over $\mathbb{N}$

### 1. FACT

“+” is **NOT** (first-order) definable from “0” and “S” (*difficult!*)

### 2. FACT

“<” is (first-order) definable from “+” (*easy: try it!*)

## first-order definability over $\mathbb{N}$

### 1. FACT

“+” is **NOT** (first-order) definable from “0” and “S” (*difficult!*)

### 2. FACT

“<” is (first-order) definable from “+” (*easy: try it!*)

### 3. FACT

“+” is **NOT** (first-order) definable from “<”, “0”, and “S” (*difficult!*)

# first-order definability over $\mathbb{N}$

## 1. FACT

“+” is **NOT** (first-order) definable from “0” and “S” (*difficult!*)

## 2. FACT

“<” is (first-order) definable from “+” (*easy: try it!*)

## 3. FACT

“+” is **NOT** (first-order) definable from “<”, “0”, and “S” (*difficult!*)

## 4. FACT

“.” is **NOT** (first-order) definable from “0”, “S”, and “+”  
(no need to mention “<”) (*difficult!*)

# first-order definability over $\mathbb{N}$

## 1. FACT

“+” is **NOT** (first-order) definable from “0” and “S” (*difficult!*)

## 2. FACT

“<” is (first-order) definable from “+” (*easy: try it!*)

## 3. FACT

“+” is **NOT** (first-order) definable from “<”, “0”, and “S” (*difficult!*)

## 4. FACT

“.” is **NOT** (first-order) definable from “0”, “S”, and “+”  
(no need to mention “<”) (*difficult!*)

## 5. FACT

“+” is (first-order) definable from “<” and “.” (*tricky: try hint below!*)

# first-order definability over $\mathbb{N}$

## 1. FACT

“+” is **NOT** (first-order) definable from “0” and “S” (*difficult!*)

## 2. FACT

“<” is (first-order) definable from “+” (*easy: try it!*)

## 3. FACT

“+” is **NOT** (first-order) definable from “<”, “0”, and “S” (*difficult!*)

## 4. FACT

“.” is **NOT** (first-order) definable from “0”, “S”, and “+”  
(no need to mention “<”) (*difficult!*)

## 5. FACT

“+” is (first-order) definable from “<” and “.” (*tricky: try hint below!*)

*Hint.* Use the following equivalence for all  $m, n, p \in \mathbb{N}$

$(p = 0) \vee (p = m + n)$  iff

$$(m \cdot p + 1) \cdot (n \cdot p + 1) = p^2 \cdot (m \cdot n + 1) + 1$$

## first-order definability over $\mathbb{N}$

- ▶ is “pr” definable from  $\{0, S, <, +, \cdot\}$ ?

## first-order definability over $\mathbb{N}$

- ▶ is “pr” definable from  $\{0, S, <, +, \cdot\}$ ?

**YES**  $\text{pr}(n)$  is true iff  $\varphi(n)$  is true, where  $\varphi(x)$  is the WFF

$$\varphi(x) \stackrel{\text{def}}{=} \neg(x \approx 1) \wedge \forall y \forall z [ (x \approx y \cdot z) \rightarrow (y \approx 1 \vee z \approx 1) ]$$

- ▶ is “ $\uparrow$ ” definable from  $\{0, S, <, +, \cdot\}$ ?



## first-order definability over $\mathbb{N}$

- ▶ is “pr” definable from  $\{0, S, <, +, \cdot\}$ ?

**YES**  $\text{pr}(n)$  is true iff  $\varphi(n)$  is true, where  $\varphi(x)$  is the WFF

$$\varphi(x) \stackrel{\text{def}}{=} \neg(x \approx 1) \wedge \forall y \forall z [ (x \approx y \cdot z) \rightarrow (y \approx 1 \vee z \approx 1) ]$$

- ▶ is “ $\uparrow$ ” definable from  $\{0, S, <, +, \cdot\}$ ?

**YES**  $m = n \uparrow p$  iff  $\varphi(m, n, p)$  is true, where  $\varphi(x, y, z)$  is the WFF ... *(not very difficult: try it!)*

(THIS PAGE INTENTIONALLY LEFT BLANK)