

1 Problem 1

Find all integers $n > 1$ with the property that for each positive divisor d of n , we also have that

$$(d + 2) \mid (n + 2)$$

Solution. We first show no even integers satisfy our property. If n is even, there exists some $k \in \mathbb{N}$ such that $2k = n$, so $k \mid n$. Furthermore, $n + 2$ is even as well, so $\frac{n+2}{2}$ is an integer, specifically $\frac{n+2}{2} = \frac{n}{2} + 1 = k + 1$. So $2(k + 1) = n + 2 \implies k + 1 \mid n + 2$. Furthermore, $k + 1$ must be the greatest possible divisor of $n + 2$ that is not equal to $n + 2$, since if a divisor d was greater than $k + 1$, there is some integer $1 < m < 2$ where $dm = n + 2$, but no such integer m exists. But $k + 2 > k + 1$, and $k + 2 = \frac{n}{2} + 2 \neq n + 2$ when $n > 1$, so $k + 2 \nmid n + 2$. So there exists a divisor of n such that two more than it is not a divisor of $n + 2$.

Now, we consider odd composite integers n . Then there exists an integers $1 < d, q < n$ (not necessarily distinct) such that $qd = n$ (so $d \mid n$). Without loss of generality, let $d \geq q$. Then $n + 2 = qd + 2$. Note that both q and d must be odd, otherwise n would be divisible by 2 and would be even, which would be against our assumption. We can write $n + 2 = qd + 2 + 2q - 2q = q(d + 2) - 2(q - 1)$, so

$$(n + 2)/(d + 2) = q - 2(q - 1)/(d + 2)$$

But since $d \geq q$, $d + 2 > q - 1$ so $d + 2 \nmid q - 1$. Furthermore, $q - 1$ is even so $2(q - 1)$ is even, but $d + 2$ is odd, so $d + 2 \nmid 2(q - 1)$, so $2(q - 1)/(d + 2)$ is not an integer, thus our term on the right is not an integer. But then $d + 2 \nmid n + 2$. Thus odd composite integers do not satisfy our property either.

Finally, consider the only remaining possibility, when n is an odd prime number. The only such divisors of this is n and 1. $n + 2 \mid n + 2$ trivially. If and only if $1 + 2 = 3 \nmid n + 2$, we have our desired property then. So if n is prime and of the form $n = 3j - 2$ for some $j \in \mathbb{N}$, then n must satisfy our property. We have shown that no other such n can satisfy our property, thus this is all the possible solutions. \square

2 Problem 2

Find all positive integers m and n such that

$$2^m - 3^n = 7$$

Solution. We can rearrange our equation to get

$$2^m = 7 + 3^n \tag{1}$$

Obviously, any m that satisfies the above equation will also satisfy $2^m = 2 \cdot 2^{m-1} \equiv 7 \pmod{3}$ (since $3 \mid 3^n$ for any n). That is to say, the set of solutions M_1 to equation (1) (elements in M_1 are of the form 2^m) is a subset of the set of solutions M_2 to our subsequent relation, $M_1 \subset M_2$. If X is the set of solutions $x \in \mathbb{Z}$ to $2x \equiv 7 \pmod{3}$, then clearly $M_2 \subset X$.

Recall proposition 7.2 (B) from the course notes: if $a, b, m \in \mathbb{Z}$ with $m \neq 0$ and $d = \gcd(a, m)$, then if $d \mid b$, the congruence equation $ax \equiv b \pmod{m}$ has exactly d solutions. Since 2 and 3 are coprime, we have that $d = 1$, so $d \mid 7$, thus $2x \equiv 7 \pmod{3}$ has exactly one solution. Thus, X has exactly one element. Therefore, since $M_1 \subset X$, equation (1) has at most one solution.

We can verify that there does exist such a solution, namely when $m = 4$ and $n = 2$, then we have $2^4 - 3^2 = 16 - 9 = 7$. \square

3 Problem 3

Let $k \in \mathbb{N}$. Show that there exists k consecutive positive integers with the property that no integer from this set is of the form $a^2 + b^2$ for some $a, b \in \mathbb{Z}$.

Solution. Let $k \in \mathbb{N}$ be arbitrary. Let m be the product of the squares of the first k primes q of the form $q = 4j + 3$ (for some $j \in \mathbb{N}$), i.e. $m = \prod_{i=1}^k q_i^2$.

Note that we can always k -many q_i . To prove this, for the sake of contradiction, assume there are only $r < k$ many such primes of this form, q_1, q_2, \dots, q_r (where $q_1 < q_2 < \dots < q_r$). Note that $n = 4q_1q_2 \dots q_r - 1$ is of the form $4j + 3$, but $n > q_r$, so by assumption, n cannot be prime, so is composite. Further, note that none of the q_i and 2 divide n , thus all the primes in the prime factor decomposition of n is of the form $4j + 1$, thus $n \equiv 1 \pmod{4}$ which is a contradiction. Thus n is a prime greater than q_r and $n = 4j - 1$. We can do this indefinitely to get k many primes of the form $4j + 3$.

Now, consider the system

$$\begin{aligned} x &\equiv q_1 - 1 \pmod{q_1^2} \\ x &\equiv q_2 - 2 \pmod{q_2^2} \\ &\vdots \\ x &\equiv q_k - k \pmod{q_k^2} \end{aligned}$$

By the Chinese Remainder Theorem, there exists a unique solution to the system modulo m , which we'll call x_0 . Then, let $x_i = x_0 + i$ for $1 \leq i \leq k$. Note that $x_i = q_i + nq_i^2$, thus $q_i \mid x_i$ but $q_i^2 \nmid x_i$ (and so $q_i^s \nmid x_i$ for all $s \geq 2$). Thus, there are k consecutive integers x_1, x_2, \dots, x_k whose prime number decomposition that contain a prime of the form $4j + 3$ with exponent 1. But by Theorem 13.4 from the notes, for all $a, b \in \mathbb{Z}$, $a^2 + b^2 \neq x_i$ for all $1 \leq i \leq k$, since the exponent of q_i is $\gamma_i = 1$, which is not even. \square

4 Problem 4

As always, for each positive integer m , we have that $d(m)$ is the number of the positive divisors of m ; also, we let $\phi(m)$ be the corresponding value of the Euler ϕ -function. Then compute the following limits:

$$\lim_{n \rightarrow \infty} \frac{n!}{d(n!)\phi(n!)}$$

$$\lim_{n \rightarrow \infty} \frac{n!}{2^{d(n!)}}$$

Solution.

Limit 1: Note that the prime factor decomposition of $n!$ contains every prime that came before it, since if p is prime and $p < n$, $p \mid n!$ by definition of factorial. Thus, if we enumerate the primes in order (ie. $p_1 = 2$, $p_2 = 3$, etc.), let p_r be the greatest prime less than or equal to n . Then we can write $n! = \prod_{i=1}^r p_i^{\alpha_i}$ thus

$$\frac{n!}{d(n!)\phi(n!)} = \prod_{i=1}^r \frac{p_i^{\alpha_i}}{d(p_i^{\alpha_i})\phi(p_i^{\alpha_i})} = \prod_{i=1}^r \frac{p_i^{\alpha_i}}{(\alpha_i + 1)p_i^{\alpha_i-1}(p_i - 1)} = \prod_{i=1}^r \frac{p_i}{(\alpha_i + 1)(p_i - 1)}$$

Note that if n is an integer such that $n > 3$, we have $3n - 2n > 3 \implies 3(n - 1) > 2n \implies \frac{n}{n-1} < \frac{3}{2}$. Thus for $i \geq 3$, $p_i \geq 5$, and since $\alpha_i \geq 1$, we have

$$\frac{p_i}{(\alpha_i + 1)(p_i - 1)} \leq \frac{p_i}{2(p_i - 1)} < \frac{3}{4}$$

for all $i \geq 3$.

If $\pi(n)$ denotes the prime counting function (the number of prime numbers less than n), we have that

$$\frac{n!}{d(n!)\phi(n!)} \leq \frac{p_1}{(\alpha_1 + 1)(p_1 - 1)} \frac{p_2}{(\alpha_2 + 1)(p_2 - 1)} \left(\frac{3}{4}\right)^{r-2} = \frac{p_1}{(\alpha_1 + 1)(p_1 - 1)} \frac{p_1}{(\alpha_1 + 1)(p_1 - 1)} \left(\frac{3}{4}\right)^{\pi(n)-2}$$

But since there are infinitely many primes, $\pi(n)$ approaches infinity as n does, thus $\left(\frac{3}{4}\right)^{\pi(n)-2} \rightarrow 0$ as $n \rightarrow \infty$. But since $\frac{n!}{d(n!)\phi(n!)}$ is bounded below by zero, by squeeze theorem, we have that

$$\lim_{n \rightarrow \infty} \frac{n!}{d(n!)\phi(n!)} = 0$$

as well.

Limit 2: Let $a_n = \frac{n!}{2^{d(n!)}}$. Then

$$\frac{a_{n+1}}{a_n} = \frac{(n+1)!/2^{d((n+1)!)}}{n!2^{d(n!)}} = (n+1)/2^{d(n!)-d((n+1)!)}$$

Thus, it is sufficient to show that $2^{d((n+1)!)-d(n!)} > 2n+2$ for sufficiently large n , because then $(n+1)/2^{d((n+1)!)-d(n!)} < (n+1)/(2n+2) = \frac{1}{2}$, so $a_n < C \left(\frac{1}{2}\right)^n$ (where C is a constant). Since $C \left(\frac{1}{2}\right)^n \rightarrow 0$ as $n \rightarrow \infty$, and $a_n > 0$ for all n , by squeeze theorem, we would have $\lim_{n \rightarrow \infty} a_n = 0$.

We show first show that $d((n+1)!)-d(n!) \geq 2^{\pi(n)-1}$ (where $\pi(n)$ is number of primes $\leq n$). To see this, let $n! = \prod_{i=1}^{\pi(n)} p_i^{\alpha_i}$ be the unique prime factorization of $n!$, where p_i is the i th prime number (e.g. $p_1 = 2$). We deal with two case: when $n+1$ is prime, and when it is composite. First, let $n+1$ be prime. Then $(n+1)! = p_{\pi(n)+1} \prod_{i=1}^{\pi(n)} p_i^{\alpha_i}$. Then let $\beta_i \in \{0, 1\}$, we have that $p_{\pi(n)+1} \prod_{i=1}^{\pi(n)} p_i^{\beta_i} \mid p_{\pi(n)+1} = n+1 \mid (n+1)!$, but since $p_{n+1} \nmid n!$, we have $p_{\pi(n)+1} \prod_{i=1}^{\pi(n)} p_i^{\beta_i} \nmid n!$. There are a total of two possibilities for $\pi(n)$ many β_i , thus there are $2^{\pi(n)}$ ways to choose the β_i , so there are at least $2^{\pi(n)}$ many divisors of $(n+1)!$ but not of $n!$. Thus

$$d((n+1)!)-d(n!) \geq 2^{\pi(n)} \geq 2^{\pi(n)-1}$$

which proves the $n + 1$ is prime case. Now let $n + 1$ be composite. Then we have the unique prime factorization $(n + 1)! = \prod_{i=1}^{\pi(n)} p_i^{\beta_i}$ where there is at least one i such that $\gamma_i > \alpha_i$ (by unique factorization). Denote this γ_i by γ_j (and p_j). Then using $\beta_i \in \{0, 1\}, i \neq j$ as before, let

$$m = p_j^{\gamma_j} \prod_{\substack{i=1 \\ i \neq j}}^{\pi(n)} p_i^{\beta_i}$$

Similar to before, we have $m \mid (n + 1)!$, but $m \nmid n!$ since $p_j^{\gamma_j} \nmid n!$. There are $\pi(n) - 1$ ways to choose the β_i (similar to the prime case), so there are at least $2^{\pi(n)-1}$ many divisors of $(n + 1)!$ but not of $n!$. Thus

$$d((n + 1)!) - d(n!) \geq 2^{\pi(n)-1}$$

Thus, we have that $2^{d((n+1)!)-d(n!)} \geq 2^{2^{\pi(n)-1}}$.

Note that for all $k \in \mathbb{N}$ such that $k \geq 5$, $2^{k-1} > 2k + 2$ (since $2^{5-1} > 2 \cdot 5 + 2$, and the exponential grows faster than linear thereafter). Since $\pi(n) \geq 5$ when $n \geq 11$, we have $2^{2^{\pi(n)-1}} > 2^{2\pi(n)+2}$ when $n \geq 11$.

Finally, we show that $2^{2\pi(n)} > n$. Consider all $m \in \mathbb{N}$ such that $m \leq n$. We can rewrite a unique prime factorization $m = p_1^{\eta_1} p_2^{\eta_2} \cdots p_{\pi(n)}^{\eta_{\pi(n)}}$ where η_i may be 0. By the division algorithm, we have $\eta_i = 2q_i + r_i$ where $r_i \in \{0, 1\}$ and r_i, q_i are unique. Thus we can uniquely write $m = u^2 p_1^{r_1} \cdots p_{\pi(n)}^{r_{\pi(n)}}$ where $u \in \mathbb{N}$ such that $u = p_1^{q_1} \cdots p_{\pi(n)}^{q_{\pi(n)}}$. See that $u \leq \sqrt{m}$. We use the notation $\mathbb{N}_j = \{i : i \in \mathbb{N}, i \leq j\}$. Now define a map

$$\phi : \mathbb{N}_n \rightarrow \mathbb{N}_{\lfloor \sqrt{n} \rfloor} \times \mathcal{P}(\{p_i : i \in \mathbb{N}_{\pi(n)}\})$$

with $m \mapsto (u, \{p_i : i \in \mathbb{N}_{\pi(n)}, r_i = 1\})$. By uniqueness, this map is injective. Thus we have that $|\mathbb{N}_n| \leq |\mathbb{N}_{\lfloor \sqrt{n} \rfloor}| |\mathcal{P}(\{p_i : i \in \mathbb{N}_{\pi(n)}\})|$, i.e. $n \leq 2^{\pi(n)} \sqrt{n}$. We want to show the strict inequality. We now deal with when n is a perfect square, and when n is not. If n is not a perfect square, we have that

$$n \leq \lfloor \sqrt{n} \rfloor 2^{\pi(n)} < 2^{\pi(n)} \sqrt{n}$$

Now let n be a perfect square. Note the pre-image of $(\sqrt{n}, \{p_i : i \in \mathbb{N}_{\pi(n)}, r_i = 1\})$ is the m of the form $m = np_1 p_2 \cdots p_{\pi(n)}$, but then $m > n$, which is a contradiction. But then the pre-image is empty, thus $n \neq 2^{\pi(n)} \sqrt{n}$ giving us the strict inequality. Thus, regardless of n , we have $n \neq 2^{\pi(n)} \sqrt{n}$, and squaring both sides (both are positive) gives us $2^{2\pi(n)} > n$.

Finally, we have that

$$2^{d((n+1)!)-d(n!)} > 2^{2\pi(n)+2} > 2 \cdot 2^{2\pi(n)} + 2 > 2n + 2$$

for $n \geq 11$, as desired. Thus we have shown the desired limit as observed before. \square