

Math 321 Assignment 1
Due Friday, January 19 at 9 am

Instructions

- (i) Homework should be submitted using Canvas. Include your name and SID.
 - (ii) Solutions should be well-crafted, legible and written in complete English sentences. You will be graded both on accuracy as well as the quality of exposition.
 - (iii) Theorems stated in the text or proved in lecture do not need to be reproved. Any other statement should be justified rigorously.
-

In this homework, we will need several definitions. Let $I = [a, b]$ be an interval and $k \geq 0$ be an integer. If $f: I \rightarrow \mathbb{R}$ is a function that is k -times differentiable on I , then we define

$$\|f\|_{C^k(I)} = \sum_{j=0}^k \sup_{x \in I} |f^{(j)}(x)|.$$

This quantity is called the “ C^k norm of f .” We define $C^k(I)$ to be the set of functions $f: I \rightarrow \mathbb{R}$ that satisfy the following two properties. (i): f is k -times differentiable on I , and (ii): $f^{(k)}$ is continuous on I . We define a metric on $C^k(I)$ as follows: $d(f, g) = \|f - g\|_{C^k(I)}$, i.e.

$$d(f, g) = \sum_{j=0}^k \sup_{x \in I} |f^{(j)}(x) - g^{(j)}(x)|. \quad (1)$$

It is straightforward to verify that this is indeed a metric, but you do not have to do so for this homework.

1. Let $f(t) = e^t$; recall that f is monotone increasing, $f'(t) = f(t)$, and $f(0) = 1$. Let $P_n(t)$ be the n -th order Taylor polynomial of f at the point $x_0 = 0$, as discussed in lecture. Let $I = [-1, 1]$ and let $k \geq 1$ be an integer.

Using Taylor’s theorem, prove that the sequence $\{P_n\}$ converges to f in the metric space $C^k(I)$.

Hints (i) Compute the Taylor polynomial $P_n(t)$. (ii) What is the derivative of P_n ? (iii) What are the higher derivatives of P_n ? (iv) How can you estimate each term in (1)?

2. Let $f(t) = e^t$. Let $P_n(t)$ be the n -th order Taylor polynomial of f at the point $x_0 = 0$.

a) Let $n \geq 1$. Prove that $n!P_n(1)$ is an integer.

b) Using part a) and Taylor’s theorem, prove that Euler’s number e is irrational. You may use the fact that e^t is strictly monotone increasing, and $0 < e < 3$.

Hint: if e were rational, then we could write $e = m/n$

The next problem concerns monotone increasing functions, and will help prepare us for the Riemann–Stieltjes integral. Let $\alpha: [0, 1] \rightarrow \mathbb{R}$ be increasing. Recall from last term that for every $c \in [0, 1]$, $\lim_{x \searrow c} \alpha(x)$ and $\lim_{x \nearrow c} \alpha(x)$ always exist. Thus α is continuous at c if and only if $\lim_{x \searrow c} \alpha(x) = \lim_{x \nearrow c} \alpha(x)$. If α is not continuous at c , then $\lim_{x \nearrow c} \alpha(x) < \lim_{x \searrow c} \alpha(x)$, and we say α has a *jump discontinuity* at c .

3. Let $\alpha: [0, 1] \rightarrow \mathbb{R}$ be monotone increasing. Prove that the set of points $c \in [0, 1]$ where α is not continuous is either finite (possibly empty), or countably infinite.