

C H A P T E R 6

Linear Transformations

Section 6.1	Introduction to Linear Transformations	207
Section 6.2	The Kernel and Range of a Linear Transformation	212
Section 6.3	Matrices for Linear Transformations.....	217
Section 6.4	Transition Matrices and Similarity	224
Section 6.5	Applications of Linear Transformations	228
Review Exercises	233
Project Solutions	241

CHAPTER 6

Linear Transformations

Section 6.1 Introduction to Linear Transformations

2. (a) The image of \mathbf{v} is

$$\begin{aligned} T(0, 4) &= (0, 2(4) - 0, 4) \\ &= (0, 8, 4). \end{aligned}$$

- (b) If $T(v_1, v_2) = (v_1, 2v_2 - v_1, v_2) = (2, 4, 3)$, then

$$\begin{aligned} v_1 &= 2 \\ 2v_2 - v_1 &= 4 \\ v_2 &= 3 \end{aligned}$$

which implies that $v_1 = 2$ and $v_2 = 3$. So, the preimage of \mathbf{w} is $(2, 3)$.

4. (a) The image of \mathbf{v} is

$$T(2, 3, 0) = (3 - 2, 2 + 3, 2(2)) = (1, 5, 4).$$

- (b) If $T(v_1, v_2, v_3) = (v_2 - v_1, v_1 + v_2, 2v_1) = (-11, -1, 10)$,

then

$$\begin{aligned} v_2 - v_1 &= -11 \\ v_1 + v_2 &= -1 \\ 2v_1 &= 10 \end{aligned}$$

which implies that $v_1 = 5$ and $v_2 = -6$. So, the preimage of \mathbf{w} is $\{(5, -6, t) : t \text{ is any real number}\}$.

6. (a) The image of \mathbf{v} is

$$T(2, 1, 4) = (2(2) + 1, 2 - 1) = (5, 1).$$

- (b) If $T(v_1, v_2, v_3) = (2v_1 + v_2, v_1 - v_2) = (-1, 2)$, then

$$\begin{aligned} 2v_1 + v_2 &= -1 \\ v_1 - v_2 &= 2, \end{aligned}$$

which implies that $v_1 = \frac{1}{3}$, $v_2 = -\frac{5}{3}$, and $v_3 = t$, where t is any real number. So, the preimage of \mathbf{w} is $\{(\frac{1}{3}, -\frac{5}{3}, t) : t \text{ is any real number}\}$.

16. T preserves addition.

$$\begin{aligned} T(A_1) + T(A_2) &= T\left(\begin{bmatrix} a_1 & b_1 \\ c_1 & d_1 \end{bmatrix}\right) + T\left(\begin{bmatrix} a_2 & b_2 \\ c_2 & d_2 \end{bmatrix}\right) \\ &= a_1 + b_1 + c_1 + d_1 + a_2 + b_2 + c_2 + d_2 \\ &= (a_1 + a_2) + (b_1 + b_2) + (c_1 + c_2) + (d_1 + d_2) = T(A_1 + A_2) \end{aligned}$$

T preserves scalar multiplication.

$$T(kA) = ka + kb + kc + kd = k(a + b + c + d) = kT(A)$$

Therefore, T is a linear transformation.

8. (a) The image of \mathbf{v} is

$$\begin{aligned} T(2, 4) &= \left(\frac{\sqrt{3}}{2}(2) - \frac{1}{2}(4), 2 - 4, 4\right) \\ &= (\sqrt{3} - 2, -2, 4). \end{aligned}$$

- (b) If $T(v_1, v_2) = \left(\frac{\sqrt{3}}{2}v_1 - \frac{1}{2}v_2, v_1 - v_2, v_2\right) = (\sqrt{3}, 2, 0)$,

then

$$\begin{aligned} \frac{\sqrt{3}}{2}v_1 - \frac{1}{2}v_2 &= \sqrt{3} \\ v_1 - v_2 &= 2 \\ v_2 &= 0 \end{aligned}$$

which implies that $v_1 = 2$ and $v_2 = 0$. So, the preimage of \mathbf{w} is $(2, 0)$.

10. T is *not* a linear transformation because it does not preserve addition nor scalar multiplication.

For example,

$$\begin{aligned} T(1, 1) + T(1, 1) &= (1, 1) + (1, 1) \\ &= (2, 2) \neq (2, 4) = T(2, 2). \end{aligned}$$

12. T is *not* a linear transformation because it does not preserve addition. For example,

$$\begin{aligned} T(1, 1, 1) + T(1, 1, 1) &= (2, 2, 2) + (2, 2, 2) \\ &= (4, 4, 4) \\ &\neq (3, 3, 3) = T(2, 2, 2). \end{aligned}$$

14. T is *not* a linear transformation because it does not preserve addition nor scalar multiplication. For example,

$$\begin{aligned} T(1, 1) + T(1, 1) &= (1, 1, 1) + (1, 1, 1) \\ &= (2, 2, 2) \neq (4, 4, 4) = T(2, 2). \end{aligned}$$

18. T is not a linear transformation. T does not preserve addition.

$$T(A_1) + T(A_2) = T\begin{bmatrix} a_1 & b_1 \\ c_1 & d_1 \end{bmatrix} + T\begin{bmatrix} a_2 & b_2 \\ c_2 & d_2 \end{bmatrix} = b_1^2 + b_2^2 \neq (b_1 + b_2)^2 = T(A_1 + A_2)$$

20. Let A and B be two elements of $M_{3,3}$ (two 3×3 matrices) and let c be a scalar. First

$$T(A + B) = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -10 \end{bmatrix} (A + B) = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -10 \end{bmatrix} A + \begin{bmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -10 \end{bmatrix} B = T(A) + T(B)$$

by Theorem 2.3, part 2 and

$$T(cA) = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -10 \end{bmatrix} (cA) = c \begin{bmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -10 \end{bmatrix} A = cT(A)$$

by Theorem 2.3, part 4. So, T is a linear transformation.

22. T preserves addition.

$$\begin{aligned} T(a_0 + a_1x + a_2x^2) + T(b_0 + b_1x + b_2x^2) &= (a_1 + 2a_2x) + (b_1 + 2b_2x) \\ &= (a_1 + b_1) + 2(a_2 + b_2)x \\ &= T((a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2) \end{aligned}$$

T preserves scalar multiplication.

$$T(c(a_0 + a_1x + a_2x^2)) = T(ca_0 + ca_1x + ca_2x^2) = ca_1 + 2ca_2x = c(a_1 + 2a_2x) = cT(a_0 + a_1x + a_2x^2)$$

Therefore, T is a linear transformation.

24. Because $(2, 0) = \frac{2}{3}(1, 2) - \frac{4}{3}(-1, 1)$, you have

$$\begin{aligned} T(2, 0) &= T\left[\frac{2}{3}(1, 2) - \frac{4}{3}(-1, 1)\right] \\ &= \frac{2}{3}T(1, 2) - \frac{4}{3}T(-1, 1) \\ &= \frac{2}{3}(1, 0) - \frac{4}{3}(0, 1) \\ &= \left(\frac{2}{3}, -\frac{4}{3}\right) \end{aligned}$$

Similarly, $(0, 3) = (1, 2) + (-1, 1)$, which gives

$$\begin{aligned} T(0, 3) &= T[(1, 2) + (-1, 1)] \\ &= T(1, 2) + T(-1, 1) \\ &= (1, 0) + (0, 1) \\ &= (1, 1). \end{aligned}$$

26. Because $(2, -1, 0)$ can be written as

$$(2, -1, 0) = 2(1, 0, 0) - 1(0, 1, 0) + 0(0, 0, 1),$$

you can use Property 4 of Theorem 6.1 to write

$$\begin{aligned} T(2, -1, 0) &= 2T(1, 0, 0) - T(0, 1, 0) + 0T(0, 0, 1) \\ &= 2(2, 4, -1) - (1, 3, -2) + (0, 0, 0) \\ &= (3, 5, 0). \end{aligned}$$

28. Because $(-2, 4, -1)$ can be written as

$$(-2, 4, -1) = -2(1, 0, 0) + 4(0, 1, 0) - 1(0, 0, 1),$$

you can use Property 4 of Theorem 6.1 to write

$$\begin{aligned} T(-2, 4, -1) &= -2T(1, 0, 0) + 4T(0, 1, 0) - T(0, 0, 1) \\ &= -2(2, 4, -1) + 4(1, 3, -2) - (0, -2, 2) \\ &= (0, 6, -8). \end{aligned}$$

30. Because $(0, 2, -1)$ can be written as

$$(0, 2, -1) = \frac{3}{2}(1, 1, 1) - \frac{1}{2}(0, -1, 2) - \frac{3}{2}(1, 0, 1),$$

you can use Property 4 of Theorem 6.1 to write

$$\begin{aligned} T(0, 2, -1) &= \frac{3}{2}T(1, 1, 1) - \frac{1}{2}T(0, -1, 2) - \frac{3}{2}T(1, 0, 1) \\ &= \frac{3}{2}(2, 0, -1) - \frac{1}{2}(-3, 2, -1) - \frac{3}{2}(1, 1, 0) \\ &= \left(3, -\frac{5}{2}, -1\right). \end{aligned}$$

32. Because $(-2, 1, 0)$ can be written as

$$(-2, 1, 0) = 2(1, 1, 1) + (0, -1, 2) - 4(1, 0, 1),$$

you can use Property 4 of Theorem 6.1 to write

$$\begin{aligned} T(-2, 1, 0) &= 2T(1, 1, 1) + T(0, -1, 2) - 4T(1, 0, 1) \\ &= 2(2, 0, -1) + (-3, 2, -1) - 4(1, 1, 0) \\ &= (-3, -2, -3). \end{aligned}$$

34. Because the matrix has 2 columns, the dimension of R^n is 2. Because the matrix has 3 rows, the dimension of R^m is 3. So, $T: R^2 \rightarrow R^3$.

36. Because the matrix has five columns, the dimension of R^n is 5. Because the matrix has two rows, the dimension of R^m is 2. So, $T: R^5 \rightarrow R^2$.

38. Because the matrix has five columns, the dimension of R^n is 5. Because the matrix has three rows, the dimension of R^m is 3. So, $T: R^5 \rightarrow R^3$.

$$40. (a) \quad T(2, 4) = \begin{bmatrix} 1 & 2 \\ -2 & 4 \\ -2 & 2 \end{bmatrix} \begin{bmatrix} 2 \\ 4 \end{bmatrix} = \begin{bmatrix} 10 \\ 12 \\ 4 \end{bmatrix} = (10, 12, 4)$$

(b) The preimage of $(-1, 2, 2)$ is given by solving the equation

$$T(v_1, v_2) = \begin{bmatrix} 1 & 2 \\ -2 & 4 \\ -2 & 2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix}$$

for $\mathbf{v} = (v_1, v_2)$. The equivalent system of linear equations

$$\begin{aligned} v_1 + 2v_2 &= -1 \\ -2v_1 + 4v_2 &= 2 \\ -2v_1 + 2v_2 &= 2 \end{aligned}$$

has the solution $v_1 = -1$ and $v_2 = 0$. So, $(-1, 0)$ is the preimage of $(-1, 2, 2)$ under T .

(c) Because the system of linear equations represented by the equation

$$\begin{bmatrix} 1 & 2 \\ -2 & 4 \\ -2 & 2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

has no solution, $(1, 1, 1)$ has no preimage under T .

$$42. (a) \quad T(1, 0, -1, 3, 0) = \begin{bmatrix} -1 & 2 & 1 & 3 & 4 \\ 0 & 0 & 2 & -1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -1 \\ 3 \\ 0 \end{bmatrix} = \begin{bmatrix} 7 \\ -5 \end{bmatrix} = (7, -5).$$

$$(b) \quad \text{The preimage of } (-1, 8) \text{ is determined by solving the equation } T(v_1, v_2, v_3, v_4, v_5) = \begin{bmatrix} -1 & 2 & 1 & 3 & 4 \\ 0 & 0 & 2 & -1 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \\ v_5 \end{bmatrix} = \begin{bmatrix} -1 \\ 8 \end{bmatrix}.$$

The equivalent system of linear equations has the solution $v_1 = 5 + 2r + \frac{7}{2}s + 4t$, $v_2 = r$, $v_3 = 4 + \frac{1}{2}s$, $v_4 = s$, and $v_5 = t$, where r , s , and t are any real numbers. So, the preimage is given by the set of vectors

$$\left\{ \left(5 + 2r + \frac{7}{2}s + 4t, r, 4 + \frac{1}{2}s, s, t \right) : r, s, t \text{ are real numbers} \right\}.$$

$$44. (a) \quad T(0, 1, 0, 1, 0) = \begin{bmatrix} 0 & 2 & 0 & 2 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 1 & 2 & 2 & 2 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \\ 0 \end{bmatrix} = (4, 0, 4)$$

(b) The preimage of $(0, 0, 0)$ is determined by solving the equation as shown.

$$T(v_1, v_2, v_3, v_4, v_5) = \begin{bmatrix} 0 & 2 & 0 & 2 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 1 & 2 & 2 & 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \\ v_5 \end{bmatrix} = (0, 0, 0)$$

The equivalent system of linear equations has the solution $v_1 = -t$, $v_2 = -s$, $v_3 = 0$, $v_4 = s$, and $v_5 = t$, where s and t are any real numbers. So, the preimage is given by the set of vectors $\{(-t, -s, 0, s, t)\}$.

(c) The preimage of $(1, -1, 2)$ is determined by solving the equation as shown.

$$T(v_1, v_2, v_3, v_4, v_5) = \begin{bmatrix} 0 & 2 & 0 & 2 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 1 & 2 & 2 & 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \\ v_5 \end{bmatrix} = (1, -1, 2)$$

The equivalent system of linear equations has the solution $v_1 = -3 - t$, $v_2 = \frac{1}{2} - s$, $v_3 = 2$, $v_4 = s$, and $v_5 = t$, where s and t are real numbers. So, the preimage is given by the set of vectors $\{(-3 - t, \frac{1}{2} - s, 2, s, t)\}$.

46. If $\theta = 45^\circ$, then T is given by

$$\begin{aligned} T(x, y) &= (x \cos \theta - y \sin \theta, x \sin \theta + y \cos \theta) \\ &= \left(\frac{\sqrt{2}}{2}x - \frac{\sqrt{2}}{2}y, \frac{\sqrt{2}}{2}x + \frac{\sqrt{2}}{2}y \right). \end{aligned}$$

Solving $T(x, y) = \mathbf{v} = (1, 1)$, you have

$$\frac{\sqrt{2}}{2}x - \frac{\sqrt{2}}{2}y = 1 \quad \text{and} \quad \frac{\sqrt{2}}{2}x + \frac{\sqrt{2}}{2}y = 1.$$

So, $x = \sqrt{2}$ and $y = 0$, and the preimage of \mathbf{v} is $(\sqrt{2}, 0)$.

$$48. \quad \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \begin{bmatrix} 12 \\ 5 \end{bmatrix} = \begin{bmatrix} 13 \\ 0 \end{bmatrix}$$

You then obtain the following system of equations.

$$12a - 5b = 13$$

$$12b + 5a = 0$$

Solving the second equation for a gives $a = \frac{-12}{5}b$.

Substituting this back into the first equation produces

$$12\left(\frac{-12}{5}b\right) - 5b = 13$$

$$\frac{-144}{5}b - 5b = 13$$

$$\frac{-169}{5}b = 13$$

$$b = \frac{-5}{13}.$$

Substituting $b = \frac{-5}{13}$ into $a = \frac{-12}{5}b$ you obtain

$$a = \frac{12}{13}.$$

50. If $\mathbf{v} = (x, y, z)$ is a vector into R^3 , then

$T(\mathbf{v}) = (0, y, z)$. In other words, T maps every vector in R^3 to its orthogonal projection in the yz -plane.

54. T is not a linear transformation.

Consider $A = I_n$. Then $T(A) = 1$, but $T(2A) = 2^n \neq 2T(A)$.

$$56. T\left(\begin{bmatrix} 1 & 3 \\ -1 & 4 \end{bmatrix}\right) = T\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + 3T\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} - T\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + 4T\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix} + 3\begin{bmatrix} 0 & 2 \\ 1 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} + 4\begin{bmatrix} 3 & -1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 12 & -1 \\ 7 & 4 \end{bmatrix}$$

58. This statement is true because D_x is a linear transformation and therefore preserves addition and scalar multiplication.

60. This statement is false because $\cos \frac{x}{2} \neq \frac{1}{2} \cos x$ for all x .

62. If $D_x(g(x)) = e^x$, then $g(x) = e^x + C$.

64. If $D_x(g(x)) = \frac{1}{x}$, then $g(x) = \ln x + C$.

66. Solve the equation $\int_0^1 p(x)dx = 1$ for $p(x)$ in P_2 .

$$\int_0^1 (a_0 + a_1x + a_2x^2)dx = 1 \Rightarrow \left[a_0x + a_1\frac{x^2}{2} + a_2\frac{x^3}{3} \right]_0^1 = 1 \Rightarrow a_0 + \frac{1}{2}a_1 + \frac{1}{3}a_2 = 1.$$

Letting $a_2 = -3b$ and $a_1 = -2a$ be free variables, $a_0 = 1 + a + b$, and $p(x) = (1 + a + b) - 2ax - 3bx^2$.

68. (a) False. This function does not preserve addition nor scalar multiplication. For example,
 $f(3x) = 27x^3 \neq 3f(x)$.

- (b) False. If $f: R \rightarrow R$ is given by $f(x) = ax + b$ for some $a, b \in R$, then it preserves addition and scalar multiplication if and only if $b = 0$.

70. (a) $T(x, y) = T[x(1, 0) + y(0, 1)]$
 $= xT(1, 0) + yT(0, 1)$
 $= x(0, 1) + y(1, 0) = (y, x)$

- (b) T is a reflection about the line $y = x$.

52. T is a linear transformation.

T preserves addition.

$$\begin{aligned} T(A + B) &= (A + B)X - X(A + B) \\ &= AX + BX - XA - XB \\ &= (AX - XA) + (BX - XB) \\ &= T(A) + T(B) \end{aligned}$$

T preserves scalar multiplication.

$$\begin{aligned} T(cA) &= (cA)X - X(cA) \\ &= c(AX) - c(XA) \\ &= c(AX - XA) \\ &= cT(A) \end{aligned}$$

72. Use the result of Exercise 71(a) as follows.

$$T(3, 4) = \left(\frac{3+4}{2}, \frac{3+4}{2} \right) = \left(\frac{7}{2}, \frac{7}{2} \right)$$

$$\begin{aligned} T(T(3, 4)) &= T\left(\frac{7}{2}, \frac{7}{2}\right) \\ &= \left(\frac{1}{2}\left(\frac{7}{2} + \frac{7}{2}\right), \frac{1}{2}\left(\frac{7}{2} + \frac{7}{2}\right) \right) = \left(\frac{7}{2}, \frac{7}{2} \right) \end{aligned}$$

T is projection onto the line $y = x$.

74. To show that $T: V \rightarrow W$ is a linear transformation, show that $T: V \rightarrow W$ preserves addition and scalar multiplication by using the definition:

$$(1) T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$$

and

$$(2) T(c\mathbf{u}) = cT(\mathbf{u}),$$

where c is any nonzero constant.

76. (a) Because $T(0, 0) = (-h, -k) \neq (0, 0)$, a translation cannot be a linear transformation.

$$\begin{aligned} \text{(b)} \quad T(0, 0) &= (0 - 2, 0 + 1) = (-2, 1) \\ T(2, -1) &= (2 - 2, -1 + 1) = (0, 0) \\ T(5, 4) &= (5 - 2, 4 + 1) = (3, 5) \end{aligned}$$

(c) Because $T(x, y) = (x - h, y - k) = (x, y)$ implies $x - h = x$ and $y - k = y$, a translation has no fixed points.

78. There are many possible examples. For instance, let $T: R^3 \rightarrow R^3$ be given by $T(x, y, z) = (0, 0, 0)$. Then if $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is any set of linearly independent vectors, their images $T(\mathbf{v}_1), T(\mathbf{v}_2), T(\mathbf{v}_3)$ form a dependent set.

80. Let T be defined by $T(\mathbf{v}) = \langle \mathbf{v}, \mathbf{v}_0 \rangle$. Then because

$$\begin{aligned} T(\mathbf{v} + \mathbf{w}) &= \langle \mathbf{v} + \mathbf{w}, \mathbf{v}_0 \rangle = \langle \mathbf{v}, \mathbf{v}_0 \rangle + \langle \mathbf{w}, \mathbf{v}_0 \rangle = T(\mathbf{v}) + T(\mathbf{w}) \\ \text{and } T(c\mathbf{v}) &= \langle c\mathbf{v}, \mathbf{v}_0 \rangle = c\langle \mathbf{v}, \mathbf{v}_0 \rangle = cT(\mathbf{v}), T \text{ is a linear transformation.} \end{aligned}$$

82. Because

$$\begin{aligned} T(\mathbf{u} + \mathbf{v}) &= \langle \mathbf{u} + \mathbf{v}, \mathbf{w}_1 \rangle \mathbf{w}_1 + \cdots + \langle \mathbf{u} + \mathbf{v}, \mathbf{w}_n \rangle \mathbf{w}_n \\ &= (\langle \mathbf{u}, \mathbf{w}_1 \rangle \mathbf{w}_1 + \langle \mathbf{v}, \mathbf{w}_1 \rangle \mathbf{w}_1) + \cdots + (\langle \mathbf{u}, \mathbf{w}_n \rangle \mathbf{w}_n + \langle \mathbf{v}, \mathbf{w}_n \rangle \mathbf{w}_n) \\ &= (\langle \mathbf{u}, \mathbf{w}_1 \rangle \mathbf{w}_1 + \cdots + \langle \mathbf{u}, \mathbf{w}_n \rangle \mathbf{w}_n) + (\langle \mathbf{v}, \mathbf{w}_1 \rangle \mathbf{w}_1 + \cdots + \langle \mathbf{v}, \mathbf{w}_n \rangle \mathbf{w}_n) = T(\mathbf{u}) + T(\mathbf{v}) \end{aligned}$$

and

$$T(c\mathbf{u}) = \langle c\mathbf{u}, \mathbf{w}_1 \rangle \mathbf{w}_1 + \cdots + \langle c\mathbf{u}, \mathbf{w}_n \rangle \mathbf{w}_n = c\langle \mathbf{u}, \mathbf{w}_1 \rangle \mathbf{w}_1 + \cdots + c\langle \mathbf{u}, \mathbf{w}_n \rangle \mathbf{w}_n = c[\langle \mathbf{u}, \mathbf{w}_1 \rangle \mathbf{w}_1 + \cdots + \langle \mathbf{u}, \mathbf{w}_n \rangle \mathbf{w}_n] = cT(\mathbf{u}),$$

T is a linear transformation.

84. Suppose first that T is a linear transformation. Then $T(a\mathbf{u} + b\mathbf{v}) = T(a\mathbf{u}) + T(b\mathbf{v}) = aT(\mathbf{u}) + bT(\mathbf{v})$.

Second, suppose $T(a\mathbf{u} + b\mathbf{v}) = aT(\mathbf{u}) + bT(\mathbf{v})$. Then $T(\mathbf{u} + \mathbf{v}) = T(1\mathbf{u} + 1\mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$

and $T(c\mathbf{u}) = T(c\mathbf{u} + \mathbf{0}) = cT(\mathbf{u}) + T(\mathbf{0}) = cT(\mathbf{u})$.

Section 6.2 The Kernel and Range of a Linear Transformation

2. $T: R^3 \rightarrow R^3, T(x, y, z) = (x, 0, z)$

The kernel consists of all vectors lying on the y -axis.

That is, $\ker(T) = \{(0, y, 0) : y \text{ is a real number}\}$.

4. $T: R^3 \rightarrow R^3, T(x, y, z) = (-z, -y, -x)$

Solving the equation

$$T(x, y, z) = (-z, -y, -x) = (0, 0, 0) \text{ yields that trivial}$$

solution $x = y = z = 0$. So, $\ker(T) = \{(0, 0, 0)\}$.

6. $T: P_2 \rightarrow R, T(a_0 + a_1x + a_2x^2) = a_0$

Solving the equation $T(a_0 + a_1x + a_2x^2) = a_0 = 0$

yields solutions of the form $a_0 = 0$ and a_1 and a_2 are any real numbers. So,

$$\ker(T) = \{a_1x + a_2x^2 : a_1, a_2 \in R\}.$$

8. $T: P_3 \rightarrow P_2,$

$$T(a_0 + a_1x + a_2x^2 + a_3x^3) = a_1 + 2a_2x + 3a_3x^2$$

Solving the equation

$$T(a_0 + a_1 + a_2x^2 + a_3x^3) = a_1 + 2a_2x + 3a_3x^2 = 0$$

yields solutions of the form $a_1 = a_2 = a_3 = 0$ and a_0

any real number. So, $\ker(T) = \{a_0 : a_0 \in R\}$.

10. $T: R^2 \rightarrow R^2, T(x, y) = (x - y, y - x)$

Solving the equation

$$T(x, y) = (x - y, y - x) = (0, 0) \text{ yields solutions of}$$

the form $x = y$. So, $\ker(T) = \{(x, x) : x \in R\}$.

$$12. (a) \begin{bmatrix} 1 & 2 \\ -3 & -6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow \begin{aligned} x_1 + 2x_2 &= 0 \\ -3x_1 - 6x_2 &= 0 \end{aligned} \Rightarrow \begin{aligned} x_1 + 2x_2 &= 0 \\ 0 &= 0 \end{aligned}$$

Using the parameter $t = x_2$ produces the family of solutions

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} -2t \\ t \end{bmatrix} = t \begin{bmatrix} -2 \\ 1 \end{bmatrix}.$$

$$\text{So, } \ker(T) = \{t(-2, 1): t \text{ is a real number}\} \\ = \text{span}\{(-2, 1)\}.$$

(b) Transpose A and find the equivalent row-echelon form.

$$A^T = \begin{bmatrix} 1 & -3 \\ 2 & -6 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & -3 \\ 0 & 0 \end{bmatrix}$$

$$\text{So, } \text{range}(T) = \{t(1, -3): t \text{ is a real number}\} \\ = \text{span}\{(1, -3)\}.$$

16. (a) Because

$$T(\mathbf{x}) = \begin{bmatrix} 1 & 1 \\ -1 & 2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

has only the trivial solution $x_1 = x_2 = 0$, $\ker(T) = \{(0, 0)\}$.

(b) Transpose A and find the equivalent reduced row-echelon form.

$$A^T = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 2 & 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & \frac{1}{3} \\ 0 & 1 & \frac{1}{3} \end{bmatrix}$$

$$\text{So, } \text{range}(T) = \text{span}\left\{\left(1, 0, \frac{1}{3}\right), \left(0, 1, \frac{1}{3}\right)\right\}.$$

18. (a) Because

$$T(\mathbf{x}) = \begin{bmatrix} -1 & 3 & 2 & 1 & 4 \\ 2 & 3 & 5 & 0 & 0 \\ 2 & 1 & 2 & 1 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

has solutions of the form $(-10s - 4t, -15s - 24t, 13s + 16t, 9s, 9t)$,

$$\ker(T) = \text{span}\{(-10, -15, 13, 9, 0), (-4, -24, 16, 0, 9)\}.$$

(b) Transpose A and find the equivalent reduced row-echelon form.

$$A^T = \begin{bmatrix} -1 & 2 & 2 \\ 3 & 3 & 1 \\ 2 & 5 & 2 \\ 1 & 0 & 1 \\ 4 & 0 & 0 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{So, } \text{range}(T) = \text{span}\{(1, 0, 0), (0, 1, 0), (0, 0, 1)\} = R^3.$$

14. (a) Because

$$T(\mathbf{x}) = \begin{bmatrix} 1 & -2 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

has solutions of the form $(-2t, -\frac{1}{2}t, t)$ where t is any real number,

$$\ker(T) = \{t(-2, -\frac{1}{2}, 1): t \text{ is a real number}\} \\ = \text{span}\{(-2, -\frac{1}{2}, 1)\}.$$

$$\{(-2, -\frac{1}{2}, 1)\}.$$

(b) Transpose A and find the equivalent reduced row-echelon form

$$A^T = \begin{bmatrix} 1 & 0 \\ -2 & 2 \\ 1 & 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$$

$$\text{So, } \text{range}(T) \text{ is } \{(1, 0), (0, 1)\} = R^2.$$

20. (a) The kernel of T is given by the solution to the equation $T(\mathbf{x}) = \mathbf{0}$. So,

$$\ker(T) = \{(2t, -3t) : t \text{ is any real number}\}.$$

(b) $\text{nullity}(T) = \dim(\ker(T)) = 1$

- (c) Transpose T and find the equivalent reduced row-echelon form.

$$A^T = \begin{bmatrix} 3 & -9 \\ 2 & -6 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & -3 \\ 0 & 0 \end{bmatrix}$$

So, $\text{range}(T) = \{(t, -3t) : t \text{ is any real number}\}.$

(d) $\text{rank}(T) = \dim(\text{range}(T)) = 1$

22. (a) Because $T(\mathbf{x}) = \mathbf{0}$ has only the trivial solution $\mathbf{x} = (0, 0)$, the kernel of T is $\{(0, 0)\}$.

(b) $\text{nullity}(T) = \dim(\ker(T)) = 0$

- (c) Transpose A and find the equivalent row-echelon form.

$$A^T = \begin{bmatrix} 4 & 0 & 2 \\ 1 & 0 & -3 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

So, $\text{range}(T) = \{(t, 0, s) : s, t \in R\}.$

(d) $\text{rank}(T) = \dim(\text{range}(T)) = 2$

28. (a) The kernel of T is given by the solution to the equation $T(\mathbf{x}) = \mathbf{0}$. So,

$$\ker(T) = \{(-t, 0, t) : t \text{ is any real number}\}.$$

(b) $\text{nullity}(T) = \dim(\ker(T)) = 1$

- (c) Transpose A and find its equivalent reduced row-echelon form.

$$A^T = \begin{bmatrix} -\frac{1}{3} & \frac{2}{3} & -\frac{1}{3} \\ \frac{2}{3} & \frac{1}{3} & \frac{2}{3} \\ -\frac{1}{3} & \frac{2}{3} & -\frac{1}{3} \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

So, $\text{range}(T) = \{(s, 0, s), (0, t, 0) : s \text{ and } t \text{ are any real numbers}\}.$

(d) $\text{rank}(T) = \dim(\text{range}(T)) = 2$

24. (a) The kernel of T is given by the solution to the equation $T(\mathbf{x}) = \mathbf{0}$. So,

$$\ker(T) = \{(5t, t) : t \in R\}.$$

(b) $\text{nullity}(T) = \dim(\ker(T)) = 1$

- (c) Transpose A and find its equivalent row-echelon form.

$$A^T = \begin{bmatrix} \frac{1}{26} & -\frac{5}{26} \\ -\frac{5}{26} & \frac{25}{26} \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & -5 \\ 0 & 0 \end{bmatrix}$$

So, $\text{range}(T) = \{(t, -5t) : t \in R\}.$

(d) $\text{rank}(T) = \dim(\text{range}(T)) = 1$

26. (a) The kernel of T is given by the solution to the equation $T(\mathbf{x}) = \mathbf{0}$. So,

$$\ker(T) = \{(0, t, 0) : t \in R\}.$$

(b) $\text{nullity}(T) = \dim(\ker(T)) = 1$

- (c) Transpose A and find its equivalent row-echelon form.

$$A^T = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

So, $\text{range}(T) = \{(t, 0, s) : s, t \in R\}.$

(d) $\text{rank}(T) = \dim(\text{range}(T)) = 2$

30. (a) The kernel of T is given by the solution to the equation $T(\mathbf{x}) = \mathbf{0}$. So,

$$\ker(T) = \{(t, -t, s, -s) : s, t \in \mathbb{R}\}.$$

(b) $\text{nullity}(T) = \dim(\ker(T)) = 2$

- (c) Transpose A and find its equivalent row-echelon form.

$$A^T = \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 0 & 1 \\ 0 & 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$$

So, $\text{range}(T) = \mathbb{R}^2$.

(d) $\text{rank}(T) = \dim(\text{range}(T)) = 2$

32. (a) The kernel of T is given by the solution to the equation $T(\mathbf{x}) = \mathbf{0}$. So,

$$\ker(T) = \{(-t - s - 2r, 6t - 2s, r, s, t) : r, s, t \in \mathbb{R}\}.$$

(b) $\text{nullity}(T) = \dim(\ker(T)) = 3$

- (c) Transpose A and find its equivalent row-echelon form.

$$A^T = \begin{bmatrix} 3 & 4 & 2 \\ -2 & 3 & -3 \\ 6 & 8 & 4 \\ -1 & 10 & -4 \\ 15 & -14 & 20 \end{bmatrix} \Rightarrow \begin{bmatrix} 17 & 0 & 18 \\ 0 & 17 & -5 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

So, $\text{range}(T) = \{(17s, 17t, 18s - 5t) : s, t \in \mathbb{R}\}$.

(d) $\text{rank}(T) = \dim(\text{range}(T)) = 2$

42. $\text{rank}(T) + \text{nullity}(T) = \dim \mathbb{R}^4 \Rightarrow \text{nullity}(T) = 4 - 0 = 4$

44. $\text{rank}(T) + \text{nullity}(T) = \dim P_3 \Rightarrow \text{nullity}(T) = 4 - 2 = 2$

46. $\text{rank}(T) + \text{nullity}(T) = \dim M_{3,3} \Rightarrow \text{nullity}(T) = 9 - 6 = 3$

48. Because $|A| = -1 \neq 0$, the homogeneous equation $A\mathbf{x} = \mathbf{0}$ has only the trivial solution. So, $\ker(T) = \{(0, 0)\}$ and T is one-to-one (by Theorem 6.6). Furthermore, because $\text{rank}(T) = \dim(\mathbb{R}^2) - \text{nullity}(T) = 2 - 0 = 2 = \dim(\mathbb{R}^2)$, T is onto (by Theorem 6.7).

50. Because $|A| = -24 \neq 0$, the homogeneous equation $A\mathbf{x} = \mathbf{0}$ has only the trivial solution. So, $\ker(T) = \{(0, 0, 0)\}$ and T is one-to-one (by Theorem 6.6). Furthermore, because $\text{rank}(T) = \dim \mathbb{R}^3 - \text{nullity}(T) = 3 - 0 = 3 = \dim(\mathbb{R}^3)$, T is onto (by Theorem 6.7).

52. The matrix representation of $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is given by $A = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$.

The matrix in row-echelon form is $A = \begin{bmatrix} 1 & -1 \\ 0 & 0 \end{bmatrix}$.

So, you have the following.

$$\dim(\text{domain}) = 2, \text{rank}(T) = 1, \text{nullity}(T) = 1$$

Because the rank of T is not equal to the dimension of \mathbb{R}^2 , T is not onto. Because $\ker(T) \neq \{\mathbf{0}\}$, T is not one-to one.

34. Because $\text{rank}(T) + \text{nullity}(T) = 3$, and you are given $\text{rank}(T) = 1$, then $\text{nullity}(T) = 2$. So, the kernel of T is a plane, and the range is a line.

36. Because $\text{rank}(T) + \text{nullity}(T) = 3$, and you are given $\text{rank}(T) = 3$, then $\text{nullity}(T) = 0$. So, the kernel of T is the single point $\{(0, 0, 0)\}$, and the range is all of \mathbb{R}^3 .

38. The kernel of T is determined by solving $T(x, y, z) = (-x, y, z) = (0, 0, 0)$, which implies that the kernel is the single point $\{(0, 0, 0)\}$. From the equation $\text{rank}(T) + \text{nullity}(T) = 3$, you see that the rank of T is 3. So, the range of T is all of \mathbb{R}^3 .

40. The kernel of T is determined by solving $T(x, y, z) = (x, y, 0) = (0, 0, 0)$, which implies that $x = y = 0$. So, the nullity of T is 1, and the kernel is a line (the z -axis). The range of T is found by observing that $\text{rank}(T) + \text{nullity}(T) = 3$. That is, the range of T is 2-dimensional, the xy -plane in \mathbb{R}^3 .

$$54. A = \begin{bmatrix} -1 & 3 & 2 & 1 & 4 \\ 2 & 3 & 5 & 0 & 0 \\ 2 & 1 & 2 & 1 & 0 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & 0 & \frac{10}{9} & \frac{4}{9} \\ 0 & 1 & 0 & \frac{5}{3} & \frac{8}{3} \\ 0 & 0 & 1 & \frac{-13}{9} & \frac{-16}{9} \end{bmatrix}$$

So, you have the following.

$$\dim(\text{domain}) = 5, \text{rank}(T) = 3, \text{nullity}(T) = 2$$

Because the rank of T is equal to the dimension of R^3 , T is onto. Because $\ker(T) \neq \{\mathbf{0}\}$, T is not one-to-one.

56. The vector spaces isomorphic to R^6 are those whose dimension is six. That is, (a) $M_{2,3}$ (d) $M_{6,1}$ (e) P_5 and (g) $\{(x_1, x_2, x_3, 0, x_5, x_6, x_7) : x_i \in R\}$ are isomorphic to R^6 .

58. Solve the equation $T(p) = \int_0^1 p(x)dx = \int_0^1 (a_0 + a_1x + a_2x^2)dx = 0$ yielding $a_0 + a_1/2 + a_2/3 = 0$.

Letting $a_2 = -3b$, $a_1 = -2a$, you have $a_0 = -a_1/2 - a_2/3 = a + b$, and $\ker(T) = \{(a + b) - 2ax - 3bx^2 : a, b \in R\}$.

$$60. A = \begin{bmatrix} 1 & -2 & 1 & 0 \\ 0 & 1 & 2 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & 5 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

(a) $\dim(R^4) = 4$

(b) $\dim(R^3) = 3$

(c) $x_1 + 5x_3 = 0 \rightarrow x_1 = -5x_3$

$x_2 + 2x_3 = 0 \rightarrow x_2 = -2x_3$

$x_4 = 0$

So, $\ker(T) = \{(-5x_3, -2x_3, x_3, 0)\}$ and

$\dim(\ker(T)) = 1$.

(d) T is not one-to-one since the $\ker(T) \neq \{\mathbf{0}\}$.

(e) $\text{rank}(T) = 3$

$= \dim(R^3)$

So, T is onto by Theorem 6.7.

(f) T is not an isomorphism since it is not one-to-one.

70. $T^{-1}(U)$ is nonempty because $T(\mathbf{0}) = \mathbf{0} \in U \Rightarrow \mathbf{0} \in T^{-1}(U)$.

Let $\mathbf{v}_1, \mathbf{v}_2 \in T^{-1}(U) \Rightarrow T(\mathbf{v}_1) \in U$ and $T(\mathbf{v}_2) \in U$. Because U is a subspace of W ,

$$T(\mathbf{v}_1) + T(\mathbf{v}_2) = T(\mathbf{v}_1 + \mathbf{v}_2) \in U \Rightarrow \mathbf{v}_1 + \mathbf{v}_2 \in T^{-1}(U).$$

Let $\mathbf{v} \in T^{-1}(U)$ and $c \in R \Rightarrow T(\mathbf{v}) \in U$. Because U is a subspace of W , $cT(\mathbf{v}) = T(c\mathbf{v}) \in U \Rightarrow c\mathbf{v} \in T^{-1}(U)$.

If $U = \{\mathbf{0}\}$, then $T^{-1}(U)$ is the kernel of T .

62. If T is onto, then $m \geq n$.

If T is one-to-one, then $m \leq n$.

64. Theorem 6.9 tells you that if $M_{m,n}$ and $M_{j,k}$ are of the same dimension then they are isomorphic. So, you can conclude that $mn = jk$.

66. (a) False. A concept of a dimension of a linear transformation does not exist.

(b) True. See discussion on page 315 before Theorem 6.6.

(c) True. Because $\dim(P_1) = \dim(R^2) = 2$ and any two vector spaces of equal finite dimension are isomorphic (Theorem 6.9 on page 317).

68. From Theorem 6.5,

$$\text{rank}(T) + \text{nullity}(T) = n = \text{dimension of } V. T \text{ is}$$

one-to-one if and only if $\text{nullity}(T) = 0$ if and only if

$$\text{rank}(T) = \text{dimension of } V.$$

Section 6.3 Matrices for Linear Transformations

2. Because

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 2 \\ 1 \\ -4 \end{bmatrix} \quad \text{and} \quad T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix},$$

$$\text{the standard matrix for } T \text{ is } A = \begin{bmatrix} 2 & -3 \\ 1 & -1 \\ -4 & 1 \end{bmatrix}.$$

4. Because

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 5 \\ 0 \\ 4 \end{bmatrix} \quad \text{and} \quad T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 0 \\ -5 \end{bmatrix},$$

the standard matrix for T is

$$\begin{bmatrix} 5 & 1 \\ 0 & 0 \\ 4 & -5 \end{bmatrix}.$$

6. Because

$$T\left(\begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad T\left(\begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad T\left(\begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

$$\text{and } T\left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

$$\text{the standard matrix for } T \text{ is } A = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

8. Because

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 1 \\ 2 \\ 0 \end{bmatrix} \quad \text{and} \quad T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ -1 \\ 0 \\ 2 \end{bmatrix},$$

the standard matrix for T is

$$A = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 0 \\ 0 & 2 \end{bmatrix}.$$

$$\text{So, } T(\mathbf{v}) = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 3 \\ -3 \end{bmatrix} = \begin{bmatrix} 0 \\ 6 \\ 6 \\ -6 \end{bmatrix}$$

$$\text{and } T(3, -3) = (0, 6, 6, -6).$$

$$10. T(x_1, x_2, x_3, x_4) = (x_1 - x_3, x_2 - x_4, x_3 - x_1, x_2 + x_4)$$

The standard matrix is

$$A = \begin{bmatrix} 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & -1 \\ -1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}.$$

The image of \mathbf{v} is

$$A\mathbf{v} = \begin{bmatrix} 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & -1 \\ -1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ -2 \end{bmatrix} = \begin{bmatrix} -2 \\ 4 \\ 2 \\ 0 \end{bmatrix}.$$


$$\text{So, } T(\mathbf{v}) = (-2, 4, 2, 0).$$

$$12. (a) \text{ The matrix of a reflection in the line } y = x, T(x, y) = (y, x), \text{ is given by } A = [T(1, 0) : T(0, 1)] = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

(b) The image of $\mathbf{v} = (3, 4)$ is given by

$$A\mathbf{v} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \end{bmatrix}.$$

$$\text{So, } T(3, 4) = (4, 3).$$




14. (a) The matrix of a reflection in the x -axis, $T(x, y) = (x, -y)$, is given by

$$A = [T(1, 0) : T(0, 1)] = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

- (b) The image of $\mathbf{v} = (4, -1)$ is given by

$$A\mathbf{v} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 4 \\ -1 \end{bmatrix} = \begin{bmatrix} 4 \\ 1 \end{bmatrix}.$$

So, $T(4, -1) = (4, 1)$.



16. (a) The counterclockwise rotation of 120° is given by

$$\begin{aligned} T(x, y) &= (\cos(120)x - \sin(120)y, \sin(120)x + \cos(120)y) \\ &= \left(-\frac{1}{2}x - \frac{\sqrt{3}}{2}y, \frac{\sqrt{3}}{2}x - \frac{1}{2}y \right). \end{aligned}$$


So, the matrix is

$$A = [T(1, 0) : T(0, 1)] = \begin{bmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{bmatrix}.$$

- (b) The image of $\mathbf{v} = (2, 2)$ is given by

$$A\mathbf{v} = \begin{bmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{bmatrix} \begin{bmatrix} 2 \\ 2 \end{bmatrix} = \begin{bmatrix} -1 - \sqrt{3} \\ \sqrt{3} - 1 \end{bmatrix}.$$

So, $T(2, 2) = (-1 - \sqrt{3}, \sqrt{3} - 1)$.



18. (a) The clockwise rotation of 30° is given by

$$\begin{aligned} T(x, y) &= (\cos(-30)x - \sin(-30)y, \sin(-30)x + \cos(-30)y) \\ &= \left(\frac{\sqrt{3}}{2}x + \frac{1}{2}y, -\frac{1}{2}x + \frac{\sqrt{3}}{2}y \right). \end{aligned}$$


So, the matrix is

$$A = [T(1, 0) : T(0, 1)] = \begin{bmatrix} \frac{\sqrt{3}}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}.$$

- (b) The image of $\mathbf{v} = (2, 1)$ is given by

$$A\mathbf{v} = \begin{bmatrix} \frac{\sqrt{3}}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} \sqrt{3} + \frac{1}{2} \\ -1 + \frac{\sqrt{3}}{2} \end{bmatrix}.$$

$$\text{So, } T(2, 1) = \left(\sqrt{3} + \frac{1}{2}, -1 + \frac{\sqrt{3}}{2} \right).$$




20. (a) The matrix of a reflection through the yz -coordinate plane is given by

$$A = [T(1, 0, 0) : T(0, 1, 0) : T(0, 0, 1)] = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

- (b) The image of $\mathbf{v} = (2, 3, 4)$ is given by

$$A\mathbf{v} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix} = \begin{bmatrix} -2 \\ 3 \\ 4 \end{bmatrix}.$$

$$\text{So, } T(2, 3, 4) = (-2, 3, 4).$$



22. (a) The reflection of a vector
- \mathbf{v}
- through
- \mathbf{w}
- is given by

$$\begin{aligned} T(\mathbf{v}) &= 2 \operatorname{proj}_{\mathbf{w}} \mathbf{v} - \mathbf{v} \\ T(x, y) &= 2 \frac{3x + y}{10} (3, 1) - (x, y) \\ &= \left(\frac{4}{5}x + \frac{3}{5}y, \frac{3}{5}x - \frac{4}{5}y \right). \end{aligned}$$

The standard matrix for T is


$$A = [T(1, 0) : T(0, 1)] = \begin{bmatrix} \frac{4}{5} & \frac{3}{5} \\ \frac{3}{5} & -\frac{4}{5} \end{bmatrix}.$$

- (b) The image of
- $\mathbf{v} = (1, 4)$
- is

$$A\mathbf{v} = \begin{bmatrix} \frac{4}{5} & \frac{3}{5} \\ \frac{3}{5} & -\frac{4}{5} \end{bmatrix} \begin{bmatrix} 1 \\ 4 \end{bmatrix} = \begin{bmatrix} \frac{16}{5} \\ -\frac{13}{5} \end{bmatrix}.$$

$$\text{So, } T(1, 4) = \left(\frac{16}{5}, -\frac{13}{5} \right).$$

- (c)



24. (a) The standard matrix for
- T
- is

$$A = \begin{bmatrix} 1 & 2 & -3 \\ 3 & -5 & 0 \\ 0 & 1 & -3 \end{bmatrix}.$$

- (b) The image of
- $\mathbf{v} = (3, 13, 4)$
- is

$$A\mathbf{v} = \begin{bmatrix} 1 & 2 & -3 \\ 3 & -5 & 0 \\ 0 & 1 & -3 \end{bmatrix} \begin{bmatrix} 3 \\ 13 \\ 4 \end{bmatrix} = \begin{bmatrix} 17 \\ -56 \\ 1 \end{bmatrix}.$$

$$\text{So, } T(3, 13, 4) \text{ is } (17, -56, 1).$$

- (c) Using a graphing utility or a computer software program to perform the multiplication in part (b) gives the same results.

26. (a) The standard matrix for
- T
- is

$$A = \begin{bmatrix} 1 & 2 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 2 & -1 \\ 1 & 0 & 0 & 0 \end{bmatrix}.$$

- (b) The image of
- $\mathbf{v} = (0, 1, -1, 1)$
- is

$$A\mathbf{v} = \begin{bmatrix} 1 & 2 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 2 & -1 \\ 1 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ -3 \\ 0 \end{bmatrix}.$$

$$\text{So, } T(0, 1, -1, 1) = (2, 1, -3, 0).$$

- (c) Using a graphing utility or a computer software program to perform the multiplication in part (b) gives the same result.

28. The standard matrices for
- T_1
- and
- T_2
- are

$$A_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \text{and} \quad A_2 = \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

The standard matrix for $T = T_2 \circ T_1$ is

$$A = A_2 A_1 = \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = A_2$$

and the standard matrix for $T' = T_1 \circ T_2$ is

$$A' = A_1 A_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = A_2.$$

30. The standard matrices for
- T_1
- and
- T_2
- are

$$A_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad A_2 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

The standard matrix for $T = T_2 \circ T_1$ is

$$A_2 A_1 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}$$

and the standard matrix for $T' = T_1 \circ T_2$ is

$$A_1 A_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

32. The standard matrix for
- T
- is

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 0 \end{bmatrix}.$$

Because $|A| = 0$, A is not invertible, and so T is not invertible.

34. The standard matrix for
- T
- is

$$A = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}.$$

Because $|A| = -2 \neq 0$, A is invertible.

$$A^{-1} = -\frac{1}{2} \begin{bmatrix} -1 & -1 \\ -1 & 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{bmatrix}$$

So, $T^{-1}(x, y) = (\frac{1}{2}x + \frac{1}{2}y, \frac{1}{2}x - \frac{1}{2}y)$.

38. (a) The standard matrix for
- T
- is

$$A' = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \end{bmatrix}$$

and the image of \mathbf{v} under T is

$$A'\mathbf{v} = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \end{bmatrix} \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix} = \begin{bmatrix} -2 \\ -2 \end{bmatrix}.$$

So, $T(\mathbf{v}) = (-2, -2)$.

- (b) The image of each vector in
- B
- is as follows.

$$T(1, 1, 1) = (0, 0) = 0(1, 1) + 0(2, 1)$$

$$T(1, 1, 0) = (0, 1) = -1(1, 1) + (1, 2)$$

$$T(0, 1, 1) = (-1, 0) = -2(1, 1) + (1, 2)$$

$$\text{So, } [T(1, 1, 1)]_{B'} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, [T(1, 1, 0)]_{B'} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}, \text{ and } [T(0, 1, 1)]_{B'} = \begin{bmatrix} -2 \\ 1 \end{bmatrix},$$

$$\text{which implies that } A = \begin{bmatrix} 0 & -1 & -2 \\ 0 & 1 & 1 \end{bmatrix}.$$

$$\text{Then, because } [\mathbf{v}]_B = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}, [T(\mathbf{v})]_{B'} = A[\mathbf{v}]_B = \begin{bmatrix} 0 & -1 & -2 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} -2 \\ 0 \end{bmatrix}.$$

So, $T(\mathbf{v}) = -2(1, 1) + 0(1, 2) = (-2, 2)$.

36. The standard matrix for
- T
- is

$$A = \begin{bmatrix} 1 & -2 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}.$$

Because $|A| = -1 \neq 0$, A is invertible. Calculate A^{-1} by Gauss-Jordan elimination

$$A^{-1} = \begin{bmatrix} 1 & 2 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & -1 \end{bmatrix}$$

and conclude that

$$T^{-1}(x_1, x_2, x_3, x_4) = (x_1 + 2x_2, x_2, x_4, x_3 - x_4).$$

40. (a) The standard matrix for
- T
- is

$$A' = \begin{bmatrix} 1 & 1 & 1 & 1 \\ -1 & 0 & 0 & 1 \end{bmatrix}$$

and the image of $\mathbf{v} = (4, -3, 1, 1)$ under T is

$$A'\mathbf{v} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ -1 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 4 \\ -3 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ -3 \end{bmatrix} \Rightarrow T(\mathbf{v}) = (3, -3).$$

- (b) Because

$$T(1, 0, 0, 1) = (2, 0) = 0(1, 1) + (2, 0)$$

$$T(0, 1, 0, 1) = (2, 1) = (1, 1) + \frac{1}{2}(2, 0)$$

$$T(1, 0, 1, 0) = (2, -1) = -(1, 1) + \frac{3}{2}(2, 0)$$

$$T(1, 1, 0, 0) = (2, -1) = -(1, 1) + \frac{3}{2}(2, 0),$$

The matrix for T relative to B and B' is $A = \begin{bmatrix} 0 & 1 & -1 & -1 \\ 1 & \frac{1}{2} & \frac{3}{2} & \frac{3}{2} \end{bmatrix}$.

Because $\mathbf{v} = (4, -3, 1, 1) = \frac{7}{2}(1, 0, 0, 1) - \frac{5}{2}(0, 1, 0, 1) + (1, 0, 1, 0) - \frac{1}{2}(1, 1, 0, 0)$, you have

$$[T(\mathbf{v})]_{B'} = A[\mathbf{v}]_B = \begin{bmatrix} 0 & 1 & -1 & -1 \\ 1 & \frac{1}{2} & \frac{3}{2} & \frac{3}{2} \end{bmatrix} \begin{bmatrix} \frac{7}{2} \\ -\frac{5}{2} \\ 1 \\ -\frac{1}{2} \end{bmatrix} = \begin{bmatrix} -3 \\ 3 \end{bmatrix}.$$

So, $T(\mathbf{v}) = -3(1, 1) + 3(2, 0) = (3, -3)$.

42. (a) The standard matrix for
- T
- is
- $A' = \begin{bmatrix} 3 & -13 \\ 1 & -4 \end{bmatrix}$
- and the image of
- $\mathbf{v} = (4, 8)$
- under
- T
- is

$$A'\mathbf{v} = \begin{bmatrix} 3 & -13 \\ 1 & -4 \end{bmatrix} \begin{bmatrix} 4 \\ 8 \end{bmatrix} = \begin{bmatrix} -92 \\ -28 \end{bmatrix} \Rightarrow T(\mathbf{v}) = (-92, -28).$$

- (b) Because

$$T(2, 1) = (-7, -2) = -(2, 1) - (5, 1)$$

$$T(5, 1) = (2, 1)$$

the matrix for T relative to B and B' is $A = \begin{bmatrix} -1 & 0 \\ -1 & 1 \end{bmatrix}$.

Because $\mathbf{v} = (4, 8) = 12(2, 1) - 4(5, 1)$, you have $[T(\mathbf{v})]_{B'} = A[\mathbf{v}]_B = \begin{bmatrix} -1 & 0 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 12 \\ -4 \end{bmatrix} = \begin{bmatrix} -12 \\ -16 \end{bmatrix}$.

So, $T(\mathbf{v}) = -12(2, 1) - 16(5, 1) = (-92, -28)$.

44. The image of each vector in
- B
- is
- $T(1) = x^2$
- ,
- $T(x) = x^3$
- ,
- $T(x^2) = x^4$
- .

So, the matrix of T relative to B and B' is $A = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$.

46. The image of each vector in B is as follows.

$$D(e^{2x}) = 2e^{2x}$$

$$D(xe^{2x}) = e^{2x} + 2xe^{2x}$$

$$D(x^2e^{2x}) = 2xe^{2x} + 2x^2e^{2x}$$

So, the matrix of T relative to B is $A = \begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & 2 \\ 0 & 0 & 2 \end{bmatrix}$.

48. Because $5e^{2x} - 3xe^{2x} + x^2e^{2x} = 5(e^{2x}) - 3(xe^{2x}) + 1(x^2e^{2x})$,

$$A[\mathbf{v}]_B = \begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & 2 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} 5 \\ -3 \\ 1 \end{bmatrix} = \begin{bmatrix} 7 \\ -4 \\ 2 \end{bmatrix} \Rightarrow D_x(5e^{2x} - 3xe^{2x} + x^2e^{2x}) = 7e^{2x} - 4xe^{2x} + 2x^2e^{2x}.$$

50. (a) Let $T : R^n \rightarrow R^m$ be a linear transformation such that, for the standard basis vectors \mathbf{e}_i of R^n ,

$$T(\mathbf{e}_1) = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix}, T(\mathbf{e}_2) = \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix}, \dots, T(\mathbf{e}_n) = \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix}.$$

Then the $m \times n$ matrix whose n columns correspond to $T(\mathbf{e}_i)$

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

is such that $T(\mathbf{v}) = A\mathbf{v}$ for every \mathbf{v} in R^n . A is called the standard matrix of T .

- (b) Let $T_1 : R^n \rightarrow R^m$ and $T_2 : R^m \rightarrow R^p$ be linear transformations with standard matrices A_1 and A_2 , respectively. The composition $T : R^n \rightarrow R^p$, defined by $T(\mathbf{v}) = T_2(T_1(\mathbf{v}))$, is a linear transformation. Moreover, the standard matrix A for T is given by the matrix product $A = A_2A_1$.
- (c) To find the inverse of a linear transformation T , first find the standard matrix A of T . Then find the inverse of A using the techniques shown in Section 2.3.
- (d) To find the transformation matrix relative to nonstandard basis, first find $T(\mathbf{v}_1), T(\mathbf{v}_2), \dots, T(\mathbf{v}_n)$. Then determine the coordinate matrices relative to B' . Finally, form the matrix T relative to B and B' by using the coordinate matrices as

columns to produce $A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix}$

52. Because $T(\mathbf{v}) = k\mathbf{v}$ for all $\mathbf{v} \in R^n$, the standard matrix for T is the $n \times n$ diagonal matrix

$$\begin{bmatrix} k & 0 & \cdots & 0 \\ 0 & k & & \vdots \\ \vdots & & k & 0 \\ 0 & \cdots & 0 & k \end{bmatrix}.$$

54. (a) True. See discussion, under “Composition of Linear Transformations,” pages 323–324.
- (b) False. See Example 3, page 324.

56. (1 \Rightarrow 2): Let T be invertible. If $T(\mathbf{v}_1) = T(\mathbf{v}_2)$, then $T^{-1}(T(\mathbf{v}_1)) = T^{-1}(T(\mathbf{v}_2))$ and $\mathbf{v}_1 = \mathbf{v}_2$, so T is one-to-one. T is onto because for any $\mathbf{w} \in R^n$, $T^{-1}(\mathbf{w}) = \mathbf{v}$ satisfies $T(\mathbf{v}) = \mathbf{w}$.

(2 \Rightarrow 1): Let T be an isomorphism. Define T^{-1} as follows: Because T is onto, for any $\mathbf{w} \in R^n$, there exists $\mathbf{v} \in R^n$ such that $T(\mathbf{v}) = \mathbf{w}$. Because T is one-to-one, this \mathbf{v} is unique. So, define the inverse of T by $T^{-1}(\mathbf{w}) = \mathbf{v}$ if and only if $T(\mathbf{v}) = \mathbf{w}$.

Finally, the corollaries to Theorems 6.3 and 6.4 show that 2 and 3 are equivalent.

If T is invertible, $T(\mathbf{x}) = A\mathbf{x}$ implies that $T^{-1}(T(\mathbf{x})) = \mathbf{x} = A^{-1}(A\mathbf{x})$ and the standard matrix of T^{-1} is A^{-1} .

58. \mathbf{b} is in the range of the linear transformation $T: R^n \rightarrow R^m$ given by $T(\mathbf{x}) = A\mathbf{x}$ if and only if \mathbf{b} is in the column space of A .

Section 6.4 Transition Matrices and Similarity

2. The standard matrix for T is $A = \begin{bmatrix} 2 & 1 \\ 1 & -2 \end{bmatrix}$. Furthermore, the transition matrix P from B' to the standard basis B , and its

inverse are $P = \begin{bmatrix} 1 & 0 \\ 2 & 4 \end{bmatrix}$ and $P^{-1} = \begin{bmatrix} 1 & 0 \\ -\frac{1}{2} & \frac{1}{4} \end{bmatrix}$. Therefore, the matrix for T relative to B' is

$$A' = P^{-1}AP = \begin{bmatrix} 1 & 0 \\ -\frac{1}{2} & \frac{1}{4} \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 1 & -2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 2 & 4 \end{bmatrix} = \begin{bmatrix} 4 & 4 \\ -\frac{11}{4} & -4 \end{bmatrix}$$

4. The standard matrix for T is $A = \begin{bmatrix} 1 & -2 \\ 4 & 0 \end{bmatrix}$. Furthermore, the transition matrix P from B' to the standard basis B , and its

inverse, are $P = \begin{bmatrix} -2 & -1 \\ 1 & 1 \end{bmatrix}$ and $P^{-1} = \begin{bmatrix} -1 & -1 \\ 1 & 2 \end{bmatrix}$. Therefore, the matrix for T relative to B' is

$$A' = P^{-1}AP = \begin{bmatrix} -1 & -1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 & -2 \\ 4 & 0 \end{bmatrix} \begin{bmatrix} -2 & -1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 12 & 7 \\ -20 & -11 \end{bmatrix}$$

6. The standard matrix for T is $A = \begin{bmatrix} 5 & 4 \\ 4 & 5 \end{bmatrix}$. Furthermore, the transition matrix P from B' to the standard basis B , and its

inverse, are $P = \begin{bmatrix} 12 & 13 \\ -13 & -12 \end{bmatrix}$ and $P^{-1} = \begin{bmatrix} -\frac{12}{25} & -\frac{13}{25} \\ \frac{13}{25} & \frac{12}{25} \end{bmatrix}$. Therefore, the matrix for T relative to B' is

$$A' = P^{-1}AP = \begin{bmatrix} -\frac{12}{25} & -\frac{13}{25} \\ \frac{13}{25} & \frac{12}{25} \end{bmatrix} \begin{bmatrix} 5 & 4 \\ 4 & 5 \end{bmatrix} \begin{bmatrix} 12 & 13 \\ -13 & -12 \end{bmatrix} = \begin{bmatrix} 5 & -4 \\ -4 & 5 \end{bmatrix}$$

8. The standard matrix for T is $A = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$. Furthermore, the transition matrix P from B' to the standard basis B , and its

inverse, are $P = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}$ and $P^{-1} = \frac{1}{2} \begin{bmatrix} 1 & 1 & -1 \\ 1 & -1 & 1 \\ -1 & 1 & 1 \end{bmatrix}$. Therefore, the matrix for T relative to B' is

$$A' = P^{-1}AP = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

10. The standard matrix for T is $A = \begin{bmatrix} -1 & 0 & 0 \\ 1 & -1 & 0 \\ 0 & 1 & -1 \end{bmatrix}$. Furthermore, the transition matrix P from B' to the standard basis B , and

its inverse, are $P = \begin{bmatrix} 0 & -2 & 1 \\ -1 & 0 & 3 \\ 2 & 3 & 0 \end{bmatrix}$ and $P^{-1} = \begin{bmatrix} \frac{3}{5} & -\frac{1}{5} & \frac{2}{5} \\ -\frac{2}{5} & \frac{2}{15} & \frac{1}{15} \\ \frac{1}{5} & \frac{4}{15} & \frac{3}{15} \end{bmatrix}$. Therefore, the matrix for T relative to B' is

$$A' = P^{-1}AP = \begin{bmatrix} \frac{3}{5} & -\frac{1}{5} & \frac{2}{5} \\ -\frac{2}{5} & \frac{2}{15} & \frac{1}{15} \\ \frac{1}{5} & \frac{4}{15} & \frac{3}{15} \end{bmatrix} \begin{bmatrix} -1 & 0 & 0 \\ 1 & -1 & 0 \\ 0 & 1 & -1 \end{bmatrix} \begin{bmatrix} 0 & -2 & 1 \\ -1 & 0 & 3 \\ 2 & 3 & 0 \end{bmatrix} = \begin{bmatrix} -\frac{7}{5} & \frac{2}{5} & 1 \\ -\frac{1}{15} & -\frac{19}{15} & \frac{1}{3} \\ -\frac{2}{15} & -\frac{8}{15} & -\frac{1}{3} \end{bmatrix}.$$

12. The standard matrix for T is $A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 2 & 0 \\ 1 & 1 & 3 \end{bmatrix}$. Furthermore, the transition matrix P from B' to the standard basis B , and

its inverse, are $P = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 0 & 1 \\ 0 & 1 & -1 \end{bmatrix}$ and $P^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$. Therefore, the matrix for T relative to B' is

$$A' = P^{-1}AP = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 1 & 2 & 0 \\ 1 & 1 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -1 & 0 & 1 \\ 0 & 1 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 2 \end{bmatrix}.$$

14. (a) The transition matrix P from B' to B is found by row-reducing $[B : B']$ to $[I : P]$.

$$[B : B'] = \begin{bmatrix} 1 & -2 & : & 1 & 0 \\ 1 & 3 & : & -1 & 1 \end{bmatrix} \Rightarrow [I : P] = \begin{bmatrix} 1 & 0 & : & \frac{1}{5} & \frac{2}{5} \\ 0 & 1 & : & -\frac{2}{5} & \frac{1}{5} \end{bmatrix}$$

$$\text{So, } P = \begin{bmatrix} \frac{1}{5} & \frac{2}{5} \\ -\frac{2}{5} & \frac{1}{5} \end{bmatrix}.$$

- (b) The coordinate matrix for \mathbf{v} relative to B is $[\mathbf{v}]_B = P[\mathbf{v}]_{B'} = \begin{bmatrix} \frac{1}{5} & \frac{2}{5} \\ -\frac{2}{5} & \frac{1}{5} \end{bmatrix} \begin{bmatrix} 1 \\ -3 \end{bmatrix} = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$.

$$\text{Furthermore, the image of } \mathbf{v} \text{ under } T \text{ relative to } B \text{ is } [T(\mathbf{v})]_B = A[\mathbf{v}]_B = \begin{bmatrix} 3 & 2 \\ 0 & 4 \end{bmatrix} \begin{bmatrix} -1 \\ -1 \end{bmatrix} = \begin{bmatrix} -5 \\ -4 \end{bmatrix}.$$

- (c) The matrix of T relative to B' is $A' = P^{-1}AP = \begin{bmatrix} 1 & -2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 3 & 2 \\ 0 & 4 \end{bmatrix} \begin{bmatrix} \frac{1}{5} & \frac{2}{5} \\ -\frac{2}{5} & \frac{1}{5} \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ -2 & 4 \end{bmatrix}$.

- (d) The image of \mathbf{v} under T relative to B' is $P^{-1}[T(\mathbf{v})]_B = \begin{bmatrix} 1 & -2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} -5 \\ -4 \end{bmatrix} = \begin{bmatrix} 3 \\ -14 \end{bmatrix}$.

$$\text{You can also find the image of } \mathbf{v} \text{ under } T \text{ relative to } B' \text{ by } A'[\mathbf{v}]_{B'} = \begin{bmatrix} 3 & 0 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} 1 \\ -3 \end{bmatrix} = \begin{bmatrix} 3 \\ -14 \end{bmatrix}.$$

16. (a) The transition matrix P from B' to B is found by row-reducing $[B : B']$ to $[I : P]$.

$$P = \begin{bmatrix} -1 & -5 \\ 0 & -3 \end{bmatrix}$$

- (b) The coordinate matrix for \mathbf{v} relative to B is $[\mathbf{v}]_B = P[\mathbf{v}]_{B'} = \begin{bmatrix} -1 & -5 \\ 0 & -3 \end{bmatrix} \begin{bmatrix} 1 \\ -4 \end{bmatrix} = \begin{bmatrix} 19 \\ 12 \end{bmatrix}$.

Furthermore, the image of \mathbf{v} under T relative to B is $[T(\mathbf{v})]_B = A[\mathbf{v}]_B = \begin{bmatrix} 2 & 1 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 19 \\ 12 \end{bmatrix} = \begin{bmatrix} 50 \\ -12 \end{bmatrix}$.

- (c) The matrix of T relative to B' is $A' = P^{-1}AP = \begin{bmatrix} -1 & \frac{5}{3} \\ 0 & -\frac{1}{3} \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} -1 & -5 \\ 0 & -3 \end{bmatrix} = \begin{bmatrix} 2 & 18 \\ 0 & -1 \end{bmatrix}$

- (d) The image of \mathbf{v} under T relative to B' is $P^{-1}[T(\mathbf{v})]_B = \begin{bmatrix} -1 & \frac{5}{3} \\ 0 & -\frac{1}{3} \end{bmatrix} \begin{bmatrix} 50 \\ -12 \end{bmatrix} = \begin{bmatrix} -70 \\ 4 \end{bmatrix}$.

You can also find the image of \mathbf{v} under T relative to B' by $A'[\mathbf{v}]_{B'} = \begin{bmatrix} 2 & 18 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ -4 \end{bmatrix} = \begin{bmatrix} -70 \\ 4 \end{bmatrix}$.

18. (a) The transition matrix P from B' to B is found by row-reducing $[B : B']$ to $[I : P]$.

$$[B : B'] = \begin{bmatrix} 1 & 1 & -1 & 1 & 0 & 0 \\ 1 & -1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 1 & 0 & 0 & 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & 0 & \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 1 & 0 & \frac{1}{2} & 0 & \frac{1}{2} \\ 0 & 0 & 1 & 0 & \frac{1}{2} & \frac{1}{2} \end{bmatrix} = [I : P]$$

So, $P = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & \frac{1}{2} \end{bmatrix}$.

- (b) The coordinate matrix for \mathbf{v} relative to B is $[\mathbf{v}]_B = P[\mathbf{v}]_{B'} = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{3}{2} \\ \frac{3}{2} \\ 1 \end{bmatrix}$.

Furthermore, the image of \mathbf{v} under T relative to B is

$$[T(\mathbf{v})]_B = A[\mathbf{v}]_B = \begin{bmatrix} \frac{3}{2} & -1 & -\frac{1}{2} \\ -\frac{1}{2} & 2 & \frac{1}{2} \\ \frac{1}{2} & 1 & \frac{5}{2} \end{bmatrix} \begin{bmatrix} \frac{3}{2} \\ \frac{3}{2} \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{4} \\ \frac{11}{4} \\ \frac{19}{4} \end{bmatrix}.$$

- (c) The matrix of T relative to B' is $A' = P^{-1}AP$.

$$A' = P^{-1}AP = \begin{bmatrix} 1 & 1 & -1 \\ 1 & -1 & 1 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} \frac{3}{2} & -1 & -\frac{1}{2} \\ -\frac{1}{2} & 2 & \frac{1}{2} \\ \frac{1}{2} & 1 & \frac{5}{2} \end{bmatrix} \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & \frac{1}{2} \end{bmatrix} = \begin{bmatrix} \frac{1}{4} & -1 & -\frac{5}{4} \\ \frac{1}{4} & 2 & -\frac{1}{4} \\ \frac{5}{4} & 1 & \frac{15}{4} \end{bmatrix}$$

- (d) The image of \mathbf{v} under T relative to B' is

$$P^{-1}[T(\mathbf{v})]_B = \begin{bmatrix} 1 & 1 & -1 \\ 1 & -1 & 1 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{4} \\ \frac{11}{4} \\ \frac{19}{4} \end{bmatrix} = \begin{bmatrix} -\frac{7}{4} \\ \frac{9}{4} \\ \frac{29}{4} \end{bmatrix}.$$

You can also find the image of \mathbf{v} under T relative to B' by

$$A'[\mathbf{v}]_{B'} = \begin{bmatrix} \frac{1}{4} & -1 & -\frac{5}{4} \\ \frac{1}{4} & 2 & -\frac{1}{4} \\ \frac{5}{4} & 1 & \frac{15}{4} \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -\frac{7}{4} \\ \frac{9}{4} \\ \frac{29}{4} \end{bmatrix}.$$

20. A is similar to A' since

$$A' = P^{-1}AP = \begin{bmatrix} 1 & 12 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -12 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -12 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & -12 \\ 0 & 1 \end{bmatrix}.$$

22. A is similar to A' since

$$A' = P^{-1}AP = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 5 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 5 & 2 & 2 \\ 0 & 3 & 2 \\ 0 & 0 & 1 \end{bmatrix}.$$

24. The transition matrix from B' to the standard matrix has columns consisting of the vectors in B' .

$$P = \begin{bmatrix} 1 & 1 & -1 \\ 1 & -1 & 1 \\ -1 & 1 & 1 \end{bmatrix}$$

and it follows that

$$P^{-1} = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & \frac{1}{2} \end{bmatrix}.$$

So, the matrix for T relative to B' is

$$\begin{aligned} A' &= P^{-1}AP \\ &= \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} \frac{3}{2} & -1 & -\frac{1}{2} \\ -\frac{1}{2} & 2 & \frac{1}{2} \\ \frac{1}{2} & 1 & \frac{5}{2} \end{bmatrix} \begin{bmatrix} 1 & 1 & -1 \\ 1 & -1 & 1 \\ -1 & 1 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}. \end{aligned}$$

26. First, note that A and B are similar.

$$\begin{aligned} P^{-1}AP &= \begin{bmatrix} -1 & -1 & 2 \\ 0 & -1 & 2 \\ 1 & 2 & -3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} -1 & 1 & 0 \\ 2 & 1 & 2 \\ 1 & 1 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 11 & 7 & 10 \\ 10 & 8 & 10 \\ -18 & -12 & -17 \end{bmatrix} \end{aligned}$$

Now,

$$\begin{aligned} |B| &= \begin{vmatrix} 11 & 7 & 10 \\ 10 & 8 & 10 \\ -18 & -12 & -17 \end{vmatrix} \\ &= 11(-16) - 7(10) + 10(24) \\ &= -6 = |A|. \end{aligned}$$

$$28. \text{ Because } B = P^{-1}AP, \text{ and } A^4 = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}^4 = \begin{bmatrix} 1 & 0 \\ 0 & 16 \end{bmatrix},$$

$$\begin{aligned} \text{you have } B^4 &= P^{-1}A^4P \\ &= \begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 16 \end{bmatrix} \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix} \\ &= \begin{bmatrix} -74 & -225 \\ 30 & 91 \end{bmatrix}. \end{aligned}$$

30. If $B = P^{-1}AP$ and A is an idempotent matrix, then

$$\begin{aligned} B^2 &= (P^{-1}AP)^2 \\ &= (P^{-1}AP)(P^{-1}AP) \\ &= P^{-1}A^2P \\ &= P^{-1}AP \\ &= B, \end{aligned}$$

which shows that B is an idempotent matrix.

32. If $A\mathbf{x} = \mathbf{x}$ and $B = P^{-1}AP$, then $PB = AP$ and $PBP^{-1} = A$. So, $PBP^{-1}\mathbf{x} = A\mathbf{x} = \mathbf{x}$.

34. Because A and B are similar, they represent the same linear transformation with respect to different bases. So, the range is the same, and so is the rank.

36. If A is nonsingular, then so is $P^{-1}AP = B$, and

$$\begin{aligned} B &= P^{-1}AP \\ B^{-1} &= (P^{-1}AP)^{-1} = P^{-1}A^{-1}(P^{-1})^{-1} = P^{-1}A^{-1}P \end{aligned}$$

which shows that A^{-1} and B^{-1} are similar.

38. Because $B = P^{-1}AP$, you have $AP = PB$, as follows.

$$\begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{n1} & \cdots & a_{nn} \end{bmatrix} \begin{bmatrix} p_{11} & \cdots & p_{1n} \\ \vdots & & \vdots \\ p_{n1} & \cdots & p_{nn} \end{bmatrix} = \begin{bmatrix} p_{11} & \cdots & p_{1n} \\ \vdots & & \vdots \\ p_{n1} & \cdots & p_{nn} \end{bmatrix} \begin{bmatrix} b_{11} & \cdots & 0 \\ \vdots & & \vdots \\ 0 & \cdots & b_{nn} \end{bmatrix}$$

So,

$$\begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{n1} & \cdots & a_{nn} \end{bmatrix} \begin{bmatrix} p_{1i} \\ \vdots \\ p_{ni} \end{bmatrix} = b_{ii} \begin{bmatrix} p_{1i} \\ \vdots \\ p_{ni} \end{bmatrix}$$

for $i = 1, 2, \dots, n$.

40. (a) There are two ways to get from the coordinate matrix $[\mathbf{v}]_{B'}$ to the coordinate matrix $[T(\mathbf{v})]_{B'}$. One way is direct, using the matrix A' to obtain $A'[\mathbf{v}]_{B'} = [T(\mathbf{v})]_{B'}$. The second way is indirect, using the matrices P , A , and P^{-1} to obtain

$$P^{-1}AP[\mathbf{v}]_{B'} = [T(\mathbf{v})]_{B'}.$$

(b) To determine if two square matrices A and A' are similar, the equation $A' = P^{-1}AP$ must hold true for some invertible matrix P .

42. (a) True. See discussion, page 330, and note that $A' = P^{-1}AP \Rightarrow PA'P^{-1} = PP^{-1}APP^{-1} = A$.

(b) False. Unless it is a diagonal matrix, see Example 5, page 333.

Section 6.5 Applications of Linear Transformations

2. The standard matrix for T is $A = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$.

$$(a) \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 5 \\ 2 \end{bmatrix} = \begin{bmatrix} -5 \\ 2 \end{bmatrix} \Rightarrow T(5, 2) = (-5, 2)$$

$$(b) \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} -1 \\ -6 \end{bmatrix} = \begin{bmatrix} 1 \\ -6 \end{bmatrix} \Rightarrow T(-1, -6) = (1, -6)$$

$$(c) \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} a \\ 0 \end{bmatrix} = \begin{bmatrix} -a \\ 0 \end{bmatrix} \Rightarrow T(a, 0) = (-a, 0)$$

$$(d) \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ b \end{bmatrix} = \begin{bmatrix} 0 \\ b \end{bmatrix} \Rightarrow T(0, b) = (0, b)$$

$$(e) \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} c \\ -d \end{bmatrix} = \begin{bmatrix} -c \\ -d \end{bmatrix} \Rightarrow T(c, -d) = (-c, -d)$$

$$(f) \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} f \\ g \end{bmatrix} = \begin{bmatrix} -f \\ g \end{bmatrix} \Rightarrow T(f, g) = (-f, g)$$

4. The standard matrix for T is $A = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$.

$$(a) \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} -2 \\ 1 \end{bmatrix} \Rightarrow T(-1, 2) = (-2, 1)$$

$$(b) \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} -3 \\ -2 \end{bmatrix} \Rightarrow T(2, 3) = (-3, -2)$$

$$(c) \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} a \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ -a \end{bmatrix} \Rightarrow T(a, 0) = (0, -a)$$

$$(d) \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ b \end{bmatrix} = \begin{bmatrix} -b \\ 0 \end{bmatrix} \Rightarrow T(0, b) = (-b, 0)$$

$$(e) \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} e \\ -d \end{bmatrix} = \begin{bmatrix} d \\ -e \end{bmatrix} \Rightarrow T(e, -d) = (d, -e)$$

$$(f) \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} -f \\ g \end{bmatrix} = \begin{bmatrix} -g \\ f \end{bmatrix} \Rightarrow T(-f, g) = (-g, f)$$


$$\begin{aligned} 6. (a) \quad T(x, y) &= xT(1, 0) + yT(0, 1) \\ &= x(1, 1) + y(0, 1) \\ &= (x, x + y) \end{aligned}$$

(b) T is vertical shear.

8. $T(x, y) = \left(\frac{x}{4}, y\right)$


(a) Identify T as a horizontal contraction from its

standard matrix $A = \begin{bmatrix} \frac{1}{4} & 0 \\ 0 & 1 \end{bmatrix}$.



10. (a) Identify T as a vertical expansion from its standard


matrix $A = \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix}$.



12. $T(x, y) = (x + 4y, y)$

(a) Identify T as a horizontal shear from its standard


matrix $A = \begin{bmatrix} 1 & 4 \\ 0 & 1 \end{bmatrix}$.



14. $T(x, y) = (x, 9x + y)$

(a) Identify T as a vertical shear from its matrix

$A = \begin{bmatrix} 1 & 0 \\ 9 & 1 \end{bmatrix}$.



16. The reflection in the x -axis is given by

$T(x, y) = (x, -y)$. If (x, y) is a fixed point, then

$T(x, y) = (x, y) = (x, -y)$ which implies that $y = 0$.

So, the set of fixed points is $\{(t, 0) : t \text{ is real}\}$

18. The reflection in the line $y = -x$ is given by

$T(x, y) = (-y, -x)$. If (x, y) is a fixed point then

$T(x, y) = (x, y) = (-y, -x)$ which implies $-x = y$.

So, the set of fixed points is $\{(t, -t) : t \text{ is real}\}$

20. A horizontal expansion has the standard matrix $\begin{bmatrix} k & 0 \\ 0 & 1 \end{bmatrix}$

where $k > 1$.

A fixed point of T satisfies the equation

$T(\mathbf{v}) = \begin{bmatrix} k & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} kv_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \mathbf{v}$

So the fixed points of T are

$\{\mathbf{v} = (0, t) : t \text{ is a real number}\}$.

22. A vertical shear has the form $T(x, y) = (x, y + kx)$. If

(x, y) is a fixed point, then


$T(x, y) = (x, y) = (x, y + kx)$ which implies that

$x = 0$. So the set of fixed points is $\{(0, t) : t \text{ is real}\}$.

24. Find the image of each vertex under $T(x, y) = (y, x)$.

$T(0, 0) = (0, 0), \quad T(1, 0) = (0, 1),$


$T(1, 1) = (1, 1), \quad T(0, 1) = (1, 0)$



26. Find the image of each vertex under $T(x, y) = \left(x, \frac{y}{4}\right)$.

$T(0, 0) = (0, 0), \quad T(1, 0) = (1, 0),$


$T(1, 1) = \left(1, \frac{1}{4}\right), \quad T(0, 1) = \left(0, \frac{1}{4}\right)$



28. Find the image of each vertex under
- $T(x, y) = (5x, y)$
- .

$$T(0, 0) = (0, 0), \quad T(1, 0) = (5, 0),$$


$$T(1, 1) = (5, 1), \quad T(0, 1) = (0, 1)$$



30. Find the image of each vertex under
- $T(x, y) = (x, y + 3x)$
- .

$$T(0, 0) = (0, 0), \quad T(1, 0) = (1, 3),$$


$$T(1, 1) = (1, 4), \quad T(0, 1) = (0, 1)$$



32. Find the image of each vertex under
- $T(x, y) = (y, x)$
- .

$$T(0, 0) = (0, 0), \quad T(1, 0) = (0, 1),$$


$$T(1, 2) = (2, 1), \quad T(0, 2) = (2, 0)$$



34. Find the image of each vertex under
- $T(x, y) = (x, \frac{1}{2}y)$
- .

$$T(0, 0) = (0, 0), \quad T(1, 0) = (1, 0),$$


$$T(1, 2) = (1, 1), \quad T(0, 2) = (0, 1)$$



36. Find the image of each vertex under
- $T(x, y) = (2x, y)$
- .

$$T(0, 0) = (0, 0), \quad T(1, 0) = (2, 0),$$


$$T(1, 2) = (2, 2), \quad T(0, 2) = (0, 2)$$



38. Find the image of each vertex under
- $T(x, y) = (x, y + 2x)$
- .

$$T(0, 0) = (0, 0), \quad T(1, 0) = (1, 2),$$

$$T(1, 2) = (1, 4), \quad T(0, 2) = (0, 2)$$




40. Find the image of each vertex under
- $T(x, y) = (y, x)$
- .

$$(a) \quad T(0, 0) = (0, 0), \quad T(1, 2) = (2, 1),$$


$$T(3, 6) = (6, 3), \quad T(5, 2) = (2, 5)$$

$$T(6, 0) = (0, 6)$$




$$(b) \quad T(0, 0) = (0, 0), \quad T(0, 6) = (6, 0),$$

$$T(6, 6) = (6, 6), \quad T(6, 0) = (0, 6)$$




42. Find the image of each vertex under $T(x, y) = (x, x + y)$.

(a) $T(0, 0) = (0, 0)$, $T(1, 2) = (1, 3)$, $T(3, 6) = (3, 9)$,
 $T(5, 2) = (5, 7)$, $T(6, 0) = (6, 6)$




(b) $T(0, 0) = (0, 0)$, $T(0, 6) = (0, 6)$,
 $T(6, 6) = (6, 12)$, $T(6, 0) = (6, 6)$




44. Find the image of each vertex under $T(x, y) = (\frac{1}{2}x, 2y)$.

(a) $T(0, 0) = (0, 0)$, $T(1, 2) = (\frac{1}{2}, 4)$,
 $T(3, 6) = (\frac{3}{2}, 12)$, $T(5, 2) = (\frac{5}{2}, 4)$,
 $T(6, 0) = (3, 0)$



(b) $T(0, 0) = (0, 0)$, $T(0, 6) = (0, 12)$,
 $T(6, 6) = (3, 12)$, $T(6, 0) = (3, 0)$



46. The linear transformation defined by A is a vertical shear.

48. The linear transformation defined by A is a vertical contraction.

50. The linear transformation defined by A is a reflection in the y -axis followed by a horizontal contraction.

52. Because $\begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix}$ represents a vertical expansion, and $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ represents a reflection in the line $x = y$, A is a vertical expansion followed by a reflection in the line $x = y$.

54. (a) The linear transformation of $\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$ represents a reflection in the y -axis.

- (b) The linear transformation of $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ represents a reflection in the x -axis.

- (c) The linear transformation of $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ represents a reflection in the line $y = x$.

- (d) The linear transformation of $\begin{bmatrix} k & 0 \\ 0 & 1 \end{bmatrix}$, where $k > 1$, represents a horizontal expansion.

- (e) The linear transformation of $\begin{bmatrix} k & 0 \\ 0 & 1 \end{bmatrix}$, where $0 < k < 1$, represents a horizontal contraction.

- (f) The linear transformation of $\begin{bmatrix} 1 & 0 \\ 0 & k \end{bmatrix}$, where $k > 1$, represents a vertical expansion.

(g) The linear transformation of $\begin{bmatrix} 1 & 0 \\ 0 & k \end{bmatrix}$, where $0 < k < 1$, is represented by a vertical contraction.

(h) The linear transformation of $\begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix}$ represents a horizontal shear.

(i) The linear transformation of $\begin{bmatrix} 1 & 0 \\ k & 1 \end{bmatrix}$ represents a vertical shear.

(j) The linear transformation of $\begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta \\ 0 & \sin \theta & \cos \theta \end{bmatrix}$ represents a rotation about the x -axis.

(k) The linear transformation of $\begin{bmatrix} \cos \theta & 0 & \sin \theta \\ 0 & 1 & 0 \\ -\sin \theta & 0 & \cos \theta \end{bmatrix}$ represents a rotation about the y -axis.

(l) The linear transformation of $\begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$ represents a rotation about the z -axis.

56. A rotation of 60° about the x -axis is given by the matrix

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos 60^\circ & -\sin 60^\circ \\ 0 & \sin 60^\circ & \cos 60^\circ \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ 0 & \frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix}.$$

58. A rotation of 120° about the x -axis is given by the matrix

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos 120^\circ & -\sin 120^\circ \\ 0 & \sin 120^\circ & \cos 120^\circ \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ 0 & \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{bmatrix}.$$

60. Using the matrix obtained in Exercise 56, you find

$$T(1, 1, 1) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ 0 & \frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ \frac{(1 - \sqrt{3})}{2} \\ \frac{(1 + \sqrt{3})}{2} \end{bmatrix}.$$

70. The matrix is $\begin{bmatrix} \cos 60^\circ & 0 & \sin 60^\circ \\ 0 & 1 & 0 \\ -\sin 60^\circ & 0 & \cos 60^\circ \end{bmatrix} \begin{bmatrix} \cos 30^\circ & -\sin 30^\circ & 0 \\ \sin 30^\circ & \cos 30^\circ & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} & 0 & \frac{\sqrt{3}}{2} \\ 0 & 1 & 0 \\ -\frac{\sqrt{3}}{2} & 0 & \frac{1}{2} \end{bmatrix} \begin{bmatrix} \frac{\sqrt{3}}{2} & -\frac{1}{2} & 0 \\ \frac{1}{2} & \frac{\sqrt{3}}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} \frac{\sqrt{3}}{4} & -\frac{1}{4} & \frac{\sqrt{3}}{2} \\ \frac{1}{2} & \frac{\sqrt{3}}{2} & 0 \\ -\frac{3}{4} & \frac{\sqrt{3}}{4} & \frac{1}{2} \end{bmatrix}.$

$$T(1, 1, 1) = \left(\frac{3\sqrt{3} - 1}{4}, \frac{\sqrt{3} + 1}{2}, \frac{\sqrt{3} - 1}{4} \right)$$

62. Using the matrix obtained in Exercise 58, you find

$$T(1, 1, 1) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ 0 & \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ \frac{(-1 - \sqrt{3})}{2} \\ \frac{(-1 + \sqrt{3})}{2} \end{bmatrix}.$$

64. The indicated tetrahedron is produced by a -90° rotation about the z -axis.

66. The indicated tetrahedron is produced by a 180° rotation about the z -axis.

68. The indicated tetrahedron is produced by a 180° rotation about the x -axis.

72. The matrix is

$$\begin{bmatrix} \cos 135^\circ & -\sin 135^\circ & 0 \\ \sin 135^\circ & \cos 135^\circ & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos 120^\circ & -\sin 120^\circ \\ 0 & \sin 120^\circ & \cos 120^\circ \end{bmatrix} = \begin{bmatrix} -\frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} & 0 \\ \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ 0 & \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{bmatrix} = \begin{bmatrix} -\frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{4} & \frac{\sqrt{6}}{4} \\ \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{4} & \frac{\sqrt{6}}{4} \\ 0 & \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{bmatrix}$$

$$T(1, 1) = \left(\frac{\sqrt{6} - \sqrt{2}}{4}, \frac{\sqrt{6} + 3\sqrt{2}}{4}, \frac{\sqrt{3} - 1}{2} \right)$$

Review Exercises for Chapter 6

2. (a) $T(\mathbf{v}) = T(4, -1) = (3, -2)$

(b) $T(v_1, v_2) = (v_1 + v_2, 2v_2) = (8, 4)$

$$v_1 + v_2 = 8$$

$$2v_2 = 4$$

$$v_1 = 6, v_2 = 2$$

Preimage of \mathbf{w} is $(6, 2)$.

4. (a) $T(\mathbf{v}) = T(-2, 1, 2) = (-1, 3, 2)$

(b) $T(v_1, v_2, v_3) = (v_1 + v_2, v_2 + v_3, v_3) = (0, 1, 2)$

$$v_1 + v_2 = 0$$

$$v_2 + v_3 = 1$$

$$v_3 = 2$$

$$v_2 = -1, v_1 = 1$$

Preimage of \mathbf{w} is $(1, -1, 2)$.

6. (a) $T(\mathbf{v}) = T(2, -3) = 7$

(b) The preimage of \mathbf{w} is given by solving the equation

$$T(v_1, v_2) = 2v_1 - v_2 = 4.$$

The resulting linear equation $2v_1 - v_2 = 4$

has the solutions $v_1 = \frac{t+4}{2}$, where t is any real

number. So, the preimage of \mathbf{w} is

$$\left\{ \left(\frac{t+4}{2}, t \right) : t \text{ is any real number} \right\}.$$

8. T preserves addition.

$$\begin{aligned} T(x_1, y_1) + T(x_2, y_2) &= (x_1 + y_1) + (x_2 + y_2) \\ &= (x_1 + x_2) + (y_1 + y_2) \\ &= T(x_1 + x_2, y_1 + y_2) \end{aligned}$$

T preserves scalar multiplication.

$$cT(x, y) = c(x + y) = (cx) + (cy) = T(cx, cy)$$

So, T is a linear transformation with standard matrix $\begin{bmatrix} 1 & 1 \end{bmatrix}$.

10. T does not preserve addition or scalar multiplication, so, T is *not* a linear transformation.

A counterexample is

$$\begin{aligned} T(1, 1) + T(1, 0) &= (4, 1) + (4, 0) \\ &= (8, 1) \neq (5, 1) = T(2, 1). \end{aligned}$$

12. $T(x, y) = (x + y, y)$

$$\begin{aligned} T(x_1, y_1) + T(x_2, y_2) &= (x_1 + y_1, y_1) + (x_2 + y_2, y_2) \\ &= x_1 + y_1 + x_2 + y_2, y_1 + y_2 \\ &= (x_1 + x_2) + (y_1 + y_2), y_1 + y_2 \end{aligned}$$

So, T preserves addition.

$$cT(x, y) = c(x + y, y) = cx + cy, cy = T(cx, cy)$$

So, T preserves scalar multiplication.

So, T is a linear transformation with standard matrix

$$A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}.$$

14. T does not preserve addition or scalar multiplication, and so, T is *not* a linear transformation. A counterexample is

$$\begin{aligned} -2T(3, -3) &= -2(|3|, |-3|) = (-6, -6) \neq (6, 6) \\ &= T(-6, 6) = T(-2(3), -2(-3)). \end{aligned}$$

16. T preserves addition.

$$\begin{aligned}
 T(x_1, x_2, x_3) + T(y_1, y_2, y_3) &= (x_1 - x_2, x_2 - x_3, x_3 - x_1) + (y_1 - y_2, y_2 - y_3, y_3 - y_1) \\
 &= (x_1 - x_2 + y_1 - y_2, x_2 - x_3 + y_2 - y_3, x_3 - x_1 + y_3 - y_1) \\
 &= ((x_1 + y_1) - (x_2 + y_2), (x_2 + y_2) - (x_3 + y_3), (x_3 + y_3) - (x_1 + y_1)) \\
 &= T(x_1 + y_1, x_2 + y_2, x_3 + y_3)
 \end{aligned}$$

T preserves scalar multiplication.

$$\begin{aligned}
 cT(x_1, x_2, x_3) &= c(x_1 - x_2, x_2 - x_3, x_3 - x_1) \\
 &= (c(x_1 - x_2), c(x_2 - x_3), c(x_3 - x_1)) \\
 &= (cx_1 - cx_2, cx_2 - cx_3, cx_3 - cx_1) \\
 &= T(cx_1, cx_2, cx_3)
 \end{aligned}$$

So, T is a linear transformation with standard matrix $A = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ -1 & 0 & 1 \end{bmatrix}$.

18. T preserves addition.

$$\begin{aligned}
 T(x_1, y_1, z_1) + T(x_2, y_2, z_2) &= (x, 0, -y_1) + (x_2, 0, -y_2) \\
 &= (x_1 + x_2, 0, -(y_1 + y_2)) \\
 &= T(x_1 + x_2, y_1 + y_2, z_1 + z_2)
 \end{aligned}$$

T preserves scalar multiplication.

$$cT(x, y, z) = c(x, 0, -y) = (cx, 0, -cy) = T(cx, cy, cz)$$

So, T is a linear transformation with standard matrix $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & -1 & 0 \end{bmatrix}$.

20. Because $(0, 1, 1) = (1, 1, 1) - (1, 0, 0)$, you have

$$\begin{aligned}
 T(0, 1, 1) &= T(1, 1, 1) - T(1, 0, 0) \\
 &= 1 - 3 \\
 &= -2.
 \end{aligned}$$

22. Because $(2, 4) = 2(1, -1) + 3(0, 2)$, you have

$$\begin{aligned}
 T(2, 4) &= 2T(1, -1) + 3T(0, 2) \\
 &= 2(2, -3) + 3(0, 8) \\
 &= (4, -6) + (0, 24) \\
 &= (4, 18).
 \end{aligned}$$

24. (a) Because A is a 2×3 matrix, it maps R^3 into R^2 , ($n = 3, m = 2$).

(b) Because $T(\mathbf{v}) = A\mathbf{v}$ and

$$A\mathbf{v} = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 5 \\ 2 \\ 2 \end{bmatrix} = \begin{bmatrix} 7 \\ 7 \end{bmatrix},$$

it follows that $T(5, 2, 2) = (7, 7)$.

(c) The preimage of \mathbf{w} is given by the solution to the equation $T(v_1, v_2, v_3) = \mathbf{w} = (4, 2)$.

The equivalent system of linear equations

$$\begin{aligned}
 v_1 + 2v_2 - v_3 &= 4 \\
 v_1 + v_3 &= 2
 \end{aligned}$$

has the solution

$$\{(2 - t, 1 + t, t) : t \text{ is a real number}\}.$$

26. (a) Because A is a 2×2 matrix, it maps R^2 into R^2 ($n = 2, m = 2$).

(b) Because $T(\mathbf{v}) = A\mathbf{v}$ and

$$A\mathbf{v} = \begin{bmatrix} 2 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 8 \\ 4 \end{bmatrix} = \begin{bmatrix} 20 \\ 4 \end{bmatrix}, \text{ it follows that}$$

$$T(8, 4) = (20, 4).$$

- (c) The preimage of \mathbf{w} is given by the solution to the equation $T(v_1, v_2) = \mathbf{w} = (5, 2)$.

The equivalent system of linear equations

$$2v_1 + v_2 = 5$$

$$v_2 = 2, v_1 = \frac{3}{2}$$

has the solution $(\frac{3}{2}, 2)$.

28. (a) Because A is a 3×2 matrix, it maps R^2 into R^3 ($n = 2, m = 3$).

(b) Because $T(\mathbf{v}) = A\mathbf{v}$ and

$$A\mathbf{v} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \\ -1 & -3 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} -3 \\ 5 \\ -18 \end{bmatrix}, \text{ it follows that}$$

$$T(3, 5) = (-3, 5, -18).$$

- (c) The preimage of \mathbf{w} is given by the solution to the equation $T(v_1, v_2) = \mathbf{w} = (5, 2, -1)$.

The equivalent system of linear equations

$$-v_1 = 5$$

$$v_2 = 2$$


$$-v_1 - 3v_2 = -1$$

has the solution $v_1 = -5$ and $v_2 = 2$. So, the preimage is $(-5, 2)$.

30. If you translate the vertex $(5, 3)$ back to the origin $(0, 0)$, then the other vertices $(3, 5)$ and $(3, 0)$ are translated to $(-2, 2)$ and $(-2, -3)$, respectively. The rotation of 90° is given by the matrix in Exercise 29, and you have

$$\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} -2 \\ 2 \end{bmatrix} = \begin{bmatrix} -2 \\ -2 \end{bmatrix} \quad \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} -2 \\ -3 \end{bmatrix} = \begin{bmatrix} 3 \\ -2 \end{bmatrix}.$$

Translating back to the original coordinate system, the new vertices are $(5, 3)$, $(3, 1)$ and $(8, 1)$.



32. (a) The standard matrix for T is

$$A = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 1 & 2 \\ 2 & 0 & 1 \end{bmatrix}.$$

Solving $A\mathbf{v} = \mathbf{0}$ yields the solution $\mathbf{v} = \mathbf{0}$. So, $\ker(T) = \{(0, 0, 0)\}$.

- (b) Because $\ker(T)$ is dimension 0, $\text{range}(T)$ must be all of R^3 .

34. (a) The standard matrix for T is

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & -1 \end{bmatrix}.$$

Solving $A\mathbf{v} = \mathbf{0}$ yields the solution

$$\{(t, -t, t) : t \in R\}. \text{ So, } \ker(T) \text{ is } \{(1, -1, 1)\}.$$

- (b) Use Gauss-Jordan elimination to reduce A^T as follows.

$$A^T = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 0 \\ 0 & 1 & -1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix}$$

The nonzero row vectors form a basis for the range of T , $\{(1, 0, 1), (0, 1, -1)\}$.

36. To find the kernel of T , row reduce A .

$$A = \begin{bmatrix} -1 & 2 \\ 0 & -1 \\ -2 & 2 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$$

- (a) $\ker(T) = \{(0, 0)\}$

- (b) $\dim(\ker(T)) = \text{nullity}(T) = 0$

- (c) $A^T = \begin{bmatrix} -1 & 0 & -2 \\ 2 & -1 & 2 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & 2 \end{bmatrix}$

$\text{range}(T)$ is $\text{span}\{(1, 0, 2), (0, 1, 2)\}$.

- (d) $\dim(\text{range}(T)) = \text{rank}(T) = 2$

$$38. A = \begin{bmatrix} 1 & 1 & -1 \\ 1 & 2 & 1 \\ 0 & 1 & 0 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- (a) $\ker(T) = \{(0, 0, 0)\}$

- (b) $\dim(\ker(T)) = \text{nullity}(T) = 0$

- (c) $A^T = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ -1 & 1 & 0 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

$\text{range}(T)$ is $\text{span}\{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$

- (d) $\dim(\text{range}(T)) = 3$

40. $\text{Rank}(T) = \dim P_3 - \text{nullity}(T) = 6 - 4 = 2$

42. $\text{nullity}(T) = \dim(M_{3,3}) - \text{rank}(T) = 9 - 5 = 4$

44. The standard matrix for T is

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Therefore, you have

$$A^2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} = A.$$

46. The standard matrix for T , relative to $B = \{1, x, x^2, x^3\}$, is

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Therefore, you have

$$A^2 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 6 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

48. The standard matrix for T_1 and T_2 are

$$A_1 = \begin{bmatrix} 1 \\ 4 \end{bmatrix} \quad \text{and} \quad A_2 = \begin{bmatrix} 3 & 1 \end{bmatrix}.$$

The standard matrix for $T = T_1 \circ T_2$ is

$$A = A_2 A_1 = \begin{bmatrix} 3 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 4 \end{bmatrix} = \begin{bmatrix} 7 \end{bmatrix}$$

and the standard matrix for $T' = T_2 \circ T_1$ is

$$A' = A_1 A_2 = \begin{bmatrix} 1 \\ 4 \end{bmatrix} \begin{bmatrix} 3 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 12 & 4 \end{bmatrix}.$$

50. The standard matrix for T is

$$A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}.$$

A is invertible and its inverse is given by

$$A^{-1} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}.$$

52. The standard matrix for T is

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & -1 \end{bmatrix}.$$

Because A is *not* invertible, T has no inverse.

54. (a) Because $|A| = 1 \neq 0$, $\ker(T) = \{(0, 0)\}$ and T is one-to-one.

(b) Because $\text{rank}(A) = 2$, T is onto.

(c) The transformation is one-to-one and onto, and is, therefore, invertible.

56. (a) Because $|A| = 40 \neq 0$, $\ker(T) = \{(0, 0, 0)\}$, and T is one-to-one.

(b) Because $\text{rank}(A) = 3$, T is onto.

(c) The transformation is one-to-one and onto, and therefore invertible.

58. (a) The standard matrix for T is

$$A = \begin{bmatrix} 0 & 2 \\ 0 & 0 \end{bmatrix}$$

so it follows that

$$A\mathbf{v} = \begin{bmatrix} 0 & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} -1 \\ 3 \end{bmatrix} = \begin{bmatrix} 6 \\ 0 \end{bmatrix} \Rightarrow T(\mathbf{v}) = (6, 0).$$

(b) The image of each vector in B is as follows.

$$T(2, 1) = (2, 0) = -2(-1, 0) + 0(2, 2)$$

$$T(-1, 0) = (0, 0) = 0(-1, 0) + 0(2, 2)$$

Therefore, the matrix for T relative to B and B' is

$$A' = \begin{bmatrix} -2 & 0 \\ 0 & 0 \end{bmatrix}.$$

Because $\mathbf{v} = (-1, 3) = 3(2, 1) + 7(-1, 0)$,

$$[\mathbf{v}]_B = \begin{bmatrix} 3 \\ 7 \end{bmatrix} \quad \text{and} \quad A'[\mathbf{v}]_B = \begin{bmatrix} -2 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 3 \\ 7 \end{bmatrix} = \begin{bmatrix} -6 \\ 0 \end{bmatrix}.$$

So, $T(\mathbf{v}) = -6(-1, 0) + 0(2, 2) = (6, 0)$.

60. The standard matrix for T is

$$A = \begin{bmatrix} 1 & 3 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & -2 \end{bmatrix}.$$

The transition matrix from B' to B , the standard matrix, is P

$$P = \begin{bmatrix} 1 & 1 & 0 \\ 1 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad P^{-1} = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ \frac{1}{2} & -\frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

The matrix A' for T relative to B' is

$$A' = P^{-1}AP$$

$$\begin{aligned} &= \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ \frac{1}{2} & -\frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & -2 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 \\ 1 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 4 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & -2 \end{bmatrix}. \end{aligned}$$

Because, $A' = P^{-1}AP$, it follows that A and A' are similar.

62. Since $A' = P^{-1}AP$

$$\begin{aligned} &= \begin{bmatrix} 0 & 0 & 1 \\ \frac{1}{2} & 0 & -\frac{1}{2} \\ \frac{1}{2} & -1 & -\frac{1}{2} \end{bmatrix} \begin{bmatrix} 1 & 0 & 1 \\ -1 & 3 & 1 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & 2 & 0 \\ 0 & 1 & -1 \\ 1 & 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 3 \end{bmatrix}, \end{aligned}$$

A and A' are similar.

66. Suppose $\mathbf{b} = \mathbf{0}$. Then $T(\mathbf{v}) = A\mathbf{v}$. $T(\mathbf{u} + \mathbf{v}) = A(\mathbf{u} + \mathbf{v}) = A\mathbf{u} + A\mathbf{v} = T(\mathbf{u}) + T(\mathbf{v})$

$$cT(\mathbf{v}) = c(A\mathbf{v}) = (cA)\mathbf{v} = T(c\mathbf{v})$$

So, $T : R^2 \rightarrow R^2$ is a linear transformation.

Suppose T is a linear transformation. Then $T(\mathbf{u} + \mathbf{v}) = A(\mathbf{u} + \mathbf{v}) + \mathbf{b}$ and $T(\mathbf{u}) + T(\mathbf{v}) = (A\mathbf{u} + \mathbf{b}) + (A\mathbf{v} + \mathbf{b})$.

$$T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$$

$$A(\mathbf{u} + \mathbf{v}) + \mathbf{b} = (A\mathbf{u} + \mathbf{b}) + (A\mathbf{v} + \mathbf{b})$$

$$A\mathbf{u} + A\mathbf{v} + \mathbf{b} = A\mathbf{u} + A\mathbf{v} + 2\mathbf{b}$$

$$\mathbf{b} = 2\mathbf{b}$$

$$\mathbf{0} = \mathbf{b}$$

68. (a) Let $S = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $T = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$.

$$\begin{aligned} \text{Then } S + T &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \text{ and } \text{rank}(S + T) \\ &= \text{rank}(S) + \text{rank}(T). \end{aligned}$$

64. (a) Because $T(\mathbf{v}) = \text{proj}_{\mathbf{u}}\mathbf{v}$ where $\mathbf{u} = (4, 3)$, you have

$$T(\mathbf{v}) = \frac{4x + 3y}{25}(4, 3).$$

So,

$$T(1, 0) = \left(\frac{16}{25}, \frac{12}{25}\right) \text{ and } T(0, 1) = \left(\frac{12}{25}, \frac{9}{25}\right)$$


and the standard matrix for T is

$$A = \frac{1}{25} \begin{bmatrix} 16 & 12 \\ 12 & 9 \end{bmatrix}.$$

$$\begin{aligned} \text{(b) } (I - A)^2 &= \left(\frac{1}{25} \begin{bmatrix} 9 & -12 \\ -12 & 16 \end{bmatrix} \right)^2 \\ &= \frac{1}{25} \begin{bmatrix} 9 & -12 \\ -12 & 16 \end{bmatrix} = I - A. \end{aligned}$$

$$\text{(c) } A\mathbf{v} = \frac{1}{25} \begin{bmatrix} 16 & 12 \\ 12 & 9 \end{bmatrix} \begin{bmatrix} 5 \\ 0 \end{bmatrix} = \begin{bmatrix} \frac{16}{5} \\ \frac{12}{5} \end{bmatrix}$$

$$(I - A)\mathbf{v} = \frac{1}{25} \begin{bmatrix} 9 & -12 \\ -12 & 16 \end{bmatrix} \begin{bmatrix} 5 \\ 0 \end{bmatrix} = \begin{bmatrix} \frac{9}{5} \\ -\frac{12}{5} \end{bmatrix}$$



70. (a) Let $\mathbf{v} \in \text{kernel}(T)$, which implies that $T(\mathbf{v}) = \mathbf{0}$.

Clearly $(S \circ T)(\mathbf{v}) = \mathbf{0}$ as well, which shows that

$$\mathbf{v} \in \text{kernel}(S \circ T).$$

- (b) Let $\mathbf{w} \in W$. Because $S \circ T$ is onto, there exists $\mathbf{v} \in V$ such that $(S \circ T)(\mathbf{v}) = \mathbf{w}$. So,

$$S(T(\mathbf{v})) = \mathbf{w}, \text{ and } S \text{ is onto.}$$

72. Compute the images of the basis vectors under D_x .

$$D_x(1) = 0$$

$$D_x(x) = 1$$

$$D_x(\sin x) = \cos x$$

$$D_x(\cos x) = -\sin x$$

So, the matrix of D_x relative to this basis is

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{bmatrix}.$$

The range of D_x is spanned by $\{x, \sin x, \cos x\}$, whereas the kernel is spanned by $\{1\}$.

74. First compute the effect of T on the basis $\{1, x, x^2, x^3\}$.

$$T(1) = 1$$

$$T(x) = 1 + x$$

$$T(x^2) = 2x + x^2$$


$$T(x^3) = 3x^2 + x^3$$

The standard matrix for T is


$$A = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Because the $\text{rank}(A) = 4$, the $\text{rank}(T) = 4$ and $\text{nullity}(T) = 0$.


76. (a) T is a horizontal shear.



78. (a) T is a horizontal expansion.




80. (a) T is a vertical shear.



82. The image of each vertex is $T(0, 0) = (0, 0)$,


$T(1, 0) = (2, 0)$, $T(0, 1) = (0, 1)$. A sketch of the triangle and its image follows.



84. The image of each vertex is

$$T(0, 0) = (0, 0), T(1, 0) = (1, 2), T(0, 1) = (0, 1).$$

A sketch of the triangle and its image follows.



86. The transformation is a vertical shear $\begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix}$ followed by a vertical expansion $\begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$.

88. A rotation of 90° about the x -axis is given by

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos 90^\circ & -\sin 90^\circ \\ 0 & \sin 90^\circ & \cos 90^\circ \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix}.$$

$$\text{Because } A\mathbf{v} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ -1 \end{bmatrix},$$

the image of $(1, -1, 1)$ is $(1, -1, -1)$.

90. A rotation of 30° about the y -axis is given by

$$A = \begin{bmatrix} \cos 30^\circ & 0 & \sin 30^\circ \\ 0 & 1 & 0 \\ -\sin 30^\circ & 0 & \cos 30^\circ \end{bmatrix} = \begin{bmatrix} \frac{\sqrt{3}}{2} & 0 & \frac{1}{2} \\ 0 & 1 & 0 \\ -\frac{1}{2} & 0 & \frac{\sqrt{3}}{2} \end{bmatrix}.$$

Because

$$A\mathbf{v} = \begin{bmatrix} \frac{\sqrt{3}}{2} & 0 & \frac{1}{2} \\ 0 & 1 & 0 \\ -\frac{1}{2} & 0 & \frac{\sqrt{3}}{2} \end{bmatrix} \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{\sqrt{3}}{2} + \frac{1}{2} \\ -1 \\ -\frac{1}{2} + \frac{\sqrt{3}}{2} \end{bmatrix},$$

the image of $(1, -1, 1)$ is $\left(\frac{\sqrt{3}}{2} + \frac{1}{2}, -1, -\frac{1}{2} + \frac{\sqrt{3}}{2}\right)$.

94. A rotation of 60° about the x -axis is given by

$$A_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos 60^\circ & -\sin 60^\circ \\ 0 & \sin 60^\circ & \cos 60^\circ \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ 0 & \frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix}$$

while a rotation of 60° about the z -axis is given by

$$A_2 = \begin{bmatrix} \cos 60^\circ & -\sin 60^\circ & 0 \\ \sin 60^\circ & \cos 60^\circ & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} & 0 \\ \frac{\sqrt{3}}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

So, the pair of rotations is given by

$$\begin{aligned} A_2 A_1 &= \begin{bmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} & 0 \\ \frac{\sqrt{3}}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ 0 & \frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix} \\ &= \begin{bmatrix} \frac{1}{2} & \frac{\sqrt{3}}{4} & \frac{3}{4} \\ \frac{\sqrt{3}}{2} & \frac{1}{4} & -\frac{\sqrt{3}}{4} \\ 0 & \frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix}. \end{aligned}$$

92. A rotation of 120° about the y -axis is given by

$$A_1 = \begin{bmatrix} \cos 120^\circ & 0 & \sin 120^\circ \\ 0 & 1 & 0 \\ -\sin 120^\circ & 0 & \cos 120^\circ \end{bmatrix} = \begin{bmatrix} -\frac{1}{2} & 0 & \frac{\sqrt{3}}{2} \\ 0 & 1 & 0 \\ \frac{\sqrt{3}}{2} & 0 & -\frac{1}{2} \end{bmatrix}$$

while a rotation of 45° about the z -axis is given by

$$A_2 = \begin{bmatrix} \cos 45^\circ & -\sin 45^\circ & 0 \\ \sin 45^\circ & \cos 45^\circ & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} & 0 \\ \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

So, the pair of rotations is given by

$$\begin{aligned} A_2 A_1 &= \begin{bmatrix} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} & 0 \\ \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} -\frac{1}{2} & 0 & \frac{\sqrt{3}}{2} \\ 0 & 1 & 0 \\ -\frac{\sqrt{3}}{2} & 0 & -\frac{1}{2} \end{bmatrix} \\ &= \begin{bmatrix} -\frac{\sqrt{2}}{4} & -\frac{\sqrt{2}}{2} & \frac{\sqrt{6}}{4} \\ -\frac{\sqrt{2}}{4} & \frac{\sqrt{2}}{2} & \frac{\sqrt{6}}{4} \\ -\frac{\sqrt{3}}{2} & 0 & -\frac{1}{2} \end{bmatrix}. \end{aligned}$$

96. The standard matrix for
- T
- is

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos 90^\circ & -\sin 90^\circ \\ 0 & \sin 90^\circ & \cos 90^\circ \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix}.$$

Therefore, T is given by $T(x, y, z) = (x, -z, y)$. The image of each vertex is as follows.

$$T(0, 0, 0) = (0, 0, 0)$$

$$T(1, 1, 0) = (1, 0, 1)$$

$$T(0, 0, 1) = (0, -1, 0)$$

$$T(1, 1, 1) = (1, -1, 1)$$

$$T(1, 0, 0) = (1, 0, 0)$$

$$T(0, 1, 0) = (0, 0, 1)$$

$$T(1, 0, 1) = (1, -1, 0)$$

$$T(0, 1, 1) = (0, -1, 1)$$

100. (a) True. The statement is true because if T is a reflection $T(x, y) = (x, -y)$, then the standard matrix is

$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

- (b) True. The statement is true because the linear transformation $T(x, y) = (x, ky)$ has the standard matrix.

$$\begin{bmatrix} 1 & 0 \\ 0 & k \end{bmatrix}.$$

102. (a) True. D_x is a linear transformation because it

preserves addition and scalar multiplication. Further, $D_x(P_n) = P_{n-1}$ because for all natural numbers $i \geq 1$,

$$D_x(x^i) = ix^{i-1}.$$

- (b) False. If T is a linear transformation $V \rightarrow W$, then kernel of T is defined to be a set of $\mathbf{v} \in V$, such that $T(\mathbf{v}) = \mathbf{0}_W$.

- (c) True. If $T = T_2 \circ T_1$ and A_i is the standard matrix for T_i , $i = 1, 2$, then the standard matrix for T is equal $A_2 A_1$ by Theorem 6.11 on page 323.

98. The standard matrix for
- T
- is

$$\begin{bmatrix} \cos 120^\circ & -\sin 120^\circ & 0 \\ \sin 120^\circ & \cos 120^\circ & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} & 0 \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Therefore, T is given by

$$T(x, y, z) = \left(-\frac{1}{2}x - \frac{\sqrt{3}}{2}y, \frac{\sqrt{3}}{2}x - \frac{1}{2}y, z \right).$$

The image of each vertex is as follows.

$$T(0, 0, 0) = (0, 0, 0)$$

$$T(1, 0, 0) = \left(-\frac{1}{2}, \frac{\sqrt{3}}{2}, 0 \right)$$

$$T(1, 1, 0) = \left(-\frac{1}{2} - \frac{\sqrt{3}}{2}, \frac{\sqrt{3}}{2} - \frac{1}{2}, 0 \right)$$

$$T(0, 1, 0) = \left(-\frac{\sqrt{3}}{2}, -\frac{1}{2}, 0 \right)$$

$$T(0, 0, 1) = (0, 0, 1)$$


$$T(1, 0, 1) = \left(-\frac{1}{2}, \frac{\sqrt{3}}{2}, 1 \right)$$

$$T(1, 1, 1) = \left(-\frac{1}{2} - \frac{\sqrt{3}}{2}, \frac{\sqrt{3}}{2} - \frac{1}{2}, 1 \right)$$


$$T(0, 1, 1) = \left(-\frac{\sqrt{3}}{2}, -\frac{1}{2}, 1 \right)$$

Project Solutions for Chapter 6

1 Reflections in the Plane-I




1.




$$\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$$

2.



$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

3.



$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

4. $\mathbf{v} = (2, 1)$ $B = \{\mathbf{v}, \mathbf{w}\}$

$$\mathbf{w} = (-1, 2)$$

$$L(\mathbf{v}) = \mathbf{v} \quad \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} = A$$

$$L(\mathbf{w}) = -\mathbf{w} \quad \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} = A$$

$$B' = \{\mathbf{e}_1, \mathbf{e}_2\} \text{ standard basis}$$

A is a matrix of L relative to basis B .

$A' = P^{-1}AP$ matrix of L relative to the standard basis B' .


$$[B' : B] \rightarrow [I : P^{-1}] \Rightarrow P^{-1} = \begin{bmatrix} 2 & -1 \\ 1 & 2 \end{bmatrix} \Rightarrow P = \frac{1}{5} \begin{bmatrix} 2 & 1 \\ -1 & 2 \end{bmatrix}$$

$$A' = P^{-1}AP = \frac{1}{5} \begin{bmatrix} 2 & -1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ -1 & 2 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 2 & 1 \\ 1 & -2 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ -1 & 2 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 3 & 4 \\ 4 & -3 \end{bmatrix} = \begin{bmatrix} \frac{3}{5} & \frac{4}{5} \\ \frac{4}{5} & -\frac{3}{5} \end{bmatrix}$$

$$\begin{bmatrix} \frac{3}{5} & \frac{4}{5} \\ \frac{4}{5} & -\frac{3}{5} \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} \frac{3}{5} & \frac{4}{5} \\ \frac{4}{5} & -\frac{3}{5} \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \end{bmatrix}$$

$$\begin{bmatrix} \frac{3}{5} & \frac{4}{5} \\ \frac{4}{5} & -\frac{3}{5} \end{bmatrix} \begin{bmatrix} 5 \\ 0 \end{bmatrix} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$$




$$\begin{aligned} 5. \quad \mathbf{v} &= (-b, a) \\ \mathbf{w} &= (a, b) \end{aligned}$$

$$A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

$$P^{-1} = \begin{bmatrix} -b & a \\ a & b \end{bmatrix}$$

$$P = \frac{1}{a^2 + b^2} \begin{bmatrix} -b & +a \\ +a & +b \end{bmatrix}$$

$$A' = P^{-1}AP = \begin{bmatrix} -b & a \\ a & b \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} P = \begin{bmatrix} -b & -a \\ a & -b \end{bmatrix} \begin{bmatrix} -b & a \\ a & b \end{bmatrix} \frac{1}{a^2 + b^2} = \frac{1}{a^2 + b^2} \begin{bmatrix} b^2 - a^2 & -2ab \\ -2ab & a^2 - b^2 \end{bmatrix}$$



$$6. \quad 3x + 4y = 0 \quad A' = \frac{1}{3^2 + 4^2} \begin{bmatrix} 7 & -24 \\ -24 & -7 \end{bmatrix}$$

$$\frac{1}{25} \begin{bmatrix} 7 & -24 \\ -24 & -7 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \end{bmatrix} = \frac{1}{25} \begin{bmatrix} -75 \\ -100 \end{bmatrix} = \begin{bmatrix} -3 \\ -4 \end{bmatrix}$$

$$\frac{1}{25} \begin{bmatrix} 7 & -24 \\ -24 & -7 \end{bmatrix} \begin{bmatrix} -4 \\ 3 \end{bmatrix} = \frac{1}{25} \begin{bmatrix} -100 \\ 75 \end{bmatrix} = \begin{bmatrix} -4 \\ 3 \end{bmatrix}$$

$$\frac{1}{25} \begin{bmatrix} 7 & -24 \\ -24 & -7 \end{bmatrix} \begin{bmatrix} 0 \\ 5 \end{bmatrix} = \frac{1}{25} \begin{bmatrix} -24 \cdot 5 \\ -7 \cdot 5 \end{bmatrix} = \begin{bmatrix} -\frac{24}{5} \\ -\frac{7}{5} \end{bmatrix}$$

2 Reflections in the Plane-II

$$1. \quad \mathbf{v} = (0, 1) \quad \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$$


$$2. \quad \mathbf{v} = (1, 0) \quad \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

$$3. \quad \mathbf{v} = (2, 1) \quad B = \{\mathbf{v}, \mathbf{w}\}$$

$$\mathbf{w} = (-1, 2)$$

$$\left. \begin{aligned} \text{proj}_{\mathbf{v}} \mathbf{v} &= \mathbf{v} \\ \text{proj}_{\mathbf{v}} \mathbf{w} &= \mathbf{0} \end{aligned} \right\} \quad A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

$$P^{-1} = \begin{bmatrix} 2 & -1 \\ 1 & 2 \end{bmatrix}, \quad P = \frac{1}{5} \begin{bmatrix} 2 & 1 \\ -1 & 2 \end{bmatrix}$$




$$A' = P^{-1}AP = \text{matrix of } L \text{ relative to standard basis.}$$

$$= \begin{bmatrix} 2 & -1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} P = \begin{bmatrix} 2 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ -1 & 2 \end{bmatrix} \frac{1}{5} = \begin{bmatrix} \frac{4}{5} & \frac{2}{5} \\ \frac{2}{5} & \frac{1}{5} \end{bmatrix}$$

$$\begin{bmatrix} \frac{4}{5} & \frac{2}{5} \\ \frac{2}{5} & \frac{1}{5} \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad \begin{bmatrix} \frac{4}{5} & \frac{2}{5} \\ \frac{2}{5} & \frac{1}{5} \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} \frac{4}{5} & \frac{2}{5} \\ \frac{2}{5} & \frac{1}{5} \end{bmatrix} \begin{bmatrix} 5 \\ 0 \end{bmatrix} = \begin{bmatrix} 4 \\ 2 \end{bmatrix}$$



$$4. \quad \mathbf{v} = (-b, a) \quad A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\mathbf{w} = (a, b)$$

$$P^{-1} = \begin{bmatrix} -b & a \\ a & b \end{bmatrix} \quad P = \frac{1}{a^2 + b^2} \begin{bmatrix} -b & a \\ a & b \end{bmatrix}$$

$$A' = P^{-1}AP = \frac{1}{a^2 + b^2} \begin{bmatrix} b^2 & -ab \\ -ab & a^2 \end{bmatrix}$$

$$5. \quad \text{proj}_{\mathbf{v}} \mathbf{u} = \frac{1}{2}(\mathbf{u} + L(\mathbf{u})) \Rightarrow L(\mathbf{u}) = 2\text{proj}_{\mathbf{v}} \mathbf{u} - \mathbf{u}$$

$$L = 2 \text{proj}_{\mathbf{v}} - I$$

$$= 2 \frac{1}{a^2 + b^2} \begin{bmatrix} b^2 & -ab \\ -ab & a^2 \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$= \frac{1}{a^2 + b^2} \left(\begin{bmatrix} 2b^2 & -2ab \\ -2ab & 2a^2 \end{bmatrix} - \begin{bmatrix} a^2 + b^2 & 0 \\ 0 & a^2 + b^2 \end{bmatrix} \right)$$

$$= \frac{1}{a^2 + b^2} \begin{bmatrix} b^2 - a^2 & -2ab \\ -2ab & a^2 - b^2 \end{bmatrix}$$

