

CLASS NOTES
FOR
ELEMENTARY STATISTICS
LETU MATH-1423

Nicholas Capo

September 5, 2012

Contents

1	Introduction	5
1.1	Data	5
1.1.1	Data Sets	5
1.1.2	Types of Data	5
1.2	Sample Mean and Median	5
1.2.1	Definition	5
1.3	Measures of Variability	6
1.3.1	Standard Deviation	6
1.4	Descriptive Statistics	6
1.4.1	Quartiles	6
1.4.2	Range and Interquartile Range	6
1.4.3	Box and Whisker Plot	7
1.5	Stem and Leaf Plots	7
1.5.1	Key Notation	8
1.5.2	Double Stem and Leaf	8
1.6	Frequency Distribution	8
1.6.1	Midpoint of a Class	8
1.6.2	Relative Frequency	8
1.7	Scatter Plots	8
1.8	Homework	8

This document comprises classroom notes from Statistics Class at [LeTourneau University](#), in the Fall of 2012.

Although the author will attempt to be complete and correct in these notes, it is the readers responsibility to learn and understand the material. The author assumes no responsibility for the completeness or accuracy of this content.

The latest version of this document is available at <https://bitbucket.org/nicholascapo/StasisticsNotes/src/tip/StatisticsNotes.pdf>

Chapter 1

Introduction

Definition of Statistics

“Statistics is the science of collecting, organizing, analyzing, and interpreting data in order to make decisions.”

1.1 Data

1.1.1 Data Sets

Population The collection of all outcomes, responses, measurements, or counts, that are of interest.

Sample A subset of the population.

Parameter A number that describes a population characteristic.

Statistic A number that describes a sample characteristic.

1.1.2 Types of Data

Qualitative Data Attributes, labels, or non-numerical entries.

Quantitative Data Numerical measurements or counts.

1.2 Sample Mean and Median

1.2.1 Definition

Sample Mean The average of the sample data points, however it may not be a data point.

$$\bar{x} = \sum_{i=1}^n \frac{x_i}{n} = \frac{x_1 + x_2 + x_3 \cdots x_n}{n}$$

Sample Median The middle value of the data.

$$\tilde{x} = \begin{cases} x_{(\frac{n+1}{2})} & \text{if } n \text{ is odd} \\ \frac{1}{2}(x_{\frac{n}{2}} + x_{\frac{n}{2}+1}) & \text{if } n \text{ is even} \end{cases}$$

Trimmed Mean A trimmed mean is computed by trimming off the largest and smallest set of values. For example a 10% trimmed mean is found by eliminating the largest 10% and smallest 10% and computing the mean of the remaining values. This may be useful for data that contains possible outliers. Denoted by $x_{tr(\text{percent})}$

1.3 Measures of Variability

1.3.1 Standard Deviation

Sample Variance

$$s^2 = \sum_{i=1}^n \frac{(x_i - \bar{x})^2}{n-1}$$

Sample Standard Deviation

$$s = +\sqrt{s^2}$$

The standard deviation is 0 when all the data points are the same.

1.4 Descriptive Statistics

1.4.1 Quartiles

Quartiles approximately divide an ordered data set into four equal parts.

First Quartile, Q_1 About 25% of the data fall on or below Q_1

Second Quartile, Q_2 About 50% of the data fall on or below Q_2

Third Quartile, Q_3 About 75% of the data fall on or below Q_3

1.4.2 Range and Interquartile Range

Range

$$\text{range} = \text{max value} - \text{min value}$$

Interquartile Range

$$IQR = Q_3 - Q_1$$

To help find outliers, compute $1.5 \times IQR$, and any values that lie outside the interval $[Q_1 - 1.5 \times IQR, Q_3 + 1.5 \times IQR]$ is a possible (and probable) outlier.

1.4.3 Box and Whisker Plot

Exploratory Data Analysis Tool

- Requires
 - Min
 - Q_1
 - Median
 - Q_3
 - Max

Example

Example Data	[1, 2, 3, 4, 5, 6, 11]
Min	1
Median	4.0
Max	6
Outlier	11




Figure 1.1: Example Box And Whisker Plot

1.5 Stem and Leaf Plots

These look like a sideways histogram

Data: [31, 21, 32, 33, 41, 42, 58, 25, 21]

Stem	Leaf	Key: $a b = ab$
2	1,1,5	
3	1,2,3	
4	1,2	
5	8	

1.5.1 Key Notation

Key: 4—5 = 45 Key: 4—5 = 4.5

1.5.2 Double Stem and Leaf

Separate the leaves into two groups, (0-4, and 5-9)

Data: [31, 21, 32, 33, 41, 42, 58, 25, 21]

Stem	Leaf	Key: $a b = ab$
2	1,1	
2	5	
3	1,2,3	
4	1.2	
4		
5		
5	8	

1.6 Frequency Distribution

A table that shows classes or intervals of data with a count of the number of entries in each class.

1.6.1 Midpoint of a Class

Average of the class limits.

$$\frac{(\text{lower class limit}) + (\text{upper class limit})}{2}$$

1.6.2 Relative Frequency

$$\frac{\text{class frequency}}{\text{sample size}} = \frac{f}{n}$$

1.7 Scatter Plots

Each entry in one data set corresponds to one entry in a second set, one-to-one mapping.

1.8 Homework

- p. 13 #'s 1.5, 1.6
- p. 17 #'s 1.11, 1.12
- p. 31 #'s 1.18, 1.19, 1.20, 1.29, 1.30

Bibliography

- [1] R.E. Walpole et al. *Probability and Statistics for Engineers and Scientists*. Pearson Education, 2010. ISBN: 9780321629111. URL: <http://books.google.com/books?id=tzZxRQAACAAJ>.