1	Neither kin selection nor familiarity explain affiliative biases towards maternal siblings in wild
2	mountain gorillas
3	
4	Nicholas M. Grebe ¹ , Jean Paul Hirwa ² , Tara S. Stoinski ² , Linda Vigilant ³ , & Stacy Rosenbaum ¹
5	
6	¹ Department of Anthropology, University of Michigan
7	² Dian Fossey Gorilla Fund International
8	³ Department of Primatology, Max Planck Institute for Evolutionary Anthropology
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	

Abstract

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

Evolutionary theories predict that sibling relationships will reflect a complex balance of cooperative and competitive dynamics. In most mammals, dispersal and death patterns mean that sibling relationships occur in a relatively narrow window during development, and/or only with same-sex individuals. Besides humans, one notable exception are mountain gorillas, in which nonsex biased dispersal, relatively stable group composition, and the long reproductive tenures of alpha males mean that animals routinely reside with both same and opposite-sex, and full and half siblings, throughout their lives. Using nearly 40,000 hours of observation data collected over 14 years on 699 sibling and 1258 non-sibling pairs of wild mountain gorillas, we demonstrate that individuals have strong affiliative preferences for full and maternal siblings over paternal siblings or unrelated animals, consistent with an inability to discriminate paternal kin. Intriguingly, however, aggression data imply the opposite. Aggression rates were statistically indistinguishable among all types of dyads except one: in mixed-sex dyads, non-siblings engaged in substantially more aggression than siblings of any type. This pattern suggests mountain gorillas may be capable of distinguishing paternal kin, but nonetheless choose not to affiliate with them over non-kin. A preference for maternal kin occurs despite gorillas not possessing low male reproductive skew, the key characteristic believed to underlie such biases. These results call into question reasons for strong maternal kin biases when paternal kin are identifiable, familiar, and similarly likely to be long-term social partners, and suggest behavioral mismatches at play during a transitional period in mountain gorilla society.

Key words: paternal kin, maternal kin, kin selection, kin recognition, primatology

Introduction

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

In humans and non-human animals alike, sibling social relationships are marked by continuous dynamics of conflict and cooperation that begin before birth (Trivers, 1974), and can persist throughout an entire lifespan, with important fitness consequences for the individuals involved (Lu, 2007; Hudson & Trillmich, 2008; Nitsch et al., 2013). While classical frameworks of sibling interactions emphasized competition among brood- or litter-mates for limited parental resources during times of dependency (e.g., Mock & Parker, 1997), subsequent developments across numerous academic disciplines (demography: e.g. Sear & Mace, 2008; Nitsch et al., 2013; sociology: e.g. Steelman et al., 2002; Lu, 2007; behavioral ecology: e.g. Silk, 2002; Hudson & Trillmich, 2008; developmental psychology: e.g. Lamb et al., 2014) have explored the full arc of sibling competition and cooperation across the lifespan and demonstrated the complexity and diversity inherent to sibling relationships. In understanding the evolution of human sibling relationships in particular, comparative studies of our primate cousins have identified a number of factors predictive of how siblings cooperate and compete. Inconsistent results within and between species, along with the remarkable flexibility of human social systems, however, limits the translational value of many primate models. Here, we address these gaps by presenting an extensive longitudinal study of wild mountain gorillas (Gorilla beringei beringei), an endangered ape whose unique, flexible social structure serves as a valuable comparative model to humans. Classic models of kin selection predict that the social/mating structure of animal groups creates patterns of relatedness between group members, which then selects for kin recognition mechanisms that manifest in differences in cooperative and/or affiliative behavior (Hamilton, 1964; Grafen, 1990; Mateo, 2015). This straightforward idea has spawned a large body of work on kin discrimination in primates, with notably mixed results. Some studies support the existence of

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

sophisticated kin discrimination (Wu et al., 1980; Smith et al., 2003; Widdig et al., 2002, Pfefferle et al., 2014), while others do not, instead favoring simple familiarity as the primary determinant of interaction patterns (Fredrickson & Sackett, 1984; Schaub, 1996; Erhart et al., 1997; Wikberg et al., 2014; Godoy et al., 2016). Inconsistent evidence has led some to suggest that non-monogamous primates evince matrilineal, but not patrilineal, sibling kin discrimination (Mitani et al., 2000; Chapais, 2001; Langergraber et al., 2007). Yet other perspectives challenge this clean distinction, suggesting that complex interactions between familiarity and kin discrimination structure social bonds across primates (see e.g. Silk, 2002; Widdig et al., 2002; Silk, 2009; Lynch et al., 2017). As one of the main contributors to familiarity, age differences within sibling and non-sibling dyads might influence social dynamics. On one hand, siblings close in age might be more likely to compete for limited parental resources (Tung et al., 2016; Salmon & Hehman, 2021); on the other hand, as longer-lasting co-residents within the same family environment, they might also be expected to form stronger affiliative bonds than siblings distant in age (though, again, this may not apply equally to maternal and paternal sibships; Widdig et al., 2002). It is unclear to what extent age proximity effects are restricted to genetic relatives. Female rhesus macaques appear to bias affiliation towards similarly-aged peers, even when unrelated to them (Widdig et al., 2001). Among female baboons, even in individuals not related through the matriline, dyadic bond strength weakened with increasing age differences; however, when restricting analyses to females unrelated through both the matriline and patriline, effects of age differences attenuated sharply (Smith et al., 2003; Silk et al., 2006). These results once again imply social familiarity (as indexed by age differences) and kin discrimination are both important for predicting sibling relationship qualities (Godoy et al., 2016), though their additive and/or interactive effects remain poorly defined.

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

Finally, the sex makeup of the dyad might influence interaction styles due to the differential benefits males and females receive from interactions with brothers, sisters, and unrelated partners (e.g. Lonsdorf et al., 2018). From the perspective of males, especially in species who engage in aggressive intrasexual competition, other males, brothers included, can represent important allies (e.g. Meikle & Vessey, 1981; Goodall, 1986) or rivals (Daly & Wilson, 1988) during status-striving efforts in adulthood. In either case, assessing physical capacities or formidability would aid in these efforts. Accordingly, rough-and-tumble play between males might serve important functions as a rehearsal for intrasexual competition in adulthood (Gray, 2019), suggesting such a behavior should occur most often in male-male relationships—a prediction supported by research on male-dominant primates (e.g., Brown & Dixson, 2000; Maestripieri & Ross, 2004). While male-male interaction patterns might generally differ from those of other sex configurations, these differences may themselves partially depend on kinship: in chimpanzees, some evidence suggests that fraternal relationships among adolescents and adults are more affiliative and cooperative than relationships between unrelated males (e.g. Mitani, 2009; Sandel et al., 2020). From the female perspective, evidence for fraternal influences on fitness outcomes is mixed: some research suggests no effect, except perhaps in the case of maternal death (Engh et al., 2009), while one demographic study of humans reports benefits of older brothers on women's lifetime fitness (Nitsch et al., 2013). From the perspective of both males and females, sisters may represent important future alloparental helpers, either for the individual themselves (e.g. Hamilton et al., 1982; Gould, 2000; Hobaiter et al., 2014), or the individual's offspring (e.g. Johnson et al., 1980; Nishida, 1983; but see Silk et al., 2006), and thus cultivating relationships with sisters via affiliative interactions might be beneficial for both males and females. These kinds of sex-biased interactions might additionally depend on age differences between siblings (Lonsdorf et al., 2018). Lastly, for females in particular, sororal

relationships may exert important influences on future rank and resource acquisition outcomes (Charpentier et al., 2008; Lea et al., 2014; cf. Engh et al., 2009).

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

Addressing the issues reviewed above, and understanding the nature and evolution of cooperative social relationships in primates more generally, requires long-term investigations of species that reveal how individuals respond behaviorally to socioecological variation (e.g. Alberts & Altmann, 2012). With this principle in mind, mountain gorillas in particular are a compelling comparative model for the evolution and development of human sibling relationships. Long-term monitoring of wild mountain gorillas by the Dian Fossey Gorilla Fund has revealed social structures marked by extensive diversity in relatedness, age proximity, and sex makeup infrequently observed in other primate groups (Robbins et al., 2009; Roy et al., 2014). Mountain gorillas regularly form multifemale, single-male groups, as well as multi-female, multi-male groups in which multiple males reproduce, though paternity data and unsophisticated paternal kin discrimination mechanisms are consistent with historically high reproductive skew (Bradley et al., 2005; Rosenbaum et al., 2015; Vigilant et al., 2015). As a result of this structure, co-resident offspring have a reasonable chance of being full siblings, paternal half-siblings, maternal half-siblings, or unrelated to one another. Like humans, both males and females, upon reaching maturity, may opt to disperse or remain in their natal groups (Robbins et al., 2009; Stoinski et al., 2009), which permits fraternal, sororal, and mixedsex relationships that can last for an entire lifespan.

In the present study, we use nearly 40,000 hours of behavioral data spanning 14 years to describe patterns of interactions between siblings and demographically-comparable non-sibling dyads in social groups of wild mountain gorillas. Using extensive maternity and genetic paternity data available for 157 identifiable individuals, we examine whether full siblings, maternal half-siblings, paternal half-siblings, and unrelated co-residents exhibit differing patterns of affiliation

(playing, grooming, and time spent in close proximity) and agonism (contact and non-contact aggression) in line with models of kin selection, after adjusting for the potential mediating presence of mothers in these interactions. Evidence of relatively weak kin discrimination among gorilla fathers and offspring (Rosenbaum et al., 2015, though see also Vigilant et al., 2015 for evidence of father-daughter inbreeding avoidance) might suggest correspondingly weak social bias toward paternal half-siblings, along with strong bias toward full siblings and maternal half-siblings; we test this speculation for the first time in this species. We also investigate whether familiarity, as indexed by age differences between social partners, predict patterns of affiliative or agonistic behavior, and whether these patterns differ between kin categories—particularly paternal and maternal kin (as some evidence from cercopithecine monkeys suggests; e.g. Widdig et al., 2002).

Lastly, we explore sex makeup as a third axis of variation potentially relevant for understanding sibling and co-resident social relationships. We test whether male-male, female-female, and mixed-sex sibling relationships are characterized by differing rates and types of social interactions, and whether these sex category differences are restricted to kin. Via these comparisons, we ask whether differences are consistent with the kinds of benefits siblings might be expected to deliver later in life: for example, among males, are fraternal relationships marked by higher rates of playing and fighting, and sororal relationships higher rates of grooming?; among females, are sororal relationships marked by the highest rates of grooming compared to any other dyad configuration?; are affiliative patterns unique to siblings, or are comparable trends found in unrelated dyads?

Results

Affiliative Behaviors. In our full sample of 1957 unique dyads spanning 7,858 dyad-years, full siblings (n=43 dyads) played and groomed each other significantly more than did paternal siblings (n=555 dyads) or non-siblings (n=1258 dyads; all comparisons p < 0.001; Figure 1A, 1B). Maternal

siblings (n=101 dyads) played significantly less than full siblings, but groomed at comparable rates. Age differences (in our sample, mean: 5.90 years; SD: 4.57 years; range: 0-23.5 years) interacted with relatedness in predicting grooming (p=0.023), but not play (p=0.076). Play consistently dropped for siblings and non-siblings alike as age differences increased (γ ranging from -0.23 – -0.28, all p < 0.001; Figure 2A). By contrast, grooming rates were relatively unrelated to age differences between partners (γ ranging from -0.07 – 0.02, all p > 0.05; Figure 2B).

Figure 1. Box and dot plots comparing relatedness categories (A, B) and sex categories (C, D) for play rates (left) and grooming rates (right).






Figure 2. Estimated rates of play (A) and grooming (B) across a range of age differences, separated by relatedness category. Bars represent 95% confidence intervals for rates of behavior at a given age difference.




Male-male dyads (n=520) played more than either mixed-sex (n=981) or female-female dyads (n=456); conversely, female-female dyads groomed each other more than either mixed-sex or male-male dyads (all p < 0.001; Figure 1C, 1D). These patterns too were significantly moderated by age differences (p < 0.002), but not relatedness (p = 0.078 and 0.112). Play dropped rapidly with increasing age differences ($\gamma = -0.29 - -0.21$) for all sex configurations (all p < 0.001; Figure 3A). Grooming was steadily low in male-male and mixed-sex dyads ($\gamma = -0.01$ and -0.02, p > 0.45), though it dropped with increasing age differences in female-female dyads ($\gamma = -0.09$, $\gamma = 0.001$), such that differences between sex categories became indistinguishable after approximately a 10-year age difference (Figure 3B).

Figure 3. Estimated rates of play (A) and grooming (B) across a range of age differences, separated by sex category. Bars represent 95% confidence intervals for rates of behavior at a given age difference.



Time spent in proximity. The time dyads spent in close proximity (<2 m) with each other also varied between relatedness categories (p < 0.001), with maternal siblings and full siblings once again spending more time near each other than non-siblings, who themselves spent more time in close proximity than paternal siblings did (all comparisons p < 0.001; Figure 4A). However, these patterns too were moderated by age differences (p < 0.001). Proximity decreased with increasing age differences in maternal siblings and paternal siblings ($\gamma = -0.08$ and -0.09, p < 0.001), but did not decrease significantly in full siblings or non-siblings ($\gamma = -0.06$ and 0.01, p > 0.19). Thus, while all classes of siblings spent more time near each other than non-siblings when near in age, even when adjusting for their mother's presence, this distinction was partially reversed at large age differences, when paternal siblings spent much less time near each other than any other dyad category (Figure 4B).


Figure 4. Box and dot plots (A) and estimated trends across a range of age differences (B) for the time gorilla dyads spent in close proximity, separated by relatedness category. Bars in (B) represent 95% confidence intervals for rates of proximity at a given age difference.



Competitive Behaviors. Neither relatedness nor sex category on their own significantly predicted rates of aggressive behavior (p = 0.205 and 0.763, respectively). However, our model did reveal a significant sex makeup × relatedness interaction term (p = 0.049; Figure 5A). Decomposing this interaction, among female-female and male-male dyads, there were no statistically significant contrasts between relatedness categories. In mixed-sex dyads, non-siblings engaged in substantially more aggression than any sibling category (all p < 0.031).

Dyads more distant in age engaged in less aggression than dyads that were closer in age, regardless of relatedness or sex makeup ($\chi = -2.11$, p = 0.035; Figure 5B).

Figure 5. Box and dot plots (A) and estimated trends across a range of age differences (B) for aggression within gorilla dyads, separated by relatedness and sex category. Bars in (B) represent 95% confidence intervals for rates of aggression at a given age difference.



Discussion

In a comprehensive examination of dyadic mountain gorilla social relationships spanning 14 years and nearly 40,000 hours of observation, we find complex patterns of affiliation and competition within gorilla pairs that speak to sex-, age-, and relatedness-specific social biases. In general, siblings affiliated with each other more and spent more time together than non-siblings, even when accounting for the presence of mothers. But within siblings, affiliative patterns further varied: full and maternal siblings were in most cases much more affiliative than paternal siblings, whose behavior more closely resembled that of non-siblings. We consistently observed a trend for male-male dyads to play more, for female-female dyads to groom more, and for mixed-sex dyads to fall intermediate between these groups. Examining competitive behaviors, on the other hand, revealed a more narrow sibling bias. Aggression was most common in mixed-sex non-sibling dyads, and larger age differences similarly predicted less aggression across all dyad types.

At the broadest level, our results support the existence of affiliative biases towards kin in mountain gorillas. While past research has been largely equivocal about the extent of kin discrimination that relies on mechanisms beyond familiarity (Wikberg et al., 2014; Godoy et al., 2016; Lynch et al., 2017)—and indeed, our results do support a role of familiarity in structuring social interactions—our findings are unlikely to be entirely explained by mere exposure for at least three reasons:

First, gorilla social groups are tight-knit and cohesive compared to their close ape relatives (Goodall, 1986; Remis, 1997; Morrison et al., 2021), such that all individuals in a group, related or not, are very likely quite familiar with one another. Second, our results pertaining to prosocial biases towards siblings are not fully explained by familiarity or exposure, as indexed by age differences (Widdig et al., 2001; Smith et al., 2003; Lynch et al., 2017). We observed clear biases towards kin at all but the largest age differences—and some siblings in our data set were 20 or more years apart in

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

age—even though sibling and non-sibling age-mates in the same social group would typically be expected to possess close familiarity. Finally, while mothers undoubtedly mediate social interactions of offspring, especially for immature individuals, biases towards siblings persist even when adjusting for the frequency of her presence during interaction periods. Jointly, these considerations suggest a sibling bias in mountain gorillas subject to influence, but not determination, by demographic factors, which we interpret as aiding in the development of sibling relationships that exist across timescales rarely observed in other non-human primates.

The observation that full and maternal siblings groomed, played, and spent more time near each other than paternal siblings or non-siblings, who tended to affiliate at comparable rates, further suggests that mountain gorillas, like several other primate species studied (Langergraber et al., 2007; Silk et al., 2006; Lynch et al., 2017), evince much stronger maternal than paternal kin bias (see also Rosenbaum et al., 2015, which found little evidence for paternal kin discrimination among fathers and offspring). Interestingly, this "asymmetric bias" in affiliation seems to persist despite mountain gorillas lacking a key element of the social structure hypothesized to underlie it in other species: namely, low reproductive skew (Galezo et al., 2022). Thus, one question concerns why mountain gorillas do not appear to more strongly favor paternal siblings. Current evidence indicates that single-male gorilla groups across research sites are entirely genetically polygynous (reviewed in Rosenbaum & Silk, 2022), and while there can be considerable temporal variation, reproductive skew is generally much higher in multi-male gorilla groups than in, for example, chimpanzee, savannah baboon, or rhesus macaque groups (Vigilant et al. 2015, Surbeck et al. 2017, Alberts et al. 2003, Widdig et al. 2004). We propose that, despite possessing a mating system quite unlike these other primate species, mountain gorillas still exhibit a comparable maternal sibling bias due to a mismatch between their historical mating structure—which we speculate consisted of highly polygynous one-male units-and their contemporary social structure of tight-knit, often multi-male

groups. In other words, while individuals in highly polygynandgrous groups might find it too difficult to detect and adjust affiliation toward paternal kin, perhaps mountain gorillas fail to do so because, until very recently, it was unnecessary. If co-residency was enough to identify patneral kin with reasonable accuracy, a more sophisticated recognition mechanism would be unlikely to evolve.

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

Notably, while we see little evidence that mountain gorillas show a prosocial bias towards paternal siblings, patterns of aggressive behavior suggest there may still be kin recognition mechanisms at play for all sibling types. Aggression remained low across most combinations of relatedness and sex configurations, with one exception: mixed-sex interactions among non-siblings. This pattern is consistent with males deploying aggression in the context of mate attraction or coercion. Past research in gorillas suggests male aggression towards females may have a number of non-mutually exclusive functions within those domains: to police female-female aggression, to discourage female dispersal or mate choice, or to indicate protective ability or overall condition (Robbins, 2009; Breuer et al., 2016). The fact that this kind of aggression was observed less frequently among related male-female pairs is another observation consistent with accurate kin discrimination. It also suggests active inbreeding avoidance, to the extent that aggression truly serves a mate attraction function. While death and dispersal have been suggested to obviate the need for sophisticated inbreeding avoidance mechanisms in some primates (e.g. baboons; Galezo et al., 2022), such an explanation is unlikely to apply to contemporary mountain gorillas. Living with opposite-sex relatives after sexual maturity is a routine occurrence in this species. Prior research confirms strong inbreeding avoidance between father-daughter dyads in this species (Vigilant et al. 2015), but further work is needed to investigate the extent to which male mate choice is manifested via female-directed aggression, and whether females, for their part, possess additional mechanisms to avoid mating with kin, including paternal siblings.

Together, these observations—prosocial biases towards kin that do not appear to be fully explained by familiarity; a stronger maternal than paternal sibling prosocial bias; and avoidance of intersexual aggression across all sibling types—both speak to key questions about the development of great ape sibling relationships and present two additional puzzles for interpretation. First, traditional mechanistic explanations for sibling biases that typically appeal to exposure during developmental periods appear largely inconsistent with our results and the nature of mountain gorilla sociality, in which siblings and non-siblings, and maternal and paternal siblings, are all likely to have significant exposure to one another during development. It is possible that early-life exposure effects via repeatedly sharing night nests (Fossey, 1979) function analogously to the manner in which corresidence duration serves as a key component of kin recognition in humans (Lieberman et al., 2007), or that preferential mother-father relationships post-birth might lead to social preferences among siblings (Rosenbaum et al., 2016). Individuals may also possess some degree of phenotype matching ability (Widdig, 2007; Parr et al., 2010; Langergraber, 2012; Pfefferle et al., 2014).

Second, the lack of evidence for a prosocial bias towards paternal siblings is not readily reconciled with clear behavioral evidence of reduced aggression within these same dyads. This remarkable disjunct between apparent sibling recognition and sibling bias suggests that from a mountain gorilla's perspective, paternal siblings are known entities that nevertheless are less attractive social partners than maternal siblings, despite each being equal relatives. There may be multiple, non-mutually exclusive explanations for this dynamic. Perhaps the presence of paternal siblings provides fewer benefits to an individual than do other sibling types—this possibility, while previously suggested (e.g. Cords et al., 2018), has not been systematically investigated and is an ideal target for future research. A mismatch between historical and current social structure might also lead to inconsistent, weakened kin recognition among paternal siblings that manifests in the contrasting patterns we report. Ultimately, disentangling these potential explanations within a species that only

exists in the wild may depend on the opportunity to study long-term mating patterns and the impacts of "natural experiments" such as early maternal loss or adoption (most often carried out by adult males in this species; Fossey, 1979, Morrison et al. 2021).

Conclusion

Our analyses of sibling relationships in mountain gorillas provide extensive, large-scale information on the dynamics of cooperation and competition in a primate society where, as in humans, potential social partners vary greatly in the genes, developmental stage, and biological sex they share with each other. We find a selective sibling bias for prosocial behaviors, in that siblings who share matrilineal kinship affiliate at greater rates than either paternal half-siblings or non-siblings, and that this bias weakens as individuals become more distant in age. While such a result is consistent with a wide range of previous research, none of the reasons proposed for this selective bias in primates appear to apply to our population: mountain gorillas gain regular exposure to siblings of all types, across their entire lives; furthermore, patterns of aggressive behavior, in contrast to affiliation, suggest that mountain gorillas can in fact recognize paternal siblings, though they evidently do not favor them as cooperative partners. Ultimately, our study underscores a diversity of means, some evidently yet to be revealed, through which individuals might perceive and engage in sibling relationships to achieve fitness outcomes.

Materials and Methods

Our study subjects came from a population of habituated wild mountain gorillas living in Volcanoes National Park, Rwanda, that have been monitored nearly continuously for the last 54 years by the Dian Fossey Gorilla Fund. Using focal follow and scan data collected by researchers and staff, we compiled a dataset of all available dyadic gorilla behavior spanning the years of 2003 to 2017. We then supplemented this dataset with demographic and relatedness data (for maternal

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

relatedness, via direct observation; for paternal relatedness, via genetic paternity determination—see e.g. Vigilant et al., 2015) on individuals pulled from long-term records. From this combined dataset, we excluded interactions with infants <1 year of age at time of observation, parent-offspring interactions, and interactions between dyads for which we could not calculate relatedness from available data. This yielded a final, curated dataset containing 157 unique individuals (75F, 82M; average age at time of observation = 9.75 years) and 38,996 total hours of observation. Composition of dyads. Our dataset of behavior from 157 individuals contained 1957 unique dyad pairs. Of these dyads, 1258 shared neither a mother nor father ("non-siblings"), 555 shared a father but not a mother ("paternal siblings"), and 43 shared both a mother and a father ("full siblings"). In addition to dyads known to share a mother but not a father (n = 50), there were a number of dyads with the same mother, but with paternity data missing for one or both individuals (n = 51). To maximize sample size, we combined these two groups into the category of "maternal" siblings"; due to this analytic choice, this category can be effectively conceived of as "at least maternal siblings". See Tables S1 and S2 in Supplementary Materials for analyses using only confirmed maternal siblings, which were very similar to those reported below. Mixed-sex dyads were the most common sex category in our dataset (n = 981), followed by male-male (n = 520) and female-female (n = 456). Dyads differed in age by an average of 5.90 years (SD: 4.57 years; range: 0 - 23.5 years); for reference, the average interbirth interval in mountain gorillas is 3.9 years (Eckardt et al., 2016). We used this age difference variable as our primary index of familiarity between individuals. While we also had information on the natal groups of individuals, which could also serve as a potential index of familiarity, we do not focus on this variable in analyses, as it did not allow us to disambiguate between relatedness and familiarity-dyads of individuals who grew up in different natal groups were virtually never (n = 3) siblings in our dataset.

Behavioral Measures. We evaluated five different categories of dyadic behaviors as outcome variables: grooming, playing, non-contact aggression, contact aggression, and time spent in close (2m) proximity. We operationalized these behaviors from standardized definitions used in previous publications about this gorilla population (see e.g. Rosenbaum et al., 2015). Trained observers regularly undergo interobserver reliability tests. The former four behavioral categories were evaluated as counts (corrected for exposure time; see Data Analysis) within the dyad during focal observations, regardless of directionality, while the latter category of time in close proximity was evaluated by counting the number of instantaneous scan samples in which a dyad was observed within 2 m of each other (also corrected for exposure time).

Data Analysis. We conducted all analyses in R (version 4.1.2). Our main statistical models for each behavioral outcome consisted of cross-classified generalized linear mixed models (conducted using the glmmTMB package; Brooks et al., 2017) that included random intercepts for each individual within the dyad, as well as the dyad itself. Given low incidences of many behaviors, we aggregated behaviors into annual counts, making the dyad-year the fundamental unit of analysis (total n = 7858). Even with annual aggregation, instances of aggression were uncommon. Therefore, counts of contact and non-contact aggression were summed into a single category for analysis (see Tables S3 and S4 and Figures S1 and S2 in Supplementary Materials for results with individual aggression categories, which were qualitatively similar to those reported below).

In models predicting each behavioral outcome, we included terms for relatedness, age difference, and sex makeup, as well as two-way interactions between relatedness and sex makeup, relatedness and age difference, and sex makeup and age difference. As mothers plausibly mediate many of the social behaviors we examined, especially early in life, we also included the average proportion of observations with mothers in close proximity, and this variable's interaction with

relatedness, as covariates in all models. In models containing significant main effect or interaction terms, we decompose omnibus comparisons and report targeted marginal effects and contrasts using the *emmeans* package (Lenth, 2022), with all reported *p*-values corrected for false discovery rate. We modeled our count outcomes as rates with a negative binomial family in *glmmTMB* and offset term for exposure time (either logged hours of observation, or logged sum of scans for both individuals, per dyad-year). For each model, we verified model fit by inspecting the deviation, dispersion, and outliers of residuals using the *DHARMa* package (Hartig, 2022). All data and code necessary to reproduce our results are available publicly at https://osf.io/6qgj5.

Acknowledgments

The authors are grateful to Winnie Eckardt and Robin Morrison for their expertise and assistance with data. We thank the Rwandan government and the Rwanda Development Board for their long-term support of the research, monitoring, and protection activities of The Dian Fossey Gorilla Fund's Karisoke Research Center. We are deeply indebted to all Karisoke field staff for their tireless support in collecting long-term behavioral and demographic data. This study was supported by the University of Michigan, the NSF Graduate Research Fellowship Program and Doctoral Dissertation Improvement Grant No 1122321, and the donors who support The Dian Fossey Gorilla Fund.

Competing Interests

The authors report no competing interests.

398 399 References 400 401 Alberts, S. C., & Altmann, J. (2012). The Amboseli Baboon Research Project: 40 years of continuity 402 and change. In Long-term field studies of primates (pp. 261-287). Springer, Berlin, Heidelberg. 403 Alberts, S. C., Watts, H. E., & Altmann, J. (2003). Queuing and queue-jumping: long-term patterns 404 of reproductive skew in male savannah baboons, Papio cynocephalus. Animal Behaviour, 405 *65*(4), 821-840. 406 Bradley, B. J., Robbins, M. M., Williamson, E. A., Steklis, H. D., Steklis, N. G., Eckhardt, N., ... & 407 Vigilant, L. (2005). Mountain gorilla tug-of-war: silverbacks have limited control over 408 reproduction in multimale groups. Proceedings of the National Academy of Sciences, 102(26), 9418-409 9423. 410 Breuer, T., Robbins, A. M., & Robbins, M. M. (2016). Sexual coercion and courtship by male 411 western gorillas. Primates, 57(1), 29-38. 412 Brooks M.E., Kristensen K, van Benthem K.J., Magnusson A, Berg C.W., Nielsen A, Skaug H.J., 413 Maechler M, Bolker B.M. (2017). glmmTMB Balances Speed and Flexibility Among 414 Packages for Zero-inflated Generalized Linear Mixed Modeling. The R Journal, 9(2), 378–400. 415 Brown, G. R., & Dixson, A. F. (2000). The development of behavioural sex differences in infant 416 rhesus macaques (Macaca mulatta). Primates, 41(1), 63-77. 417 Chapais, B. (2001). Primate nepotism: what is the explanatory value of kin selection?. *International* 418 *Journal of Primatology*, 22(2), 203-229.

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

Charpentier, M. J. E., Tung, J., Altmann, J., & Alberts, S. C. (2008). Age at maturity in wild baboons: genetic, environmental and demographic influences. *Molecular Ecology*, 17(8), 2026-2040. Cords, M., Minich, T., Roberts, S. J., & Sleator, C. (2018). Evidence for paternal kin bias in the social affiliation of adult female blue monkeys. American Journal of Primatology, 80(5), e22761. Daly, M., & Wilson, M. (1988). Homicide. Eckardt, W., Fawcett, K., & Fletcher, A. W. (2016). Weaned age variation in the Virunga mountain gorillas (Gorilla beringei beringei): Influential factors. Behavioral Ecology and Sociobiology, 70, 493-507. Engh, A. L., Hoffmeier, R. R., Seyfarth, R. M., & Cheney, D. L. (2009). O brother, where art thou? The varying influence of older siblings in rank acquisition by female baboons. Behavioral Ecology and Sociobiology, 64(1), 97-104. Erhart, E. M., Coelho Jr, A. M., & Bramblett, C. A. (1997). Kin recognition by paternal halfsiblings in captive Papio cynocephalus. American Journal of Primatology, 43(2), 147-157. Fossey, D. (1979). Development of the mountain gorilla (Gorilla gorilla beringei), through the first thirty-six months. In D. A. Hamburg, E. R. McCown (Eds.), Perspectives on human evolution (Vol. 5, pp. 139–184) The Great Apes. Redwood City, California: The Benjamin-Cummings Publishing Company. Fredrickson, W. T., & Sackett, G. P. (1984). Kin preferences in primates (Macaca nemestrina): relatedness or familiarity?. Journal of Comparative Psychology, 98(1), 29. Galezo, A. A., Nolas, M. A., Fogel, A. S., Mututua, R. S., Warutere, J. K., Siodi, I. L. I., ... & Alberts, S. C. (2022). Mechanisms of inbreeding avoidance in a wild primate. Current Biology, 32(7), 1607-1615.

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

Godov, I., Vigilant, L., & Perry, S. E. (2016). Cues to kinship and close relatedness during infancy in white-faced capuchin monkeys, Cebus capucinus. Animal Behaviour, 116, 139-151. Goodall, J. (1986). The Chimpanzees of Gombe: Patterns of Behavior. Gould, L. (2000). Adoption of a wild orphaned ringtailed lemur infant by natal group members: adaptive explanations. Primates, 41(4), 413-419. Grafen, A. (1990). Do animals really recognize kin?. Animal Behaviour, 39(1), 42-54. Gray, P. (2019). Evolutionary functions of play: Practice, resilience, innovation, and cooperation. In P. K. Smith & J. L. Roopnarine (Eds.), The Cambridge handbook of play: Developmental and disciplinary perspectives (pp. 84–102). Cambridge University Press. Hamilton III, W. J., Busse, C., & Smith, K. S. (1982). Adoption of infant orphan chacma baboons. Animal Behaviour, 30(1), 29-34. Hamilton, W. D. (1964). The genetical evolution of social behaviour. II. Journal of Theoretical Biology, 7(1), 17-52. Hartig, F. (2022). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models. R package version 0.4.5. Hobaiter, C., Schel, A. M., Langergraber, K., & Zuberbühler, K. (2014). 'Adoption' by maternal siblings in wild chimpanzees. *PloS one*, 9(8), e103777. Hudson, R., & Trillmich, F. (2008). Sibling competition and cooperation in mammals: challenges, developments and prospects. Behavioral Ecology and Sociobiology, 62(3), 299-307. Johnson, C., Koerner, C., Estrin, M., & Duoos, D. (1980). Alloparental care and kinship in captive social groups of vervet monkeys (Cercopithecus aethiops sahaeus). Primates, 21(3), 406-415.

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

Lamb, M. E., Sutton-Smith, B., Sutton-Smith, B., & Lamb, M. E. (Eds.). (2014). Sibling relationships: Their nature and significance across the lifespan. New York, NY: Psychology Press. Langergraber K.A. (2012). Cooperation among kin. In: Mitani J.C., Call J., Kappeler P.M., Palombit R.A., Silk J.B (Eds). The evolution of primate societies. The University of Chicago Press, London, p. 491-513. Langergraber, K. E., Mitani, J. C., & Vigilant, L. (2007). The limited impact of kinship on cooperation in wild chimpanzees. Proceedings of the National Academy of Sciences, 104(19), 7786-7790. Lenth, R.V. (2022). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.7.2. https://CRAN.R-project.org/package=emmeans Lonsdorf, E. V., Stanton, M. A., & Murray, C. M. (2018). Sex differences in maternal sibling-infant interactions in wild chimpanzees. *Behavioral Ecology and Sociobiology*, 72(7), 1-12. Lea, A. J., Learn, N. H., Theus, M. J., Altmann, J., & Alberts, S. C. (2014). Complex sources of variance in female dominance rank in a nepotistic society. Animal Behaviour, 94, 87-99. Lu, P. C. (2007). Sibling relationships in adulthood and old age: A case study of Taiwan. Current *Sociology*, *55*(4), 621-637. Lynch, E. C., Di Fiore, A., Lynch, R. F., & Palombit, R. A. (2017). Fathers enhance social bonds among paternal half-siblings in immature olive baboons (Papio hamadryas anubis). Behavioral Ecology and Sociobiology, 71(8), 1-11. Maestripieri, D., & Ross, S. R. (2004). Sex differences in play among western lowland gorilla (Gorilla gorilla gorilla) infants: implications for adult behavior and social structure. American Journal of Physical Anthropology, 123(1), 52-61.

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

Mateo, J. M. (2015). Perspectives: Hamilton's legacy: Mechanisms of kin recognition in humans. Ethology, 121(5), 419-427. Meikle, D. B., & Vessey, S. H. (1981). Nepotism among rhesus monkey brothers. *Nature*, 294(5837), 160-161. Mitani, J. C. (2009). Male chimpanzees form enduring and equitable social bonds. *Animal Behaviour*, 77(3), 633-640. Mitani, J. C., Merriwether, D. A., & Zhang, C. (2000). Male affiliation, cooperation and kinship in wild chimpanzees. Animal Behaviour, 59(4), 885-893. Mock, D. & Parker, G. (1997). The evolution of sibling rivalry. Oxford University Press. Morrison, R. E., Mushimiyimana, Y., Stoinski, T. S., & Eckardt, W. (2021). Rapid transmission of respiratory infections within but not between mountain gorilla groups. Scientific Reports, 11, 19622. Nishida, T. (1983). Alloparental behavior in wild chimpanzees of the Mahale Mountains, Tanzania. Folia Primatologica, 41(1-2), 1-33. Nitsch, A., Faurie, C., & Lummaa, V. (2013). Are elder siblings helpers or competitors? Antagonistic fitness effects of sibling interactions in humans. Proceedings of the Royal Society B: Biological Sciences, 280(1750), 20122313. Parr, L. A., Heintz, M., Lonsdorf, E., & Wroblewski, E. (2010). Visual kin recognition in nonhuman primates (Pan troglodytes and Macaca mulatta): inbreeding avoidance or male distinctiveness?. *Journal of Comparative Psychology*, 124(4), 343.

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

Pfefferle, D., Kazem, A. J., Brockhausen, R. R., Ruiz-Lambides, A. V., & Widdig, A. (2014). Monkeys spontaneously discriminate their unfamiliar paternal kin under natural conditions using facial cues. Current Biology, 24(15), 1806-1810. Remis, M. J. (1997). Ranging and grouping patterns of a western lowland gorilla group at Bai Hokou, Central African Republic. American Journal of Primatology, 43(2), 111-133. Robbins, M.M. (2009). Male aggression toward females in mountain gorillas: courtship or coercion. In: Muller M.N., Wrangham R. (eds.) Sexual coercion in primates. Harvard University Press, Cambridge. Robbins, A. M., Stoinski, T. S., Fawcett, K. A., & Robbins, M. M. (2009). Socioecological influences on the dispersal of female mountain gorillas—evidence of a second folivore paradox. Behavioral Ecology and Sociobiology, 63(4), 477-489. Rosenbaum, S., Hirwa, J. P., Silk, J. B., Vigilant, L., & Stoinski, T. S. (2015). Male rank, not paternity, predicts male-immature relationships in mountain gorillas, Gorilla beringei beringei. Animal Behaviour, 104, 13-24. Rosenbaum, S., Hirwa, J. P., Silk, J. B., Vigilant, L., & Stoinski, T. S. (2016). Infant mortality risk and paternity certainty are associated with postnatal maternal behavior toward adult male mountain gorillas (Gorilla beringei beringei). Plos One, 11(2), e0147441. Rosenbaum, S., & Silk, J. B. (2022). Pathways to paternal care in primates. Evolutionary Anthropology. Roy, J., Gray, M., Stoinski, T., Robbins, M. M., & Vigilant, L. (2014). Fine-scale genetic structure analyses suggest further male than female dispersal in mountain gorillas. BMC Ecology, 14(1), 1-16.

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

Salmon, C. A., & Hehman, J. A. (2021). Good Friends, Better Enemies? The Effects of Sibling Sex, Co-Residence, and Relatedness on Sibling Conflict and Cooperation. Evolutionary Psychological Science, 7(4), 327-337. Sandel, A. A., Langergraber, K. E., & Mitani, J. C. (2020). Adolescent male chimpanzees (Pan troglodytes) form social bonds with their brothers and others during the transition to adulthood. American Journal of Primatology, 82(1), e23091. Schaub, H. (1996). Testing kin altruism in long-tailed macaques (Macaca fascicularis) in a foodsharing experiment. *International Journal of Primatology*, 17(3), 445-467. Sear, R., & Mace, R. (2008). Who keeps children alive? A review of the effects of kin on child survival. Evolution and Human Behavior, 29(1), 1-18. Silk, J. B. (2002). Kin selection in primate groups. *International Journal of Primatology*, 23(4), 849-875. Silk, J. B. (2009). Nepotistic cooperation in non-human primate groups. Philosophical Transactions of the Royal Society B: Biological Sciences, 364(1533), 3243-3254. Silk, J. B., Altmann, J., & Alberts, S. C. (2006). Social relationships among adult female baboons (Papio cynocephalus) I. Variation in the strength of social bonds. Behavioral Ecology and *Sociobiology*, *61*(2), 183-195. Smith, K., Alberts, S. C., & Altmann, J. (2003). Wild female baboons bias their social behaviour towards paternal half-sisters. Proceedings of the Royal Society of London. Series B: Biological Sciences, 270(1514), 503-510. Steelman, L. C., Powell, B., Werum, R., & Carter, S. (2002). Reconsidering the effects of sibling configuration: Recent advances and challenges. Annual Review of Sociology, 28(1), 243-269.

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

Stoinski, T. S., Vecellio, V., Ngaboyamahina, T., Ndagijimana, F., Rosenbaum, S., & Fawcett, K. A. (2009). Proximate factors influencing dispersal decisions in male mountain gorillas, Gorilla beringei beringei. Animal Behaviour, 77(5), 1155-1164. Surbeck, M., Langergraber, K. E., Fruth, B., Vigilant, L., & Hohmann, G. (2017). Male reproductive skew is higher in bonobos than chimpanzees. Current Biology, 27(13), R640-R641. Trivers, R. L. (1974). Parent-offspring conflict. *Integrative and Comparative Biology*, 14(1), 249-264. Tung, I., Archie, E. A., Altmann, I., & Alberts, S. C. (2016). Cumulative early life adversity predicts longevity in wild baboons. Nature Communications, 7(1), 1-7. Vigilant, L., Roy, J., Bradley, B. J., Stoneking, C. J., Robbins, M. M., & Stoinski, T. S. (2015). Reproductive competition and inbreeding avoidance in a primate species with habitual female dispersal. Behavioral Ecology and Sociobiology, 69(7), 1163-1172. Widdig, A. (2007). Paternal kin discrimination: the evidence and likely mechanisms. *Biological Reviews*, *82*(2), 319-334. Widdig, A., Bercovitch, F. B., Jürgen Streich, W., Sauermann, U., Nürnberg, P., & Krawczak, M. (2004). A longitudinal analysis of reproductive skew in male rhesus macaques. *Proceedings of* the Royal Society of London. Series B: Biological Sciences, 271(1541), 819-826. Widdig, A., Nürnberg, P., Krawczak, M., Streich, W. J., & Bercovitch, F. B. (2001). Paternal relatedness and age proximity regulate social relationships among adult female rhesus macaques. Proceedings of the National Academy of Sciences, 98(24), 13769-13773. Widdig, A., Nürnberg, P., Krawczak, M., Streich, W. J., & Bercovitch, F. (2002). Affiliation and aggression among adult female rhesus macaques: a genetic analysis of paternal cohorts. Behaviour, 371-391.

Wikberg, E. C., Ting, N., & Sicotte, P. (2014). Familiarity is more important than phenotypic similarity in shaping social relationships in a facultative female dispersed primate, Colobus vellerosus. *Behavioural Processes*, 106, 27-35.
Wu, H. M., Holmes, W. G., Medina, S. R., & Sackett, G. P. (1980). Kin preference in infant *Macaca nemestrina*. *Nature*, 285(5762), 225-227.

Supplementary Analyses for "Neither kin selection nor familiarity explain affiliative biases towards maternal siblings in wild mountain gorillas"

Nicholas M. Grebe, Jean Paul Hirwa, Tara S. Stoinski, Linda Vigilant, & Stacy Rosenbaum

Tables S1 – S2. Results for analyses using a 'stricter' categorization of maternal siblings (n = 50; see Methods for details of categorization; total dyad-years: 7625).

Table S1. Omnibus statistics for target parameters (full model results available from data and code posted publicly at https://osf.io/6qgj5). Effects p < 0.05 bolded.

	Effect							
	Relatedness	Sex Category	Age Differences	Rel.× Sex Category	Rel.× Age Diff.	Sex × Age Diff.		
Play		$\chi^2(2) = 13.64, p = 0.001$						
Grooming	$\chi^2(3) = 16.05, p < 0.001$	$\chi^2(2) = 10.88, p = 0.004$	$\chi^2(1) = 2.08,$ $p = 0.149$	$\chi^{2}(6) = 6.88$ $p = 0.331$	$\chi^2(3) = 8.44,$ $p = 0.038$	$\chi^2(2) = 13.94, p < 0.001$		
Proximity	$\chi^2(3) = 17.64, p < 0.001$	$\chi^2(2) = 1.69,$ $p = 0.423$	$\chi^2(1) = 1.58,$ $p = 0.209$	$\chi^2(6) = 7.74,$ $p = 0.258$	$\chi^2(3) = 63.81, \\ p < 0.001$	$\chi^2(2) = 0.30,$ $p = 0.985$		
Aggression	$\chi^2(3) = 1.46,$ $p = 0.691$	$\chi^2(2) = 0.54,$ $p = 0.762$	$\chi^2(1) = 4.30, p = 0.022$	$\chi^{2}(6) = 11.52, p = 0.073$	$\chi^2(3) = 2.68,$ $p = 0.443$	$\chi^2(2) = 6.67,$ $p = 0.036$		

Table S2. Estimated marginal means and standard errors across relatedness and sex categories.

		Relate	edness	Sex Category			
	Full Siblings	Maternal Half	Paternal Half	Non- Siblings	Female - Female	Male - Male	Mixed - Sex
Play	2.91 (0.58)	1.83 (0.42)	1.42 (0.17)	1.14 (0.11)	1.17 (0.22)	2.95 (0.50)	1.46 (0.20)
Grooming	2.51 (0.60)	1.61 (0.45)	0.30 (0.04)	0.34 (0.04)	1.49 (0.30)	0.48 (0.10)	0.73 (0.11)
Proximity	44.9 (5.5)	42.6 (5.9)	15.8 (1.0)	18.9 (1.0)	30.2 (3.2)	25.9 (2.6)	26.4 (2.1)
Aggression	0.43 (0.08)	0.64 (0.13)	0.56 (0.04)	0.76 (0.05)	0.50 (0.07)	0.68 (0.09)	0.60 (0.06)

Tables S3 – S4, Figures S1-S2. Results for analyses assessing contact aggression and non-contact aggression separately (total dyad-years: 7822).

Table S3. Omnibus statistics for target parameters (full model results available from data and code posted publicly at https://osf.io/6qgj5).

	Effect								
	Relatedness	Sex Category	Age Differences	Rel.× Sex Category	Rel.× Age Diff.	Sex × Age Diff.			
	** ' '	$\chi^2(2) = 1.57,$ $p = 0.456$		$\chi^2(6) = 10.54, p = 0.104$		$\chi^2(2) = 3.74,$ $p = 0.154$			
Non- Contact Aggression	6 - 0 309	$\chi^2(2) = 3.35,$ $p = 0.187$,, ,	$\chi^{2}(6) = 10.23,$ $p = 0.116$,, ,	$\chi^2(2) = 2.22,$ $p = 0.330$			

Table S4. Estimated marginal means and standard errors across relatedness and sex categories.

		Relate	dness	Sex Category			
	Full Siblings	Maternal Half	Paternal Half	Non- Siblings	Female - Female	Male - Male	Mixed - Sex
Contact Aggression	0.23 (0.04)	0.22 (0.04)	0.28 (0.02)	0.34 (0.02)	0.22 (0.03)	0.28 (0.04)	0.30 (0.03)
Non- Contact Aggression	0.10 (0.03)	0.18 (0.04)	0.18 (0.02)	0.23 (0.02)	0.18 (0.03)	0.18 (0.03)	0.13 (0.02)

Figure S1. Box and dot plots showing estimated non-contact aggression within gorilla dyads, separated by relatedness and sex category.




Figure S2. Box and dot plots showing estimated contact aggression within gorilla dyads, separated by relatedness and sex category.

