

# Appendix S5 - Extracting PRISM Climate Data

*Nicholas J Clark, David J Harris, Ceridwen Fraser*

This appendix describes how to calculate mean PRISM climate variables for 40km-radius buffers around BBS sample sites. These matrices can be accessed in the `BBS.occurrences` library using `data('Bird.phy.distances')` and `data('Site.descriptors')`.

First, we need to download the raw climate data for each point in the dataset. Specify a file path for downloading PRISM climate data, and set this as the option for the `prism` library to store and access files

```
dir.create("/PRISM/ppt")
filepath <- "/PRISM/ppt"
options(prism.path = filepath)
```

Download monthly min temp data for the entire USA, starting the year prior to the first year of surveys

```
prism::get_prism_monthlys(type = "ppt", year = 2002:2009,
  mon = 1:12, keepZip = F)
```

Load the species occurrence dataset, which lists the binary occurrences of 303 avian species (stored in columns) across sites

```
library(BBS.occurrences)
data("Site.descriptors")
```

We need to extract unique sample sites and convert these points to a spatial dataframe so that they can be intersected with the `prism` rasters

```
library(dplyr)
points <- Site.descriptors %>% dplyr::ungroup() %>%
  dplyr::select(Latitude, Longitude) %>% dplyr::distinct()

points.spdf <- sp::SpatialPointsDataFrame(coords = points[,
  c("Longitude", "Latitude")], data = points, proj4string = CRS("+proj=longlat +ellps=WGS84 +no_defs"))
```

Extract mean estimates for PRISM climate variables (for precipitation, in this case) from 40km buffers around each point. If space for storing the raw files is an issue, use argument `delete_files = TRUE` to ensure files are deleted once calculations are complete

```
buffer <- 40000
points.ppt <- meanPRISM(points = points, points_spdf = points.spdf,
  var_name = "Precipitation", buffer = buffer, filepath = filepath,
  n_cores = 3, delete_files = FALSE)
save(points.ppt, file = "./Analysis_data/points.ppt.rda")
```

Repeat the above procedure for each climate variable (`temp.min`, `temp.mean`, and `temp.max`). Load all the saved climate datasets and join together in a single `dataframe`

```
load("./Analysis_data/points.ppt.rda")
load("./Analysis_data/points.tmean.rda")
load("./Analysis_data/points.tmin.rda")
load("./Analysis_data/points.tmax.rda")

monthly_climate_data = points.ppt %>% left_join(points.tmax) %>%
  left_join(points.tmean) %>% left_join(points.tmin)
```

Process to return climate variables describing conditions during the previous Spring season and during the previous year to be used as predictors in analyses. Ignore warnings here, these are just to showcase that a small number of specific points have no available measurements for certain months

```
climate.predictors <- processPRISM(monthly_climate_data)
```

Finally, join these predictors to the `Site.descriptors` file and save. Note, the climate predictors and other site description variables can be directly accessed in the `BBS.occurrences` library using `data('Site.descriptors')`.

```
Site.descriptors <- Site.descriptors %>% dplyr::ungroup() %>%  
  left_join(climate.predictors)
```