

# FYS3150 Computational Physics - Project 4

Nicholas Karlsen

Using the Metropolis algorithm to study a system of weakly coupled paramagnets, the Ising model. Derived exact solutions of expectation values for a 2x2 lattice and found that the required  $\sim 10^5$  Monte-Carlo cycles to consistent results with  $\sigma \sim 10^{-3}$ .

## INTRODUCTION

## THEORY, ALGORITHMS AND METHODS

### The Ising Model

In its simplest case, a paramagnet is described as a lattice of noninteracting dipole moments, each in a state up or down. Where the energy of the system is proportional to the sum of the spins, up or down of each noninteracting constituent.

The Ising model is a natural extension to this, by adding local interaction between directly neighboring spins (See fig. 1), such that the energy of the system is instead proportional to the sum of the interactions of all coupled spins, which gives a more accurate description of the system compared to the simple case of an ideal paramagnet.

Or more precisely, the energy is given by Eqn. 1, where  $s_i \in \{-1, 1\}$  and  $J$  is a coupling constant.

$$E = -J \sum_{\langle kl \rangle} s_k s_l \quad (1)$$

In this sum, the notation  $\langle kl \rangle$  means that we take the sum over each coupling  $s_k s_l$ . Consider again Fig. 1. The contribution,  $\epsilon_i$ , to the total energy due the interactions at spin-site  $i$  is the sum of its interactions,  $\epsilon_i = s_i(s_{\uparrow} + s_{\downarrow} + s_{\leftarrow} + s_{\rightarrow})$  where the arrows denote the spin-site directly neighboring site  $i$  in the direction of the arrow. Now wish to compute the energy contribution due to the interactions at the spin-site left of site  $i$ . In this case, the interaction between it, and spin-site  $i$  has already been computed, and must not be counted again.




FIG. 1: Section of a 2D lattice, where each spin-site is represented by a dot. In the Ising model, spin-site  $i$  will only interact with its directly neighbouring spin-sites, connected by dotted lines in this diagram

Further, in this project the lattice will be treated as pseudo-continuous where any spin-sites located at an edge, will also interact with the spin-site on the opposing edge, illustrated in Fig. 2, which strictly speaking constitutes having periodic boundary conditions. If instead we had fixed boundary conditions, there would simply be no such across-lattice interactions, and spin-sites located at the edges would then only interact with 3 other spin-sites (2 in the case of corner sites) in the 2D case.

Much simpler, the net magnetization,  $\mathcal{M}$ , of the system is taken as the sum over all spins

$$\mathcal{M} = \sum_i s_i$$




FIG. 2: Spin-site  $i$ , located at the left edge of the lattice interacting with its direct neighbors, as well as the spin site on the opposite edge of the lattice

### Boltzmann Statistics

From this model we can derive many thermodynamic quantities by the use of Boltzmann statistics, where we

have the probability of the system being in some specific macro-state<sup>1</sup> at temperature<sup>2</sup>

$$\mathcal{P}(E_s) = \frac{1}{Z} e^{-\beta E_s} \quad (2)$$

With  $\beta \equiv \frac{1}{k_B T}$ , the thermodynamic beta and  $Z$ , the partition function, acting as a normalization factor given by

$$Z = \sum_s e^{-\beta E_s} \quad (3)$$

From this, the expectation value of some quantity  $X$ , with micro-states  $X_s$  can be computed by

$$\langle X \rangle = \sum_s X_s \mathcal{P}(E_s) = \frac{1}{Z} \sum_s X_s e^{-\beta E_s} \quad (4)$$

Using this, we can compute the specific heat capacity,  $C_V$  by computing the expectation values of  $E, E^2$  [1], related by the following expression

$$C_V = \frac{\langle E^2 \rangle - \langle E \rangle^2}{k_B T^2} \quad (5)$$

Where  $\langle X^2 \rangle - \langle X \rangle^2$  is recognized as the variance of a set of randomly distributed quantities,  $\sigma_X^2$ .

In a similar fashion, the magnetic susceptibility<sup>3</sup>,  $\chi$ , is calculated from  $\sigma_{|\mathcal{M}|}^2$  [1]

$$\chi = \frac{\sigma_{|\mathcal{M}|}^2}{k_B T} \quad (6)$$

For more details on the Ising model, or Boltzmann statistics refer to Schroeder [2, Ch 6, 8.2], Hjorth-Jensen [1], or similar introductory texts on the topic.

#### Analytic solution of 2x2 Lattice

Consider now a 2x2 lattice, each with spin  $\pm 1$ . The energy of the system for a particular micro-state is given by Eqn. 1, so for a  $2 \times 2$  lattice we have

$$E = -J \sum_{\langle kl \rangle} s_{kl} = -J(2s_1s_2 + 2s_1s_3 + 2s_2s_4 + 2s_3s_4) \quad (7)$$

The multiplicity, that is number of possible combinations of micro-states for a system consisting of 4 spins is  $2^4 = 16$ .




FIG. 3: A  $2 \times 2$  lattice of spin-sites

If we first consider the micro-state in which all the spins are in parallel, we get energy  $E = -8J$ . Thus, this particular macro-state is at least 2-fold degenerate<sup>4</sup>. The different micro-states and their degeneracies are summed up in table I from Hjorth-Jensen [1]

TABLE I: Possible configurations of 2x2 lattice

No. spin up	Degeneracy	$E$	$\mathcal{M}$
4	1	-8J	4
3	4	0	2
2	4	0	0
2	2	8J	0
1	4	0	-2
0	1	-8J	-4

See that there are only 3 distinct energy macro-states,  $E = -8J, 0, 8J$  corresponding to 2, 12 and 2 of the possible micro-states respectively.

It is then easy to see that the partition function for the system will take the following form

$$Z = 2e^{+8J\beta} + 2e^{-8J\beta} + 12 = 4 \cosh(8J\beta) + 12 \quad (8)$$

From this, we derive analytic values for  $\langle E \rangle, \langle |\mathcal{M}| \rangle, C_V$  and  $\chi$ .

First, compute the expectation value of the energy using an alternate, simpler expression from Schroeder [2, Ch 6.2]

$$\langle E \rangle = -\frac{1}{Z} \frac{\partial}{\partial \beta} Z = -\frac{8J \sinh(8J\beta)}{\cosh(8J\beta) + 3} \quad (9)$$

Similarly, the expectation value of  $E^2$  is computed easily by another expression found in Schroeder [2, Ch 6.2]

$$\langle E^2 \rangle = -\frac{1}{Z} \frac{\partial^2}{\partial \beta^2} Z = -\frac{64J^2 \cosh(8J\beta)}{\cosh(8J\beta) + 3} \quad (10)$$

From this, we compute the variance of  $E$

$$\begin{aligned} \sigma_E^2 &= \langle E^2 \rangle - \langle E \rangle^2 \\ &= \frac{-64J^2}{\cosh(8J\beta) + 3} \left( \cosh(8J\beta) + \frac{\sinh^2(8J\beta)}{\cosh(8J\beta) + 3} \right) \end{aligned} \quad (11)$$

<sup>1</sup> Macroscopic states being characterized by their energy, meaning i could just as well have written "... the probability of the system having some specific energy"

<sup>2</sup> Contained within  $\beta$  in this notation

<sup>3</sup> Using the absolute magnetization to compute  $\chi$ , in line with the problem text for 4e

<sup>4</sup> Two micro-states corresponds to the same observable macro-state.

With which we can compute  $C_V$  using Eqn. 5. Compute expectation values of  $|\mathcal{M}|$ ,  $|\mathcal{M}|^2$  using Eqn. 4 and Table I

$$\begin{aligned}\langle |\mathcal{M}| \rangle &= \frac{1}{Z} \sum_s |\mathcal{M}_s| e^{-\beta E_s} \\ &= \frac{8e^{8J\beta} + 16e^0}{Z} = \frac{2e^{8J\beta} + 4}{\cosh(8J\beta) + 3}\end{aligned}\quad (12)$$

$$\begin{aligned}\langle |\mathcal{M}|^2 \rangle &= \frac{1}{Z} \sum_s |\mathcal{M}_s|^2 e^{-\beta E_s} \\ &= \frac{1}{Z} (16e^{8J} + 32e^0) = \frac{4e^{8J\beta} + 8}{\cosh(8J\beta) + 3}\end{aligned}\quad (13)$$

We then compute the variance of the net magnetization

$$\begin{aligned}\sigma_{\mathcal{M}}^2 &= \langle |\mathcal{M}|^2 \rangle - \langle |\mathcal{M}| \rangle^2 \\ &= \frac{1}{\cosh(8J\beta) + 3} \left( 4e^{8J\beta} + 8 - \frac{4e^{16J\beta} + 16}{\cosh(8J\beta) + 3} \right)\end{aligned}\quad (14)$$

Which can be used to compute the magnetic susceptibility using Eqn 6.

## The Metropolis Algorithm

### Metropolis Algorithm

1 Initialize State

## RESULTS AND DISCUSSIONS

See in Fig 4

## CONCLUSIONS

- 
- [1] M. Hjorth-Jensen, “Computational physics lectures: Statistical physics and the ising model,” (2017).
  - [2] D. V. Schroeder, *An Introduction to Thermal Physics*, 1st ed. (Pearson, 1999).




FIG. 4: (a-d) Computed results from running the algorithm 3000 times per number of monte-carlo cycles for select expectation values with an ordered initial state of a 2x2 lattice of spin ups. (e-h) The standard deviation (Std.) of the computed results.