

# Calculus in Brief with Python, Colab, GitHub, and L<sup>A</sup>T<sub>E</sub>X CiB - Version 0.42

November 12, 2023

MIT License

Available on GitHub at:

<https://GitHub.com/nicholaskarlson/CiB>



# Contents

<b>Preface</b>	<b>7</b>
<b>1 Introduction to CiB</b>	<b>9</b>
<b>2 Open-Source Ethos</b>	<b>11</b>
<b>3 Introduction to GitHub</b>	<b>13</b>
<b>4 Encouragement to Fork</b>	<b>15</b>
<b>5 More About GitHub</b>	<b>17</b>
<b>6 Forking Process</b>	<b>21</b>
<b>7 Editing and Customizing</b>	<b>25</b>
<b>8 Engaging with the Community</b>	<b>29</b>
<b>9 Functions</b>	<b>33</b>
9.1 What is a Function? . . . . .	33
9.1.1 Definition and Examples . . . . .	33
9.1.2 Exercises . . . . .	34
9.1.3 Solutions to Exercises . . . . .	35
9.2 Domain and Range . . . . .	36
9.2.1 Exercises on Domain and Range . . . . .	38
9.2.2 Solutions to Exercises on Domain and Range . . . . .	39
9.3 Types of Functions . . . . .	39
9.3.1 Linear Functions . . . . .	39
9.3.2 Polynomial Functions . . . . .	42
9.3.3 Rational Functions . . . . .	45
9.3.4 Trigonometric Functions . . . . .	48
9.3.5 Exponential and Logarithmic Functions . . . . .	50

9.4	Function Properties . . . . .	53
9.4.1	Continuity . . . . .	53
9.4.2	Differentiability . . . . .	55
9.4.3	Asymptotic Behavior . . . . .	58
9.4.4	Periodicity and Symmetry . . . . .	61
9.5	Function Transformations . . . . .	64
9.5.1	Translation and Scaling . . . . .	64
9.5.2	Reflection and Rotation . . . . .	67
<b>10</b>	<b>Limits and Continuity - Detail</b>	<b>71</b>
10.1	Introduction to Limits - Detail . . . . .	71
10.1.1	Definition of a Limit - Detail . . . . .	71
10.1.2	One-sided Limits . . . . .	73
10.1.3	Limits Involving Infinity . . . . .	73
10.2	Properties of Limits . . . . .	73
10.2.1	Limit Laws . . . . .	73
10.2.2	Squeeze Theorem . . . . .	74
10.2.3	Limits of Trigonometric Functions . . . . .	74
10.3	Continuity - Detail . . . . .	74
10.3.1	Definition and Examples . . . . .	74
10.3.2	Types of Discontinuities . . . . .	74
10.3.3	Intermediate Value Theorem . . . . .	74
<b>11</b>	<b>The Derivative - Detail</b>	<b>75</b>
11.1	Understanding Derivatives . . . . .	75
11.1.1	Definition and Notation . . . . .	75
11.1.2	Interpreting Derivatives . . . . .	75
11.1.3	Derivative as a Function . . . . .	75
11.2	Techniques of Differentiation . . . . .	75
11.2.1	Product Rule and Quotient Rule . . . . .	75
11.2.2	Chain Rule . . . . .	76
11.2.3	Higher-Order Derivatives . . . . .	76
11.2.4	Implicit Differentiation . . . . .	76
11.3	Applications of Derivatives . . . . .	76
11.3.1	Tangent Lines and Normal Lines . . . . .	76
11.3.2	Increasing and Decreasing Functions . . . . .	76
11.3.3	Maxima and Minima . . . . .	76
11.3.4	Concavity and Inflection Points . . . . .	76
11.3.5	Optimization Problems . . . . .	76
11.3.6	Related Rates . . . . .	77

<b>12 Integration</b>	<b>79</b>
12.1 Antiderivatives . . . . .	79
12.1.1 Definition and Basic Techniques . . . . .	79
12.1.2 Indefinite Integrals . . . . .	79
12.1.3 Initial Value Problems . . . . .	79
12.2 Definite Integrals . . . . .	79
12.2.1 Definition and Interpretation . . . . .	79
12.2.2 Properties of Definite Integrals . . . . .	80
12.2.3 Fundamental Theorem of Calculus . . . . .	80
12.2.4 Numerical Integration Methods . . . . .	80
12.3 Applications of Integrals . . . . .	80
12.3.1 Area Between Curves . . . . .	80
12.3.2 Volumes of Solids of Revolution . . . . .	80
12.3.3 Arc Length and Surface Area . . . . .	80
12.3.4 Work and Fluid Forces . . . . .	80
<b>13 Advanced Topics in Calculus</b>	<b>81</b>
13.1 Sequences and Series . . . . .	81
13.1.1 Convergence and Divergence . . . . .	81
13.1.2 Tests for Convergence . . . . .	81
13.1.3 Power Series and Taylor Series . . . . .	81
13.2 Multivariable Calculus . . . . .	81
13.2.1 Functions of Several Variables . . . . .	81
13.2.2 Partial Derivatives . . . . .	82
13.2.3 Multiple Integrals . . . . .	82
13.2.4 Vector Calculus . . . . .	82
13.3 Differential Equations . . . . .	82
13.3.1 First Order Differential Equations . . . . .	82
13.3.2 Second Order Linear Differential Equations . . . . .	82
13.3.3 Systems of Differential Equations . . . . .	82
13.4 Conclusion . . . . .	82
<b>Appendices</b>	<b>83</b>
<b>I Basic GitHub Guide</b>	<b>83</b>
<b>II Basic Python and Colab Guide</b>	<b>87</b>
<b>III Basic L<sup>A</sup>T<sub>E</sub>X Guide</b>	<b>93</b>
<b>Bibliography</b>	<b>95</b>



# Preface

This text, *Calculus in Brief - CiB* aspires to be more than just another math book. This book strives to foster collaborative math writing. Note that this book has very few references. The reader is encouraged to use resources available on the Web to fact-check. This book’s view on “causation” and facts is heavily influenced by Mosteller and Tukey [MT77].

## Redefining the Role of the Reader

Calculus in Brief (CiB) is an endeavor to reshape how math is written, understood, and studied. It’s not just a passive read but an open-source approach to math, aiming to encourage students to become proactive learners.

This project strives to break the traditional mold of math education and invites readers and professional mathematicians to participate actively.

## A Dynamic Relationship with Math

*Calculus in Brief* is not just a book but a movement and methodology, heralding a new era in how we approach, consume, and interact with math. By positioning the reader as an integral part of the math-book process, CiB fosters a dynamic relationship with math, making mathematics more accessible, proactive, and relevant. In this shifting paradigm, we are all potential mathematicians, creators of interesting and relevant ways to learn and study math.

Please fork the LaTeX source code for CiB (available on GitHub) and create your own book that chooses the facts and exercises most relevant to you! Also, starring the CiB project on GitHub would be greatly appreciated! Thanks for reading CiB!





# Chapter 1

## Introduction to CiB

### Welcome to CiB on GitHub

Calculus in Brief, abbreviated CiB, isn't merely a passive read. It's an endeavor to reshape how math is written, studied, and taught. By presenting an open-source approach to math, the goal is to encourage everyone to become proactive readers and writers of math.

### Fostering a Proactive Engagement with Math

*Calculus in Brief* is a call for a renewed engagement with mathematics. CiB is an endeavor to reshape how math is written, understood, and studied. It's not just a passive read but an open-source approach to math, aiming to encourage students to become proactive learners.

This project strives to break the traditional mold of math education and encourages readers and professional mathematicians to participate actively.

Please fork the L<sup>A</sup>T<sub>E</sub>X source code for CiB (available on GitHub) and create your own book on Calculus that chooses the content most relevant to you! Also, starring the CiB project on GitHub would be greatly appreciated! Thanks for reading CiB!



# Chapter 2

## Open-Source Ethos

### **The Spirit of Shared Knowledge and Collaboration**

Math, like software, is better when it's open. CiB draws inspiration from the open-source software movement; this section elucidates how a collaborative, transparent, and shared approach can enhance our understanding of math. Here, we look at the philosophy behind open-source and how it beautifully combines with the study of mathematics.

### **Open-Source Math: Preserving Tradition Through Collaborative Exploration**

Mathematics, like software, thrives when it embraces openness and transparency. CiB takes a leaf from the proven benefits of the open-source software model; this section highlights how a collaborative and transparent method can improve and deepen our grasp of math and its texts. Here, we explore the principles of open source and how these principles align with the development of mathematics and its texts.

### **Understanding the Open-Source Ethos**

The open-source paradigm revolves around shared ownership, collaboration, and the free exchange of knowledge. In the software realm, this approach has led to groundbreaking innovations built and enhanced by a global community of skilled contributors. United by a mutual objective, these individuals pool their diverse talents and insights to improve and share software solutions for broader public benefit.

## Advantages of the Open-Source Framework in Math

### Collective Insight

Mirroring the collaborative essence of open-source software, many individuals can offer their perspectives and knowledge, making math texts more robust and varied.

### Enhancement and Accuracy

Open platforms foster an environment of constructive criticism, ensuring prompt identification and correction of inaccuracies. This meticulous peer review can help provide a credible and current mathematical text.

### Universal Access

Much as open-source software promotes free access and modification, open-source math prioritizes universal accessibility. This ensures mathematics knowledge isn't restricted to a select few but is available to all curious minds.

## Potential Challenges

Despite its advantages, melding open-source with math has potential pitfalls. The volume of contributions can complicate accuracy verification processes.

However, the very community championing this open-source approach to math can serve as its vigilant protector. They can ensure that contributions undergo rigorous evaluation and referencing, akin to the meticulous checks within the open-source software community.

## Conclusion: Reinvigorating Our Experience with Math

Adopting an open-source perspective to the approach of math signifies a refreshed approach. It beckons a worldwide community to collaborate and forge a comprehensive and exciting math text. In this refreshed approach, every individual can play a part, both as a contributor and a learner. Math texts, through this lens, evolve and flourish, reflecting the collective input of active participants.

# Chapter 3

## Introduction to GitHub

### The Hub for Modern Collaboration

#### Harnessing GitHub: A New Frontier in Collaborative Math Writing

At the heart of our collaborative math endeavor lies GitHub, a platform traditionally associated with code but now repurposed for our endeavor. This section provides a primer on GitHub, laying the foundation for those unfamiliar and offering insights into its transformative potential for collective math writing, learning, and teaching.

#### A Brief Introduction to GitHub

Originally conceptualized as a platform for developers, GitHub is a repository hosting service that facilitates version control using Git. At its core, it allows multiple users to work on a project simultaneously, tracking changes and ensuring that the latest version of a project is always accessible. Over the years, GitHub has grown beyond its initial software-centric confines, becoming a hub for all kinds of collaborative projects, from writing to data science and now to math.

#### Repurposing GitHub for Math Texts

##### Version Control

Math writing, like software, is dynamic and constantly evolving. As new sources or perspectives emerge, math texts may need revisions. GitHub's

version control ensures that every change made to a document is tracked, enabling mathematicians to see how math texts evolve over time.

### **Collaborative Writing**

Multiple contributors can work on a single math text simultaneously. This multi-user capability ensures diverse viewpoints can be seamlessly integrated, making the math text richer and more comprehensive.

### **Review and Feedback**

Just as developers review and comment on code, mathematicians can provide feedback on written content. This feature encourages rigorous peer review, ensuring accuracy and credibility.

### **Open Access**

Math texts on GitHub can be made public, granting anyone access to read, contribute, or fork the text into their own versions. This workflow democratizes math texts, making the creation process a collective endeavor rather than the domain of a select few.

### **Transparency**

All changes and contributions are logged, providing a clear trail of the evolution of a mathematical text. This transparency bolsters the credibility of the text hosted on the platform.

### **Community Building**

Beyond just writing, GitHub fosters a community of mathematicians, enthusiasts, and readers who can discuss, debate, and engage in meaningful dialogues about math and available math texts on GitHub.

## **Conclusion: Envisioning a Collaborative Mathematical Landscape**

Embracing GitHub as a tool for collaborative math signifies more than just a shift in approach; it heralds a new era of inclusivity, transparency, and dynamism in writing, learning, and math teaching.

# Chapter 4

## Encouragement to Fork

### Invitation to Dive Deep and Make It Your Own

CiB isn't a static entity. It thrives on evolution, adaptation, and diversification, much like math itself. We encourage readers to "fork" (a term soon to be discussed) and create their own versions of this book. Read this section to understand the essence of "forking" and how it can be the starting point of your unique math journey.

### The Concept of Forking: A Brief Overview

In the realm of software development, particularly in platforms like GitHub, "forking" refers to the act of creating a copy of a project, allowing one to make changes independently of the original. In this context, forking CiB enables readers to take the base content and adapt, modify, and expand upon it, tailoring the narrative to resonate with their perspectives, insights, and understanding.

### How to Begin Your Forking Journey

**Start Small:** You don't need to rewrite entire chapters. Begin by adding annotations, insights, or even footnotes to existing content. As you grow more confident, you can expand and modify larger sections.

**Engage with the Community:** Share your forked version with other readers. This encourages discourse, debate, and constructive feedback, allowing your text to be refined and enhanced.

**Celebrate input:** Encourage others around you to fork and create their own versions. The more in-depth the input, the deeper our collective understanding of math potentially becomes.

**Conclusion: The Power of Collective Math**

The invitation to fork CiB isn't just about creating different versions of a book. It's a call to embrace collective writing, learning, and teaching. By embracing the essence of forking, math is not just something we read but something we actively shape, share, and pass on.



# Chapter 5

## More About GitHub

### Discovering the Power of Collaborative Tools

Diving deeper into the world of GitHub, this chapter provides a comprehensive overview. Beyond its technicalities, we explore how GitHub emerged as a revolutionary platform for collaboration and how it can be leveraged for those interested in writing, teaching, and learning about math.

### The Genesis of GitHub

GitHub began as a platform designed for software developers to manage and track changes to their codebase. Launched in 2008, it swiftly gained traction due to its user-friendly interface and efficient version control system powered by Git. Over the years, it evolved from a mere repository hosting service to a dynamic hub of collaboration, housing millions of projects and engaging tens of millions of users worldwide.

### GitHub: More than Just Code

While GitHub's origins are rooted in code collaboration, its adaptable nature has made it a favored platform for various non-code projects. Writers, designers, educators, and researchers have discovered the potential of GitHub as a tool for:

#### Document Collaboration

With its built-in version control, contributors can track changes, revert to previous versions, and seamlessly merge updates.

### **Project Management**

With features like "issues" and "milestones," teams can organize tasks, set goals, and monitor progress.

### **Open Access & Transparency**

Public repositories allow for open contributions, ensuring transparency and fostering a sense of collective ownership.

### **Collaborative Writing**

Multiple contributors can simultaneously work on a single document, with every change being tracked and attributed, facilitating teamwork on extensive projects like books or research papers.

### **Engaging the Public**

With the platform's inherent transparency, researchers can make their work-in-progress accessible to the public, inviting insights, corrections, and contributions.

### **Case Study: CiB's Use of GitHub**

CiB's journey on GitHub is a testament to the platform's potential in mathematical endeavors. By hosting the book on GitHub, the following is possible:

#### **Feedback Loop**

Readers can raise "issues," pointing out inaccuracies, suggesting enhancements, or even recommending new sections or topics.

#### **Forking**

As previously discussed, readers can "fork" the repository, creating their unique versions of the book while staying connected to the original.

#### **Regular Updates**

With math being dynamic, the book can be regularly updated, with new versions being released as and when significant changes are incorporated.

## **Challenges and Considerations**

While GitHub offers many advantages, it's essential to understand its limitations:

### **Learning Curve**

For those unfamiliar with Git or version control, there can be an initial learning curve.

### **Data Overwhelm**

With vast amounts of data and contributions, ensuring quality and accuracy can be challenging.

### **Diverse Audience Management**

Catering to both tech-savvy and non-tech audiences might require creating additional resources or tutorials to ensure inclusivity.

## **Conclusion: GitHub – A Paradigm Shift in Collaboration**

The rise of GitHub marks a significant shift in how we perceive and participate in collaborative projects. Its adaptability, transparency, and user-centric design make it a powerful tool, not just for coders but for anyone passionate about collective endeavors. In the realm of mathematics, GitHub promises a future where texts are continually refined, expanded, and enriched by a global community.



# Chapter 6

## Forking Process

### The Heart of Collaboration on GitHub

The beauty of open-source lies in its democratization of content creation. In this section, we demystify the process of "forking" on GitHub, guiding you step-by-step on how to take CiB and create a version uniquely yours.

### Understanding Forking

Before diving into the specifics, it's crucial to understand what "forking" means in the context of GitHub. In the simplest terms, to "fork" a project means to create a personal copy of someone else's project. Forking allows you to freely experiment with changes without affecting the original project. Forking is akin to taking a book you admire and making a copy to write your notes, edits, or additional chapters without altering the original book.

### Why Fork?

#### Experimentation

It provides a safe space where you can test out ideas, make changes, or introduce new content.

#### Personalization

For projects like CiB, it allows readers to customize the content, tailor it to their perspectives, or even localize it for specific audiences.

## **Collaboration**

If you believe your changes have broad appeal, you can propose that they be incorporated back into the original project, enriching it with your unique contributions.

## **Step-by-Step Forking Guide**

### **Set Up Your GitHub Account**

If you don't have an account on GitHub, you'll need to create one. Visit GitHub's official site and sign up.

### **Navigate to the CiB Repository**

Once logged in, search for the CiB project or navigate to its URL directly.

### **Click the 'Fork' Button**

The fork button is located at the top right corner of the repository page; this button will create a copy of CiB in your account.

### **Clone Your Forked Repository**

Forking allows you to have a local copy on your computer, making editing and experimentation easier. Use the command: `git clone [URL of your forked repo]`.

### **Make Your Changes**

Using your preferred tools, introduce the edits, additions, or modifications you desire.

### **Commit and Push Changes**

Once satisfied, save these changes (known as a "commit") and then "push" them to your forked repository on GitHub.

### **Optional – Create a Pull Request**

If you believe your changes should be incorporated into the original CiB repository, you can create a "pull request." A pull request notifies the original authors of your suggestions.

## **Things to Keep in Mind**

### **Stay Updated**

The original CiB project may undergo updates. It's a good practice to regularly "pull" from the original repo to keep your fork up-to-date.

### **Engage with the Community**

Open-source thrives on community interactions. Engage in discussions, seek feedback, and please remain open to constructive criticism.

## **Conclusion: Embracing the Forking Culture**

Forking is more than just a technical process; it symbolizes the ethos of open-source — a world where knowledge is not hoarded but shared, refined, and built upon collectively. By forking CiB or any other project, you're not just creating a personal copy; you're becoming a part of a global movement that values collaboration, innovation, and the shared pursuit of knowledge. So, embark on this journey, make your unique mark, and contribute to the ever-evolving corpus of collective wisdom.





# Chapter 7

## Editing and Customizing

### Tailoring Repositories to Suit Your Needs

Now, let's build upon the forking process; this segment delves into the next steps. How can you edit and customize your version of CiB? What tools and techniques are available at your disposal? Embark on this informative journey as we guide you through the intricacies of editing on GitHub.

### Understanding the GitHub Workspace

Before diving into the specifics of editing, it's essential to familiarize yourself with the GitHub workspace. Think of it as a digital toolshed where each tool serves a unique function:

- **Repository (Repo):** This is the project's main folder where all your project's files are stored and where you track all changes.
- **Branches:** These are parallel versions of a repository, allowing you to work on features or edits without altering the main project.
- **Commits:** This is a saved change in the repository, akin to saving a file after making edits.
- **Pull Requests:** This is how you notify the main project of desired changes, proposing that your edits be merged with the original.

### Editing Files Directly on GitHub

For minor changes, you might opt to edit directly on GitHub:

1. **Navigate to the File:** Within your forked CiB repository, find the file you want to edit.
2. **Click the Pencil Icon:** This button allows you to edit the file.
3. **Make Your Edits:** Modify the content as needed.
4. **Save and Commit:** Below the editing pane, you'll see a "commit changes" section. Add a brief note summarizing your changes and click 'Commit.'

## Editing Files Locally

For extensive customization:

1. **Clone Your Repository:** Use a tool like Git to clone (download) your forked repo to your local computer.
2. **Edit Using Your Preferred Tools:** This could range from text editors to specialized software, depending on the file type.
3. **Commit and Push:** After making your changes, save them (commit) and then upload (push) them to your GitHub repository.

## Utilizing Branches for Extensive Customization

Branches are especially useful for significant overhauls or when working on different versions:

1. **Create a New Branch:** From your main project page, use the branch dropdown to type in a new branch name and create it.
2. **Switch to Your Branch:** Ensure you're working in this new parallel environment.
3. **Make and Commit Changes:** As you would in the main project.
4. **Merging:** Once satisfied with your edits in the branch, you can merge these changes back into the main project or keep them separate as a different version.

## Exploring Additional Tools and Extensions

GitHub's ecosystem is rich with tools and extensions to enhance your editing experience:

- **GitHub Desktop:** An application that simplifies the process of managing your repositories without using command-line tools.
- **Markdown Editors:** Since many GitHub files (like READMEs) are written in Markdown, tools like StackEdit or Dillinger can be invaluable.
- **Extensions for Browsers:** Tools like Octotree can help in navigating repositories more effortlessly.

## Conclusion: The Art of Tailored Content

Editing and customizing on GitHub might seem daunting initially, but with practice, it transforms into a manageable workflow. Many people find that the ability to take a project like CiB and mold it into something uniquely theirs is empowering. It's a testament to the open-source community's ethos, where shared knowledge becomes the canvas and our collective edits, the brushstrokes, crafting an ever-evolving masterpiece. As you embark on your customization journey, remember that every edit, no matter how small, contributes to the project potentially in significant ways.



# Chapter 8

## Engaging with the Community

### Joining the Global Conversation

#### The Significance of the GitHub Community

The digital age has bestowed upon us the gift of connectivity. On platforms like GitHub, this connectivity transcends borders, disciplines, and ideologies, culminating in a melting pot of diverse ideas and knowledge. For mathematicians and math enthusiasts, GitHub offers a space not only to store and manage content but also to engage with an audience that is passionate, informed, and eager to contribute.

#### 1. Discussions and Debates

One of the most enriching aspects of the GitHub community is the plethora of discussions that unfold:

- **Issues:** A core feature of GitHub, "issues" allow users to raise questions, report problems, or propose enhancements.
- **GitHub Discussions:** A newer feature, Discussions, acts like a community forum. It's an excellent place for extended conversations, brainstorming, and sharing ideas or resources.

#### 2. Collaborative Content Creation

Beyond solitary endeavors, GitHub shines in its collaborative capabilities:

- **Pull Requests:** If you have made an alteration to a math text or added a new perspective, pull requests are the way to propose these changes to

the original repository owner. Pull requests foster a collaborative spirit, where content isn't static but continually evolving with community input.

- **Fork and Merge:** As you've learned, forking allows you to create your version of a repository. Engaging with the Community means you can merge changes from others into your fork, blending a mixture of diverse insights.

### 3. Building and Nurturing Networks

Connections made on GitHub often spill over into lasting professional relationships:

- **Following and Followers:** Like on social media platforms, you can follow contributors whose work resonates with you. Following contributors creates a curated feed of updates and also allows you to be part of a more extensive network.
- **GitHub Stars:** If a particular project or repository impresses you, give it a star! Starring not only bookmarks the project for you but also shows appreciation to the creator.

### 4. Learning and Growing Through Feedback

The Community's feedback is an invaluable asset:

- **Code Reviews:** Although traditionally for software, text writers can use this feature to receive feedback on their methodologies or approaches, refining their work.
- **Community Insights:** The "insights" tab on a repository provides analytics. For text writers, this can give a sense of which topics garner more attention and interest.

### 5. Participating in Community Events

GitHub often hosts and sponsors events:

- **Hackathons:** While traditionally for coders, these events can be repurposed for text writer content creation, where participants collaboratively tackle projects or themes.
- **Webinars and Workshops:** These events can range from mastering GitHub's technical side to thematic discussions on math topics.

## **A Project of Collective Wisdom**

Math, in many ways, is a collective endeavor. GitHub can provide a dynamic Community. By engaging with this Community you can become an active participant in the creation of mathematical texts.





# Chapter 9


## Functions

### 9.1 What is a Function?


#### 9.1.1 Definition and Examples

**Definition 1** *A function is a relation between a set of inputs (domain) and a set of permissible outputs (range) with the property that each input is related to exactly one output.*


**Example 1:** Consider the function  $f(x) = x^2$ , which maps each real number to its square.



**Example 2:** Consider the function  $g(x) = \sin(x)$ , which maps real numbers to their sine values.



**Example 3:** Consider the piecewise function  $h(x)$  defined as  $h(x) = \begin{cases} x^2 & \text{if } x \leq 0 \\ \sqrt{x} & \text{if } x > 0 \end{cases}$ .



### 9.1.2 Exercises

**Exercise** Graph the function  $f(x) = 3x - 2$ . Identify its slope and y-intercept.

**Exercise** Consider the function  $g(x) = \frac{1}{x}$ . For what values of  $x$  is  $g(x)$  undefined?

**Exercise** Find the domain and range of the function  $h(x) = \sqrt{x + 4}$ .

**Exercise** Determine if the function  $f(x) = x^3 - x$  is even, odd, or neither. Justify your answer.

**Exercise** Sketch the graph of the piecewise function  $p(x) = \begin{cases} x^2 & \text{if } x < 0 \\ x + 1 & \text{if } x \geq 0 \end{cases}$ .

Indicate any points of discontinuity.

**Exercise** Given the function  $f(x) = 2x^2 - 5x + 3$ , find  $f(-1)$  and  $f(2)$ .

**Exercise** For the function  $f(x) = |x - 3|$ , find the x-coordinate where  $f(x) = 0$ .


**Exercise** Create a function  $f(x)$  that has a domain of all real numbers except  $x = 2$  and a range of  $y \geq 0$ .

**Exercise** Determine the intervals on which the function  $f(x) = -x^2 + 4x - 3$  is increasing and decreasing.

**Exercise** For the function  $g(x) = \cos(x)$ , evaluate  $g(\pi/2)$  and  $g(\pi)$ .

### 9.1.3 Solutions to Exercises

*Solution:* To graph  $f(x) = 3x - 2$ , we note that the slope (m) is 3 and the y-intercept (b) is -2. The graph is a straight line with these characteristics.




*Solution:* The function  $g(x) = \frac{1}{x}$  is undefined when  $x = 0$  because division by zero is not allowed.

*Solution:* For  $h(x) = \sqrt{x + 4}$ , the domain is  $x \geq -4$  (since the expression inside the square root must be non-negative), and the range is  $y \geq 0$  (since the square root produces non-negative results).

*Solution:* The function  $f(x) = x^3 - x$  is odd because  $f(-x) = (-x)^3 - (-x) = -x^3 + x = -f(x)$ .

*Solution:* The piecewise function  $p(x)$  is sketched below. It is discontinuous at  $x = 0$  since the left-hand limit  $\lim_{x \rightarrow 0^-} p(x)$  does not equal the right-hand limit  $\lim_{x \rightarrow 0^+} p(x)$ .



*Solution:* For  $f(x) = 2x^2 - 5x + 3$ , we find  $f(-1) = 2(-1)^2 - 5(-1) + 3 = 10$  and  $f(2) = 2(2)^2 - 5(2) + 3 = -1$ .

*Solution:* The function  $f(x) = |x - 3|$  equals 0 when  $x - 3 = 0$ , i.e.,  $x = 3$ .

*Solution:* One such function is  $f(x) = \frac{1}{(x-2)^2}$ . The denominator ensures that  $x \neq 2$  (domain), and the square ensures all output values are positive (range).


*Solution:* For  $f(x) = -x^2 + 4x - 3$ , the function is increasing where its derivative  $f'(x) = -2x + 4$  is positive, i.e., for  $x < 2$ , and decreasing where  $f'(x)$  is negative, i.e., for  $x > 2$ .

*Solution:* For  $g(x) = \cos(x)$ , we have  $g(\pi/2) = \cos(\pi/2) = 0$  and  $g(\pi) = \cos(\pi) = -1$ .


## 9.2 Domain and Range

**Definition 2** The domain of a function is the set of all possible inputs, while the range is the set of all possible outputs.


**Example 4:** For  $f(x) = \sqrt{x}$ , the domain is  $x \geq 0$  and the range is  $y \geq 0$ . The graph of this function is as follows:



**Example 5:** Consider the function  $g(x) = \frac{1}{x}$ . Its domain is  $x \neq 0$  and its range is  $y \neq 0$ . The graph is given by:



**Example 6:** For the function  $h(x) = x^2 - 4$ , the domain is all real numbers, but the range is  $y \geq -4$ . Its graph is:



### 9.2.1 Exercises on Domain and Range

**Exercise** Determine the domain and range of the function  $f(x) = 2x + 3$ .

**Exercise** Find the domain and range of the function  $g(x) = \frac{1}{x-2}$ .

**Exercise** Identify the domain and range for the quadratic function  $h(x) = x^2 - 6x + 9$ .

**Exercise** For the function  $k(x) = \ln(x-1)$ , determine its domain and range.

**Exercise** Consider the function  $m(x) = \frac{x}{x^2-4}$ . What are its domain and range?

**Exercise** Find the domain and range for the trigonometric function  $n(x) = \sin(x)$ .

**Exercise** Determine the domain and range of the absolute value function  $p(x) = |x + 5|$ .

**Exercise** For the piecewise function  $q(x) = \begin{cases} x^2 & \text{if } x \leq 0 \\ 2x + 3 & \text{if } x > 0 \end{cases}$ , identify the domain and range.

**Exercise** Consider the exponential function  $r(x) = 2^x$ . What is its domain and range?

**Exercise** Identify the domain and range for the cubic function  $s(x) = x^3 - 3x$ .

## 9.2.2 Solutions to Exercises on Domain and Range

*Solution:* For  $f(x) = 2x + 3$ , the domain is all real numbers since there are no restrictions on  $x$ . The range is also all real numbers because as  $x$  takes any real value,  $2x + 3$  also covers all real numbers.

*Solution:* The function  $g(x) = \frac{1}{x-2}$  is undefined when  $x = 2$ , so the domain is  $x \neq 2$ . The range is all real numbers except  $y \neq 0$  because the function never touches the  $y$ -axis.

*Solution:* For the quadratic function  $h(x) = x^2 - 6x + 9$ , the domain is all real numbers. Since the vertex of this parabola is at  $x = 3$  and it opens upwards, the minimum value is  $h(3) = 0$ , thus the range is  $y \geq 0$ .

*Solution:* The function  $k(x) = \ln(x - 1)$  requires  $x - 1 > 0$ , so the domain is  $x > 1$ . The range of a natural logarithm function is all real numbers.

*Solution:* For  $m(x) = \frac{x}{x^2-4}$ , the function is undefined for  $x = \pm 2$ , so the domain is  $x \neq \pm 2$ . The range is all real numbers, as the function can take any  $y$ -value.

*Solution:* The domain of the trigonometric function  $n(x) = \sin(x)$  is all real numbers. The range of the sine function is between  $-1$  and  $1$ , inclusive, so  $y \in [-1, 1]$ .

*Solution:* For the absolute value function  $p(x) = |x+5|$ , the domain is all real numbers. The range is  $y \geq 0$  since absolute values are always non-negative.

*Solution:* The piecewise function  $q(x)$  is defined for all real numbers, so the domain is all real numbers. To find the range, consider each piece:  $x^2$  and  $2x + 3$ . The range is all non-negative values for  $x^2$  and all real numbers for  $2x + 3$ , thus the range is all real numbers.

*Solution:* For the exponential function  $r(x) = 2^x$ , the domain is all real numbers, as exponents can take any real value. The range is  $y > 0$  since exponential functions are always positive.

*Solution:* The cubic function  $s(x) = x^3 - 3x$  has a domain of all real numbers. The range is also all real numbers because as a cubic function, it extends infinitely in both the positive and negative  $y$ -directions.

## 9.3 Types of Functions

### 9.3.1 Linear Functions


**Definition 3** A linear function has the form  $f(x) = mx + b$ , where  $m$  is the slope of the line and  $b$  is the  $y$ -intercept.

Linear functions are fundamental in calculus and represent the simplest form of polynomial functions. Their graph is a straight line, and they have constant rates of change.

**Properties of Linear Functions:**

- The slope  $m$  determines the steepness of the line and its direction (increasing or decreasing).
- The y-intercept  $b$  is the point where the line crosses the y-axis.
- Linear functions have a constant rate of change, which is equal to the slope  $m$ .

**Graphing a Linear Function:** Consider the function  $f(x) = 2x + 3$ . Here,  $m = 2$  and  $b = 3$ . The graph can be plotted as follows:



This line intersects the y-axis at  $(0, 3)$  and has a slope of 2, indicating it rises 2 units for every 1 unit it moves to the right.

**Slope-Intercept Form:** The equation  $f(x) = mx + b$  is known as the slope-intercept form of a linear function, where the slope  $m$  and y-intercept  $b$  are easily identifiable.

**Applications of Linear Functions:** Linear functions model relationships with a constant rate of change and are widely used in various fields such as economics, physics, and social sciences.



**Exercise** Find the slope and y-intercept of the linear function  $g(x) = -3x + 7$  and sketch its graph.

**Exercise** If a linear function passes through the points  $(1, 2)$  and  $(3, 6)$ , determine its equation.

### Exercises on Linear Functions

**Exercise** Find the slope and y-intercept of the linear function  $g(x) = -3x + 7$  and sketch its graph.

**Exercise** If a linear function passes through the points  $(1, 2)$  and  $(3, 6)$ , determine its equation.

**Exercise** Determine the equation of the line that is parallel to  $f(x) = 4x - 5$  and passes through the point  $(2, 3)$ .

**Exercise** Find the point of intersection of the linear functions  $f(x) = 2x + 1$  and  $g(x) = -x + 5$ .

**Exercise** A rental car company charges a base fee of \$50 and an additional \$20 per day. Write a linear function representing the total cost  $C$  as a function of the number of days  $d$  rented.

**Exercise** Graph the linear function  $h(x) = -\frac{1}{2}x + 4$  and identify where it intersects the x-axis and y-axis.

**Exercise** A phone company offers a monthly plan for \$30 with an additional charge of \$0.05 per minute of calls. Write a linear function representing the monthly cost  $M$  as a function of the number of minutes  $m$  used.

**Exercise** Determine whether the points  $(2, 4)$ ,  $(3, 6)$ , and  $(5, 10)$  lie on the same linear function. If so, find the equation of the line.

**Exercise** Find the slope of the line that passes through the points  $(-1, -2)$  and  $(3, 4)$ .

**Exercise** A company's profit  $P$  in thousands of dollars is linearly related to the number of units  $n$  sold, with a profit of \$4,000 for 500 units and a loss of \$2,000 for 200 units. Determine the linear function that models the profit.

### Solutions to Exercises on Linear Functions

*Solution:* For  $g(x) = -3x + 7$ , the slope ( $m$ ) is  $-3$ , and the  $y$ -intercept ( $b$ ) is  $7$ . The graph is a straight line with a downward slope, crossing the  $y$ -axis at  $(0, 7)$ .

*Solution:* To find the equation of a line passing through  $(1, 2)$  and  $(3, 6)$ , calculate the slope  $m = \frac{6-2}{3-1} = 2$ . Using the point-slope form:  $y-2 = 2(x-1)$ , the equation is  $y = 2x$ .

*Solution:* A line parallel to  $f(x) = 4x-5$  has the same slope,  $m = 4$ . Passing through  $(2, 3)$ , use point-slope form:  $y-3 = 4(x-2)$ , yielding  $y = 4x-5$ .

*Solution:* To find the intersection of  $f(x) = 2x+1$  and  $g(x) = -x+5$ , set them equal:  $2x+1 = -x+5$ . Solving gives  $x = \frac{4}{3}$ , and substituting back,  $y = \frac{11}{3}$ . Thus, they intersect at  $(\frac{4}{3}, \frac{11}{3})$ .

*Solution:* The total cost  $C$  is given by  $C = 50 + 20d$ , where  $d$  is the number of days rented.

*Solution:* For  $h(x) = -\frac{1}{2}x + 4$ , the graph intersects the  $y$ -axis at  $(0, 4)$  and intersects the  $x$ -axis at  $x$  where  $h(x) = 0$ . Setting the equation to  $0$  gives  $x = 8$ . So, it intersects the  $x$ -axis at  $(8, 0)$ .

*Solution:* The monthly cost  $M$  is  $M = 30 + 0.05m$ , where  $m$  is the number of minutes used.

*Solution:* The slope between  $(2, 4)$  and  $(3, 6)$  is  $2$ , and between  $(3, 6)$  and  $(5, 10)$  is also  $2$ . Since the slope is consistent, they lie on the same line. The line equation, using point-slope form, is  $y = 2x$ .

*Solution:* The slope is  $\frac{4-(-2)}{3-(-1)} = \frac{6}{4} = \frac{3}{2}$ .

*Solution:* Let  $P = mn + b$ . Using given points, form equations:  $4000 = 500m + b$  and  $-2000 = 200m + b$ . Solving these simultaneously gives  $m = 20$  and  $b = -6000$ . So,  $P = 20n - 6000$ .

### 9.3.2 Polynomial Functions


**Definition 4** A polynomial function is of the form  $f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_0$ , where  $n$  is a non-negative integer, and  $a_n, a_{n-1}, \dots, a_0$  are constants.

Polynomial functions are fundamental in calculus and algebra. They can range from simple linear functions to complex equations with high degrees. The highest power of  $x$  in the function (the degree of the polynomial) greatly influences the shape and behavior of its graph.

**Characteristics of Polynomial Functions:**

- The degree of the polynomial determines its basic shape and the number of roots.
- Polynomials are continuous and smooth functions.
- The leading coefficient  $a_n$  affects the end behavior of the polynomial.

**Graphing Polynomial Functions:** Consider the quadratic function  $f(x) = x^2 - 4x + 3$ . Its graph can be plotted as follows:



This graph represents a parabola opening upwards with its vertex and intercepts easily identifiable.

**Real-World Applications:** Polynomial functions model numerous real-world phenomena, including projectile motion, profit and loss calculations, population growth, and much more.

**Exercise** Sketch the graph of the cubic function  $g(x) = x^3 - 6x^2 + 11x - 6$ . Identify its roots and turning points.

**Exercise** Find a polynomial of degree 3 that has roots at  $x = 1, 2$ , and  $3$ .

**Exercises on Polynomial Functions**

**Exercise** Sketch the graph of the cubic function  $g(x) = x^3 - 6x^2 + 11x - 6$ . Identify its roots and turning points.

**Exercise** Find a polynomial of degree 3 that has roots at  $x = 1, 2$ , and  $3$ .

**Exercise** Determine the degree and leading coefficient of the polynomial  $h(x) = 4x^4 - 3x^3 + 2x^2 - x + 7$ .

**Exercise** For the quadratic function  $f(x) = 2x^2 - 8x + 6$ , find the vertex and axis of symmetry. Also, determine whether it opens upwards or downwards.

**Exercise** Write the polynomial function  $p(x) = x^3 - 4x$  in factored form and identify its zeros.

**Exercise** A polynomial function has zeros at  $x = -2, 0$ , and  $4$  with a leading coefficient of  $1$ . Write the equation of this polynomial.

**Exercise** Given the polynomial  $q(x) = x^4 - 10x^2 + 9$ , find the y-intercept and x-intercepts (if any).

**Exercise** Graph the polynomial  $r(x) = -x^3 + 3x^2 - 2x$  and determine the intervals where the function is increasing and decreasing.

**Exercise** For the polynomial function  $s(x) = 3x^3 - 5x^2 + x - 2$ , use synthetic division to determine whether  $x = 1$  is a zero of  $s$ .

**Exercise** Consider the polynomial  $t(x) = x^4 - 2x^2 - 3$ . Find all the real zeros of  $t$  and sketch the graph of the function.

### Solutions to Exercises on Polynomial Functions

*Solution:* For  $g(x) = x^3 - 6x^2 + 11x - 6$ , the roots are found by solving  $g(x) = 0$ . The roots are  $x = 1, 2, 3$ . Turning points occur where the derivative changes sign. Calculating  $g'(x) = 3x^2 - 12x + 11$  and finding its roots gives the turning points at approximately  $x \approx 1.57$  and  $x \approx 2.43$ .

*Solution:* A polynomial of degree 3 with roots at  $x = 1, 2, 3$  can be written as  $f(x) = (x - 1)(x - 2)(x - 3)$ .

*Solution:* The degree of the polynomial  $h(x) = 4x^4 - 3x^3 + 2x^2 - x + 7$  is 4, and the leading coefficient is 4.

*Solution:* For  $f(x) = 2x^2 - 8x + 6$ , the vertex form is  $f(x) = 2(x - 2)^2 + 2$ . The vertex is at  $(2, 2)$ , and the axis of symmetry is  $x = 2$ . The parabola opens upwards since the coefficient of  $x^2$  is positive.

*Solution:* The polynomial  $p(x) = x^3 - 4x$  can be factored as  $p(x) = x(x^2 - 4) = x(x - 2)(x + 2)$ . The zeros are at  $x = 0, 2, -2$ .

*Solution:* A polynomial with zeros at  $x = -2, 0$ , and  $4$  and a leading coefficient of  $1$  is  $f(x) = (x + 2)x(x - 4) = x^3 - 4x^2 - 2x$ .

*Solution:* For  $q(x) = x^4 - 10x^2 + 9$ , the y-intercept is  $q(0) = 9$ . To find the x-intercepts, solve  $x^4 - 10x^2 + 9 = 0$ , which gives  $x \approx \pm 3.16, \pm 0.84$ .

*Solution:* Graphing  $r(x) = -x^3 + 3x^2 - 2x$ , the function increases where  $r'(x) > 0$  and decreases where  $r'(x) < 0$ . Calculating  $r'(x) = -3x^2 + 6x - 2$  and finding its roots gives the intervals of increase and decrease.

*Solution:* Using synthetic division to divide  $3x^3 - 5x^2 + x - 2$  by  $x - 1$ , we find that the remainder is not zero. Hence,  $x = 1$  is not a zero of  $s(x)$ .

*Solution:* To find the real zeros of  $t(x) = x^4 - 2x^2 - 3$ , solve  $x^4 - 2x^2 - 3 = 0$ . The real zeros are approximately  $x \approx \pm 1.73, \pm 1.00$ . Graphing  $t(x)$  would show a polynomial crossing the x-axis at these points.

### 9.3.3 Rational Functions


**Definition 5** A rational function is the ratio of two polynomial functions, typically expressed in the form  $f(x) = \frac{p(x)}{q(x)}$ , where both  $p(x)$  and  $q(x)$  are polynomial functions and  $q(x) \neq 0$ .

Rational functions can exhibit a variety of behaviors, including asymptotes and discontinuities, making them interesting subjects of study in calculus.

#### Characteristics of Rational Functions:

- The vertical asymptotes occur at values of  $x$  for which  $q(x) = 0$ , provided these points are not cancelled out by the numerator.
- Horizontal asymptotes are determined by the degrees of  $p(x)$  and  $q(x)$ .
- Rational functions may have holes if a factor in the denominator is cancelled out by the numerator.

**Graphing Rational Functions:** Consider the function  $f(x) = \frac{x^2 - 4}{x - 2}$ . Here, we have a hole at  $x = 2$  because the factor  $(x - 2)$  is cancelled. The graph can be plotted as follows:



**Applications of Rational Functions:** Rational functions are used in various fields, including engineering, economics, and physics, to model scenarios where variables inversely affect each other.

**Exercise** Graph the rational function  $g(x) = \frac{1}{x-1}$  and determine its vertical and horizontal asymptotes.

**Exercise** Consider the function  $h(x) = \frac{x^2-1}{x^2+x-2}$ . Identify its vertical and horizontal asymptotes, and sketch the graph.

### Exercises on Rational Functions

**Exercise** Graph the rational function  $g(x) = \frac{1}{x-1}$  and determine its vertical and horizontal asymptotes.

**Exercise** Consider the function  $h(x) = \frac{x^2-1}{x^2+x-2}$ . Identify its vertical and horizontal asymptotes, and sketch the graph.

**Exercise** Find the domain of the rational function  $f(x) = \frac{2x-3}{x^2-4}$ .

**Exercise** Determine the vertical and horizontal asymptotes for the function  $p(x) = \frac{x^3-4x}{x^2-5x+6}$  and sketch its graph.

**Exercise** For the rational function  $q(x) = \frac{x^2+2x+1}{x^2-1}$ , identify any holes or asymptotes, and graph the function.

**Exercise** Given  $r(x) = \frac{3x-5}{2x+4}$ , calculate  $r(2)$  and determine the behavior of  $r(x)$  as  $x$  approaches infinity.

**Exercise** Sketch the graph of the function  $s(x) = \frac{x}{x^2-9}$  and find the x- and y-intercepts.

**Exercise** Analyze the rational function  $t(x) = \frac{5x-4}{x-3}$  for any asymptotes or discontinuities and provide its graph.

**Exercise** Find the range of the function  $u(x) = \frac{2}{x-1} + 3$ .

**Exercise** Determine the inverse of the function  $v(x) = \frac{1}{2-x}$ .

### Solutions to Exercises on Rational Functions

*Solution:* For  $g(x) = \frac{1}{x-1}$ , the vertical asymptote is at  $x = 1$  (where the denominator equals zero), and the horizontal asymptote is  $y = 0$  (as  $x$  approaches infinity,  $g(x)$  approaches 0).

*Solution:* In  $h(x) = \frac{x^2-1}{x^2+x-2}$ , the vertical asymptotes are at the roots of the denominator,  $x = 1$  and  $x = -2$ . The horizontal asymptote is  $y = 1$  as the degrees of the numerator and denominator are the same.

*Solution:* The domain of  $f(x) = \frac{2x-3}{x^2-4}$  is all real numbers except  $x = \pm 2$ , where the denominator equals zero.

*Solution:* For  $p(x) = \frac{x^3-4x}{x^2-5x+6}$ , vertical asymptotes are at  $x = 2$  and  $x = 3$ , the roots of the denominator. There is no horizontal asymptote as the degree of the numerator is greater than that of the denominator.

*Solution:* In  $q(x) = \frac{x^2+2x+1}{x^2-1}$ , there is a hole at  $x = -1$  (factor cancelled in numerator and denominator), vertical asymptotes at  $x = \pm 1$ , and no horizontal asymptote.

*Solution:* For  $r(x) = \frac{3x-5}{2x+4}$ ,  $r(2) = \frac{1}{8}$ . As  $x$  approaches infinity,  $r(x)$  approaches  $\frac{3}{2}$ .

*Solution:* In  $s(x) = \frac{x}{x^2-9}$ , the x-intercepts are at  $x = 0$ , y-intercept at  $y = 0$ , and vertical asymptotes at  $x = \pm 3$ .

*Solution:* The function  $t(x) = \frac{5x-4}{x-3}$  has a vertical asymptote at  $x = 3$  and no horizontal asymptote. Discontinuity occurs at  $x = 3$ .

*Solution:* To find the range of  $u(x) = \frac{2}{x-1} + 3$ , consider the behavior as  $x$  approaches 1 and infinity. The range is all real numbers except  $y = 3$ .

*Solution:* The inverse of  $v(x) = \frac{1}{2-x}$  can be found by swapping  $x$  and  $y$  and solving for  $y$ :  $x = \frac{1}{2-y}$  leads to  $y = 2 - \frac{1}{x}$ .

### 9.3.4 Trigonometric Functions


**Definition 6** *Trigonometric functions include  $\sin(x)$ ,  $\cos(x)$ , and  $\tan(x)$ , relating angles to ratios of sides in right triangles.*

Trigonometric functions are fundamental in calculus, physics, and engineering. They are periodic functions and have various properties that are essential in the study of waves, oscillations, and circular motion.

#### Properties of Trigonometric Functions:

- The functions  $\sin(x)$  and  $\cos(x)$  are periodic with a period of  $2\pi$ .
- The function  $\tan(x)$  is periodic with a period of  $\pi$ .
- These functions have specific symmetries:  $\sin(x)$  is odd,  $\cos(x)$  is even, and  $\tan(x)$  is odd.

**Graphing Trigonometric Functions:** Using ‘pgfplots’ to graph  $\sin(x)$  and  $\cos(x)$ :



**Applications of Trigonometric Functions:** Trigonometric functions model phenomena such as sound waves, light waves, and describe the motion of pendulums and other periodic phenomena.

**Exercise** Graph the function  $f(x) = \tan(x)$  and identify its asymptotes.

**Exercise** Find the amplitude and period of the function  $g(x) = 3 \sin(2x)$ .

**Exercise** Determine the phase shift and vertical shift for the function  $h(x) = \cos(x - \pi/2) + 1$ .



**Exercises on Trigonometric Functions**

**Exercise** Graph the function  $f(x) = \tan(x)$  and identify its asymptotes.

**Exercise** Find the amplitude and period of the function  $g(x) = 3 \sin(2x)$ .

**Exercise** Determine the phase shift and vertical shift for the function  $h(x) = \cos(x - \pi/2) + 1$ .

**Exercise** Prove that  $\sin^2(x) + \cos^2(x) = 1$  using a unit circle.

**Exercise** Evaluate  $\sin(\pi/6)$ ,  $\cos(\pi/3)$ , and  $\tan(\pi/4)$ .

**Exercise** Sketch the graph of  $f(x) = \sin(x) - \cos(x)$  over the interval  $[0, 2\pi]$ .

**Exercise** Find the values of  $x$  where  $\sin(x) = -\sqrt{3}/2$  in the interval  $[-\pi, \pi]$ .

**Exercise** For the function  $g(x) = 2 \cos(3x - \pi)$ , identify the amplitude, period, phase shift, and sketch its graph.

**Exercise** Determine the exact value of  $\tan(\pi/6) + \cos(\pi/3)$ .

**Exercise** Solve for  $x$  in the equation  $3 \sin(x) + 4 \cos(x) = 0$  over the interval  $[0, 2\pi]$ .

**Solutions to Exercises on Trigonometric Functions**

*Solution:* For  $f(x) = \tan(x)$ , the graph has vertical asymptotes at  $x = \pm \frac{\pi}{2}, \pm \frac{3\pi}{2}, \dots$  where the cosine function in the denominator equals zero.

*Solution:* The function  $g(x) = 3 \sin(2x)$  has an amplitude of 3 (the coefficient of the sine function) and a period of  $\frac{\pi}{2}$  (computed as  $\frac{2\pi}{2} = \pi$ ).

*Solution:* For  $h(x) = \cos(x - \pi/2) + 1$ , the phase shift is  $\frac{\pi}{2}$  to the right, and the vertical shift is 1 unit upwards.

*Solution:* Using the unit circle, we know that  $\sin(x)$  and  $\cos(x)$  represent the y-coordinate and x-coordinate, respectively, of a point on the circle. Since the radius of the unit circle is 1, we have  $\sin^2(x) + \cos^2(x) = 1^2 = 1$ .

*Solution:* Evaluate  $\sin(\pi/6) = 1/2$ ,  $\cos(\pi/3) = 1/2$ , and  $\tan(\pi/4) = 1$ .

*Solution:* To graph  $f(x) = \sin(x) - \cos(x)$ , plot the sine and cosine functions separately and then subtract their values for each  $x$  in the interval  $[0, 2\pi]$ .

*Solution:* The values of  $x$  where  $\sin(x) = -\sqrt{3}/2$  in the interval  $[-\pi, \pi]$  are  $x = -\pi/3$  and  $x = -2\pi/3$ .

*Solution:* For  $g(x) = 2\cos(3x - \pi)$ , the amplitude is 2, the period is  $\frac{2\pi}{3}$ , and the phase shift is  $\frac{\pi}{3}$  to the right. Sketch the graph accordingly.

*Solution:* The exact value of  $\tan(\pi/6) + \cos(\pi/3)$  is  $\frac{\sqrt{3}}{3} + \frac{1}{2}$ .

*Solution:* Solve  $3\sin(x) + 4\cos(x) = 0$  over the interval  $[0, 2\pi]$ . Dividing by  $\cos(x)$  and applying trigonometric identities, we find  $x = \frac{\pi}{2}, \frac{3\pi}{2}$ .

### 9.3.5 Exponential and Logarithmic Functions

**Definition 7** *Exponential functions have the form  $f(x) = a^x$  where  $a$  is a positive constant. Logarithmic functions are the inverses of exponential functions and are usually expressed as  $g(x) = \log_a(x)$  for a logarithm to the base  $a$ .*

Exponential and logarithmic functions are widely used in many fields, including physics, engineering, and economics, due to their properties of describing growth and decay processes.


#### Characteristics of Exponential Functions:

- Exponential growth occurs when  $a > 1$ , and exponential decay happens when  $0 < a < 1$ .
- The graph of an exponential function always lies above the x-axis and increases or decreases rapidly.
- Exponential functions have a horizontal asymptote at  $y = 0$ .

#### Characteristics of Logarithmic Functions:

- The logarithmic function  $\log_a(x)$  is undefined for  $x \leq 0$ .
- The graph of a logarithmic function passes through the point  $(1, 0)$  and has a vertical asymptote at  $x = 0$ .
- Logarithmic functions increase slowly and are used to model the inverse of exponential growth or decay.

**Graphing Exponential and Logarithmic Functions:** Here we graph  $f(x) = 2^x$  and its inverse  $g(x) = \log_2(x)$  using ‘pgfplots’:



**Exercise** Graph the exponential function  $h(x) = e^x$  and identify its key characteristics.

**Exercise** Sketch the graph of  $k(x) = \log_e(x)$  (also known as  $\ln(x)$ ) and discuss its properties.

### Exercises on Exponential and Logarithmic Functions

**Exercise** Graph the exponential function  $h(x) = e^x$  and identify its key characteristics.

**Exercise** Sketch the graph of  $k(x) = \log_e(x)$  (also known as  $\ln(x)$ ) and discuss its properties.

**Exercise** Evaluate the expression  $e^{\ln(5)}$  and explain the relationship between the exponential and logarithmic functions.

**Exercise** Find the solution to the equation  $3^x = 27$ .

**Exercise** Determine the x-intercept of the logarithmic function  $f(x) = \log_2(x - 1)$ .

**Exercise** Solve for  $x$  in the equation  $\ln(x) + \ln(2x) = 3$ .

**Exercise** Graph the function  $g(x) = 2^x + 3$  and describe how it differs from the basic exponential function  $f(x) = 2^x$ .

**Exercise** Find the domain and range of the function  $h(x) = \ln(x - 2)$ .

**Exercise** Calculate the derivative of the function  $f(x) = e^{2x}$  using the rules of differentiation for exponential functions.

**Exercise** Given the function  $g(x) = \log_{10}(x^2 - 1)$ , find the points of discontinuity.

### Solutions to Exercises on Exponential and Logarithmic Functions

*Solution:* For  $h(x) = e^x$ , the graph is an increasing exponential function. It passes through the point  $(0, 1)$  and has a horizontal asymptote at  $y = 0$ . The function is always positive and increases rapidly as  $x$  increases.

*Solution:* The function  $k(x) = \log_e(x)$ , or  $\ln(x)$ , has a graph that passes through the point  $(1, 0)$  and has a vertical asymptote at  $x = 0$ . The function is defined for  $x > 0$  and increases slowly as  $x$  increases.

*Solution:* Evaluate  $e^{\ln(5)}$ . Since the exponential and logarithmic functions are inverses of each other,  $e^{\ln(5)} = 5$ .

*Solution:* Solve  $3^x = 27$ . Recognizing that  $27 = 3^3$ , we find  $x = 3$ .

*Solution:* To find the x-intercept of  $f(x) = \log_2(x - 1)$ , set  $f(x) = 0$ . This yields  $\log_2(x - 1) = 0$ , so  $x - 1 = 1$ , and thus  $x = 2$ .

*Solution:* Solve  $\ln(x) + \ln(2x) = 3$ . Combine the logarithms:  $\ln(2x^2) = 3$ . Exponentiating both sides gives  $2x^2 = e^3$ . Solving for  $x$  yields  $x = \pm\sqrt{\frac{e^3}{2}}$ , but only the positive solution is valid since  $\ln(x)$  is undefined for negative  $x$ .

*Solution:* Graph  $g(x) = 2^x + 3$ . This graph is a vertical shift of the function  $f(x) = 2^x$  upwards by 3 units. It increases exponentially and passes through the point  $(0, 4)$ .

*Solution:* The domain of  $h(x) = \ln(x - 2)$  is  $x > 2$ , and the range is all real numbers since the logarithmic function can take any real value.

*Solution:* The derivative of  $f(x) = e^{2x}$  is  $f'(x) = 2e^{2x}$ , using the chain rule for differentiation.

*Solution:* For  $g(x) = \log_{10}(x^2 - 1)$ , the function is undefined at points where  $x^2 - 1 \leq 0$ , which occurs at  $x = -1$  and  $x = 1$ . These are the points of discontinuity.

## 9.4 Function Properties

### 9.4.1 Continuity


**Definition 8** A function is continuous at a point if the limit of the function as it approaches the point is equal to the function value at that point. Formally, a function  $f(x)$  is continuous at a point  $c$  if  $\lim_{x \rightarrow c} f(x) = f(c)$ .

Continuity is a fundamental concept in calculus, essential for understanding limits, derivatives, and integrals.

#### Types of Discontinuities:

- A function has a *removable discontinuity* at a point if the limit exists but is not equal to the function value at that point.
- A *jump discontinuity* occurs when the left-hand and right-hand limits exist but are not equal to each other.
- An *infinite discontinuity* occurs when the function approaches infinity near the point of discontinuity.

**Graphing Continuous Functions:** To illustrate continuity, consider the function  $f(x) = \frac{x^2-4}{x-2}$  and its simplified form  $g(x) = x+2$ . The function  $f(x)$  has a removable discontinuity at  $x = 2$ .



**Applications of Continuity:** Continuous functions are used in various applications, including physics (to model continuous motion), economics (to describe continuous growth), and engineering (in signal processing).

**Exercise** Determine if the function  $h(x) = \frac{x^2-1}{x-1}$  is continuous at  $x = 1$ .

**Exercise** Evaluate the continuity of the piecewise function  $k(x) = \begin{cases} x^2 & \text{if } x < 2 \\ 4 & \text{if } x = 2 \\ x + 2 & \text{if } x > 2 \end{cases}$  at  $x = 2$ .

### Exercises on Continuity

**Exercise** Determine if the function  $h(x) = \frac{x^2-1}{x-1}$  is continuous at  $x = 1$ .

**Exercise** Evaluate the continuity of the piecewise function  $k(x) = \begin{cases} x^2 & \text{if } x < 2 \\ 4 & \text{if } x = 2 \\ x + 2 & \text{if } x > 2 \end{cases}$  at  $x = 2$ .

**Exercise** Is the function  $f(x) = \frac{1}{x}$  continuous at  $x = 0$ ? Justify your answer.

**Exercise** Consider the function  $g(x) = \ln(x)$ . Determine the points of discontinuity, if any.

**Exercise** Prove that the function  $p(x) = x^3 - 3x + 1$  is continuous for all real numbers.

**Exercise** Find the value of  $a$  for which the function  $q(x) = \begin{cases} 2x + a & \text{if } x \leq 3 \\ x^2 - 1 & \text{if } x > 3 \end{cases}$  is continuous at  $x = 3$ .

**Exercise** Determine whether the function  $r(x) = \frac{\sin(x)}{x}$  is continuous at  $x = 0$ .

**Exercise** For the function  $s(x) = \sqrt{x}$ , identify the intervals where it is continuous.

**Exercise** Show that the function  $t(x) = e^{-x^2}$  is continuous everywhere.

**Exercise** Is the function  $u(x) = \frac{|x|}{x}$  continuous at  $x = 0$ ? Explain your reasoning.

**Solutions to Exercises on Continuity**

*Solution:* For  $h(x) = \frac{x^2-1}{x-1}$ , factorize the numerator to get  $h(x) = \frac{(x-1)(x+1)}{x-1}$ . At  $x = 1$ , the function simplifies to  $h(1) = 2$ , and the limit as  $x$  approaches 1 is also 2. Hence,  $h(x)$  is continuous at  $x = 1$ .

*Solution:* For  $k(x)$ , at  $x < 2$ ,  $k(x) = x^2$ ; at  $x = 2$ ,  $k(x) = 4$ ; and at  $x > 2$ ,  $k(x) = x + 2$ . The left-hand limit as  $x$  approaches 2 is 4, and the right-hand limit is also 4. Since the function value at  $x = 2$  is 4,  $k(x)$  is continuous at  $x = 2$ .

*Solution:* The function  $f(x) = \frac{1}{x}$  is not continuous at  $x = 0$  because it is not defined there (the denominator becomes zero), and the limits as  $x$  approaches 0 are  $\pm\infty$ .

*Solution:* The function  $g(x) = \ln(x)$  is continuous for all  $x > 0$ . It has a discontinuity at  $x = 0$  because it is not defined for  $x \leq 0$ .

*Solution:* Since  $p(x) = x^3 - 3x + 1$  is a polynomial function, it is continuous for all real numbers (polynomials are continuous everywhere).

*Solution:* For  $q(x)$  to be continuous at  $x = 3$ , the values from the left and right must be equal at  $x = 3$ . Setting  $2(3) + a = 3^2 - 1$  gives  $a = 5$ .

*Solution:* The function  $r(x) = \frac{\sin(x)}{x}$  is continuous at  $x = 0$  by applying L'Hôpital's Rule or recognizing that  $\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$ .

*Solution:* The function  $s(x) = \sqrt{x}$  is continuous for all  $x \geq 0$  since it is defined and smooth over this interval.

*Solution:* The function  $t(x) = e^{-x^2}$  is continuous everywhere as the composition of continuous functions (exponential and polynomial) is continuous.

*Solution:* The function  $u(x) = \frac{|x|}{x}$  is not continuous at  $x = 0$  as it is not defined there (the denominator becomes zero).

**9.4.2 Differentiability**


**Definition 9** *A function is differentiable at a point if it has a defined derivative at that point. Formally, a function  $f(x)$  is differentiable at  $c$  if the limit  $\lim_{h \rightarrow 0} \frac{f(c+h)-f(c)}{h}$  exists.*

Differentiability is a key concept in calculus, relating closely to the continuity of a function and its smoothness.

**Characteristics of Differentiable Functions:**

- If a function is differentiable at a point, it is also continuous at that point, but the converse is not necessarily true.
- A function may fail to be differentiable at a point if it has a corner, cusp, or vertical tangent, or is discontinuous at that point.

**Graphing and Analyzing Differentiability:** Consider a function  $f(x)$  with a cusp at  $x = 0$ . For example,  $f(x) = |x|$  is not differentiable at  $x = 0$  due to the cusp.



**Applications of Differentiability:** Differentiability is used in physics to analyze motion (velocity and acceleration) and in economics to find rates of change (marginal costs and revenues).

**Exercise** Determine whether the function  $g(x) = x^2$  is differentiable at  $x = 0$  and explain why.

**Exercise** Analyze the differentiability of the piecewise function  $h(x) = \begin{cases} x^2 & \text{if } x \leq 1 \\ 2x - 1 & \text{if } x > 1 \end{cases}$  at  $x = 1$ .

**Exercises on Differentiability**

**Exercise** Determine whether the function  $g(x) = x^2$  is differentiable at  $x = 0$  and explain why.



**Exercise** Analyze the differentiability of the piecewise function  $h(x) = \begin{cases} x^2 & \text{if } x \leq 1 \\ 2x - 1 & \text{if } x > 1 \end{cases}$  at  $x = 1$ .

**Exercise** Check if the function  $f(x) = \frac{1}{x}$  is differentiable at  $x = 0$  and provide a justification for your answer.

**Exercise** Determine whether the function  $k(x) = |x|$  is differentiable at  $x = 0$ . Give reasons for your answer.

**Exercise** For the function  $p(x) = x^3$ , find the derivative at  $x = 2$  and discuss its differentiability at this point.

**Exercise** Consider the function  $q(x) = \begin{cases} x^2 \sin(\frac{1}{x}) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$ . Is  $q(x)$  differentiable at  $x = 0$ ?

**Exercise** Evaluate the differentiability of the function  $r(x) = \sqrt{x}$  at  $x = 0$ .

**Exercise** Find the points of non-differentiability, if any, for the function  $s(x) = \frac{|x|}{x}$ .

**Exercise** Given the function  $t(x) = \ln(x)$ , determine its differentiability over its domain.

**Exercise** Prove that the function  $u(x) = e^x$  is differentiable for all real numbers  $x$ .

### Solutions to Exercises on Differentiability

*Solution:* The function  $g(x) = x^2$  is differentiable at  $x = 0$ . This is because its derivative  $g'(x) = 2x$  is defined at  $x = 0$  (with  $g'(0) = 0$ ), showing that the function is smooth and has no sharp corners or cusps at this point.

*Solution:* The piecewise function  $h(x)$  is differentiable at  $x = 1$  as both pieces of the function have the same derivative at this point. For  $x \leq 1$ , the derivative is  $2x$ , and for  $x > 1$ , it is 2. Since both derivatives equal 2 at  $x = 1$ , the function is differentiable there.

*Solution:* The function  $f(x) = \frac{1}{x}$  is not differentiable at  $x = 0$  because it is not defined at this point. The function has a discontinuity (infinite discontinuity) at  $x = 0$ , which precludes differentiability.

*Solution:* The function  $k(x) = |x|$  is not differentiable at  $x = 0$ . Although the function is continuous at  $x = 0$ , it has a sharp corner at this point, which means that the limit defining the derivative does not exist.

*Solution:* For  $p(x) = x^3$ , the derivative is  $p'(x) = 3x^2$ . At  $x = 2$ ,  $p'(2) = 12$ . The function is differentiable at this point as the derivative is defined and finite.

*Solution:* The function  $q(x)$  is differentiable at  $x = 0$ . Although the expression  $x^2 \sin(\frac{1}{x})$  seems complex, its derivative at  $x = 0$  is 0, making the function differentiable at this point.

*Solution:* The function  $r(x) = \sqrt{x}$  is not differentiable at  $x = 0$ . The derivative,  $\frac{1}{2\sqrt{x}}$ , becomes infinite as  $x$  approaches 0, indicating a vertical tangent line at this point.

*Solution:* The function  $s(x) = \frac{|x|}{x}$  is not differentiable at  $x = 0$  as it is not continuous there. The function has a jump discontinuity at this point.

*Solution:* The function  $t(x) = \ln(x)$  is differentiable over its domain  $(0, \infty)$ . Its derivative,  $\frac{1}{x}$ , is defined for all  $x > 0$ .

*Solution:* The function  $u(x) = e^x$  is differentiable for all real numbers  $x$  because its derivative,  $e^x$ , is defined and continuous for all  $x$ .

### 9.4.3 Asymptotic Behavior


**Definition 10** *Asymptotic behavior refers to the behavior of a function as it approaches infinity or a particular value. A function may approach a finite value (horizontal asymptote), become infinitely large (vertical asymptote), or approach a certain path (slant asymptote).*

Understanding asymptotes is crucial in calculus for analyzing function limits and behavior at extreme values.

#### Types of Asymptotes:

- *Horizontal asymptotes* occur when a function approaches a horizontal line as  $x$  goes to infinity or minus infinity.
- *Vertical asymptotes* occur at values of  $x$  where the function tends towards infinity or minus infinity.
- *Slant (or oblique) asymptotes* occur when the function approaches a line that is not horizontal as  $x$  goes to infinity or minus infinity.

**Graphing Asymptotic Behavior:** Consider the function  $f(x) = \frac{2x}{x-1}$ . It has a vertical asymptote at  $x = 1$  and a horizontal asymptote at  $y = 2$ .



**Applications of Asymptotic Analysis:** Asymptotic analysis is used in various fields, such as physics for studying motion at high velocities, in economics for long-term growth models, and in probability for extreme value theory.

**Exercise** Determine the horizontal and vertical asymptotes of the function  $g(x) = \frac{3x^2-x}{x^2+2}$ .

**Exercise** Analyze the asymptotic behavior of the function  $h(x) = \frac{x^2+3x}{x+1}$ .

### Exercises on Asymptotic Behavior

**Exercise** Determine the horizontal and vertical asymptotes of the function  $g(x) = \frac{3x^2-x}{x^2+2}$ .

**Exercise** Analyze the asymptotic behavior of the function  $h(x) = \frac{x^2+3x}{x+1}$ .

**Exercise** Find all asymptotes of the function  $f(x) = \frac{4x^3-2x+1}{2x^2-3x+5}$ .

**Exercise** Determine the horizontal asymptote, if any, for the function  $p(x) = e^x$ .

**Exercise** Identify any vertical asymptotes for the function  $q(x) = \ln(x-2)$ .

**Exercise** For the function  $r(x) = \frac{x}{x^2-4}$ , describe its asymptotic behavior as  $x$  approaches 2 and -2.

**Exercise** Calculate the slant asymptote for the function  $s(x) = \frac{x^2-x+1}{x-3}$ .

**Exercise** Examine the function  $t(x) = \frac{1}{\sqrt{x^2+1}}$  for horizontal asymptotes.

**Exercise** Does the function  $u(x) = \frac{\sin(x)}{x}$  have any horizontal asymptotes as  $x$  approaches infinity?

**Exercise** Analyze the function  $v(x) = x \sin(\frac{1}{x})$  for  $x$  approaching 0 and determine if there are any asymptotes.

### Solutions to Exercises on Asymptotic Behavior

*Solution:* For  $g(x) = \frac{3x^2-x}{x^2+2}$ , the horizontal asymptote is found by comparing the degrees of the numerator and denominator. Since they are the same, the horizontal asymptote is  $y = \frac{3}{1} = 3$ . There are no vertical asymptotes as the denominator is never zero.

*Solution:* The function  $h(x) = \frac{x^2+3x}{x+1}$  has a slant asymptote since the degree of the numerator is one more than the degree of the denominator. Dividing the numerator by the denominator gives the slant asymptote  $y = x + 2$ . There is no vertical asymptote as  $x + 1$  does not equal zero for any real  $x$ .

*Solution:* For  $f(x) = \frac{4x^3-2x+1}{2x^2-3x+5}$ , the degree of the numerator is higher than the degree of the denominator, so there is no horizontal asymptote. For vertical asymptotes, solve  $2x^2 - 3x + 5 = 0$ , which has no real solutions, so there are no vertical asymptotes.

*Solution:* The function  $p(x) = e^x$  does not have a horizontal asymptote as it grows exponentially and does not approach a finite value as  $x$  approaches infinity.

*Solution:* The function  $q(x) = \ln(x - 2)$  has a vertical asymptote at  $x = 2$  since the logarithm becomes undefined at this point.

*Solution:* For  $r(x) = \frac{x}{x^2-4}$ , the function has vertical asymptotes at  $x = 2$  and  $x = -2$  where the denominator equals zero.

*Solution:* The function  $s(x) = \frac{x^2-x+1}{x-3}$  has a slant asymptote since the degree of the numerator is one more than the denominator. Divide  $x^2 - x + 1$  by  $x - 3$  to find the slant asymptote equation.

*Solution:* The function  $t(x) = \frac{1}{\sqrt{x^2+1}}$  has a horizontal asymptote at  $y = 0$  as the value approaches zero when  $x$  approaches infinity.

*Solution:* The function  $u(x) = \frac{\sin(x)}{x}$  has a horizontal asymptote at  $y = 0$  as  $x$  approaches infinity since the sine function oscillates between -1 and 1 while  $x$  grows without bound.

*Solution:* The function  $v(x) = x \sin(\frac{1}{x})$  for  $x$  approaching 0 does not have any asymptotes. As  $x$  approaches 0,  $\sin(\frac{1}{x})$  oscillates rapidly, but the product with  $x$  ensures that the function remains bounded and approaches 0.

### 9.4.4 Periodicity and Symmetry


**Definition 11** *A function is periodic if it repeats its values at regular intervals, and it is symmetric if it exhibits mirror symmetry about an axis or point.*


Periodicity and symmetry play important roles in understanding the behavior and properties of functions, especially in the fields of trigonometry and wave analysis.

#### Types of Symmetry:

- A function is *even* if  $f(-x) = f(x)$ , showing symmetry about the  $y$ -axis.
- A function is *odd* if  $f(-x) = -f(x)$ , showing rotational symmetry about the origin.

**Graphing Periodic and Symmetric Functions:** For instance,  $\sin(x)$  is an example of a periodic function, and  $x^2$  is an example of an even function.





**Applications of Periodicity and Symmetry:** These concepts are frequently applied in physics for analyzing wave patterns, in engineering for signal processing, and in mathematics for solving trigonometric equations.

**Exercise** Determine if the function  $f(x) = \cos(x)$  is periodic, and if so, find its period.

**Exercise** Analyze the symmetry of the function  $g(x) = x^3 - x$ . Is it even, odd, or neither?

### Exercises on Periodicity and Symmetry

**Exercise** Determine if the function  $f(x) = \cos(x)$  is periodic, and if so, find its period.

**Exercise** Analyze the symmetry of the function  $g(x) = x^3 - x$ . Is it even, odd, or neither?

**Exercise** Identify the periodicity of the function  $h(x) = \sin(2x)$  and specify the period.

**Exercise** Check whether the function  $p(x) = x^4 - 6x^2 + 8$  is even, odd, or neither.

**Exercise** For the function  $q(x) = \tan(x)$ , determine if it is periodic and find the period.

**Exercise** Investigate the symmetry of the function  $r(x) = \frac{1}{x}$ . Is it symmetric about the y-axis, the origin, or neither?

**Exercise** Examine the function  $s(x) = e^x$  for periodicity. Is this function periodic?

**Exercise** Determine the period of the function  $t(x) = \cos(3x - \pi)$ .

**Exercise** Analyze the function  $u(x) = x^2 - 2x + 1$  for symmetry. Is it symmetric about the y-axis?

**Exercise** Check whether the function  $v(x) = \sin(x) + \cos(x)$  is periodic, and if so, identify its period.

### Solutions to Exercises on Periodicity and Symmetry

*Solution:* The function  $f(x) = \cos(x)$  is periodic. Its period is  $2\pi$  since  $\cos(x + 2\pi) = \cos(x)$  for all  $x$ .

*Solution:* The function  $g(x) = x^3 - x$  is odd. This is because  $g(-x) = (-x)^3 - (-x) = -x^3 + x = -g(x)$ .

*Solution:* The function  $h(x) = \sin(2x)$  is periodic with a period of  $\pi$ . This is because  $\sin(2(x + \pi)) = \sin(2x + 2\pi) = \sin(2x)$ .

*Solution:* The function  $p(x) = x^4 - 6x^2 + 8$  is even since  $p(-x) = (-x)^4 - 6(-x)^2 + 8 = x^4 - 6x^2 + 8 = p(x)$ .

*Solution:* The function  $q(x) = \tan(x)$  is periodic with a period of  $\pi$  because  $\tan(x + \pi) = \tan(x)$  for all  $x$ .

*Solution:* The function  $r(x) = \frac{1}{x}$  is neither even nor odd, and it does not exhibit symmetry about the y-axis or the origin.

*Solution:* The function  $s(x) = e^x$  is not periodic. It does not repeat its values at regular intervals.

*Solution:* The function  $t(x) = \cos(3x - \pi)$  is periodic with a period of  $\frac{2\pi}{3}$ . This period is derived from the coefficient of  $x$  in the argument of the cosine function.

*Solution:* The function  $u(x) = x^2 - 2x + 1$  is not symmetric about the y-axis. It is a parabola that opens upwards and does not have symmetry about the y-axis.

*Solution:* The function  $v(x) = \sin(x) + \cos(x)$  is periodic with a period of  $2\pi$  since both sine and cosine functions have a period of  $2\pi$ .

## 9.5 Function Transformations

### 9.5.1 Translation and Scaling

**Definition 12** *Translation shifts a function horizontally or vertically, while scaling changes its size.*


Translation and scaling are fundamental operations in function manipulation, allowing for adjustments in position and size of the graph of a function.

**Types of Transformations:**

- *Horizontal Translation:* Shifting a function left or right. For instance,  $f(x)$  to  $f(x - h)$  shifts the function right by  $h$ .
- *Vertical Translation:* Moving a function up or down. For example,  $f(x)$  to  $f(x) + k$  raises the function by  $k$ .
- *Horizontal Scaling:* Stretching or compressing a function horizontally.  $f(x)$  to  $f(cx)$  scales the function horizontally.
- *Vertical Scaling:* Altering the function's height.  $f(x)$  to  $cf(x)$  scales the function vertically.


**Graphing Translated and Scaled Functions:** Let's consider the function  $f(x) = x^2$ , its translation, and scaling.

Original Function  $f(x) = x^2$






Translated Function  $f(x - 2) = (x - 2)^2$



Scaled Function  $2f(x) = 2x^2$



**Applications of Translation and Scaling:** These transformations are widely used in data analysis for adjusting graphs, in physics to model motion and waves, and in other mathematical contexts for function analysis.

**Exercise** Graph the function  $g(x) = (x - 3)^2 + 2$  and describe its translation and scaling relative to  $f(x) = x^2$ .

**Exercise** For the function  $h(x) = \frac{1}{2} \sin(2x + \pi)$ , identify the translations and scalings applied to the basic sine function.

**Exercises on Translation and Scaling**

**Exercise** Graph the function  $g(x) = (x-3)^2 + 2$  and describe its translation and scaling relative to  $f(x) = x^2$ .

**Exercise** For the function  $h(x) = \frac{1}{2}\sin(2x + \pi)$ , identify the translations and scalings applied to the basic sine function.

**Exercise** Consider the function  $p(x) = -3\cos(x - \pi/2)$ . Describe its vertical and horizontal translations, and vertical scaling.

**Exercise** Determine the translation and scaling transformations applied to the function  $f(x) = x^2$  to obtain  $q(x) = 4(x+1)^2 - 5$ .

**Exercise** Sketch the graph of  $r(x) = \frac{1}{3}(x-2)^3 + 4$  and explain the transformations applied to the basic cubic function  $f(x) = x^3$ .

**Exercise** Graph the function  $s(x) = 2\sin(\pi x - \pi/2) + 1$  and describe its periodicity, translation, and scaling.

**Exercise** Analyze the function  $t(x) = \frac{x+2}{x-1}$  for vertical and horizontal translations and any scaling factors.

**Exercise** Identify the transformations applied to the exponential function  $f(x) = e^x$  to produce  $u(x) = 2e^{-x+3} - 1$ .

**Exercise** Consider the function  $v(x) = \sqrt{x+4} - 2$ . Determine the transformations applied to the basic square root function  $f(x) = \sqrt{x}$ .

**Exercise** Graph the function  $w(x) = \ln(x-3)+2$  and describe its translation and scaling relative to the natural logarithm function  $f(x) = \ln(x)$ .

**Solutions to Exercises on Translation and Scaling**

*Solution:* The function  $g(x) = (x-3)^2 + 2$  is a translation of  $f(x) = x^2$ . It is translated 3 units to the right and 2 units up.

*Solution:* For  $h(x) = \frac{1}{2}\sin(2x + \pi)$ , the function is scaled vertically by a factor of  $\frac{1}{2}$ , horizontally by a factor of  $\frac{1}{2}$  (period is  $\pi$ ), and translated  $\frac{\pi}{2}$  units to the left.

*Solution:* The function  $p(x) = -3\cos(x - \pi/2)$  is vertically scaled by a factor of 3, inverted (due to the negative sign), and translated  $\frac{\pi}{2}$  units to the right.

*Solution:* In  $q(x) = 4(x + 1)^2 - 5$ , the function  $f(x) = x^2$  is horizontally translated 1 unit to the left, vertically scaled by a factor of 4, and translated 5 units down.

*Solution:* The function  $r(x) = \frac{1}{3}(x - 2)^3 + 4$  is a transformation of  $f(x) = x^3$  with a horizontal translation 2 units to the right, vertical scaling by  $\frac{1}{3}$ , and vertical translation 4 units up.

*Solution:* For  $s(x) = 2\sin(\pi x - \pi/2) + 1$ , the function is vertically scaled by 2, translated  $\frac{1}{2}$  unit to the right, and translated 1 unit up. The period is  $\frac{2\pi}{\pi} = 2$ .

*Solution:* The function  $t(x) = \frac{x+2}{x-1}$  involves a horizontal translation 1 unit to the right and 2 units to the left in the numerator. There's no clear vertical or horizontal scaling.

*Solution:* The function  $u(x) = 2e^{-x+3} - 1$  is translated 3 units to the right, reflected over the y-axis, vertically scaled by 2, and translated 1 unit down from  $f(x) = e^x$ .

*Solution:* In  $v(x) = \sqrt{x+4} - 2$ , the square root function  $f(x) = \sqrt{x}$  is translated 4 units to the left and 2 units down.

*Solution:* The function  $w(x) = \ln(x - 3) + 2$  is a translation of the natural logarithm function  $f(x) = \ln(x)$ , shifted 3 units to the right and 2 units up.

## 9.5.2 Reflection and Rotation

**Definition 13** *Reflection inverts a function across a line, and rotation turns it around a point.*


Reflection and rotation are key transformations in function manipulation, changing the orientation of the graph of a function.

### Types of Transformations:


- *Reflection Across the y-axis:* Reflecting a function  $f(x)$  to  $f(-x)$  inverts it across the y-axis.
- *Reflection Across the x-axis:* Reflecting a function  $f(x)$  to  $-f(x)$  inverts it across the x-axis.
- *Rotation:* Rotation involves turning the graph of a function around a specific point, often the origin.

**Graphing Reflected and Rotated Functions:** Consider the function  $f(x) = x^2$  and its reflection across the y-axis.

Original Function  $f(x) = x^2$



Reflected Function  $f(-x) = (-x)^2$



**Applications of Reflection and Rotation:** These transformations are used in various applications, such as in physics for understanding wave properties, in engineering for signal processing, and in computer graphics for image manipulation.

**Exercise** Graph the function  $g(x) = -\frac{1}{x}$  and describe its reflection relative to the basic function  $f(x) = \frac{1}{x}$ .

**Exercise** Consider the function  $h(x) = \sqrt{-x}$ . Describe the transformation applied to the basic function  $f(x) = \sqrt{x}$ .

**Exercises on Reflection and Rotation**

**Exercise** Graph the function  $g(x) = -\frac{1}{x}$  and describe its reflection relative to the basic function  $f(x) = \frac{1}{x}$ .

**Exercise** Consider the function  $h(x) = \sqrt{-x}$ . Describe the transformation applied to the basic function  $f(x) = \sqrt{x}$ .

**Exercise** Reflect the function  $p(x) = x^3$  across the x-axis and sketch the resulting graph.

**Exercise** Graph the function  $q(x) = -\sin(x)$  and explain how it is transformed from the basic sine function.

**Exercise** Analyze the rotation of the function  $r(x) = \cos(-x)$  and describe how it differs from  $f(x) = \cos(x)$ .

**Exercise** Reflect the function  $s(x) = e^x$  across the y-axis and describe the changes in its graph.

**Exercise** Graph  $t(x) = -\ln(x)$  and discuss its reflection in relation to the function  $f(x) = \ln(x)$ .

**Exercise** Consider the function  $u(x) = \tan(-x)$ . Is it a reflection or a rotation of the basic tangent function? Explain.

**Exercise** Sketch the function  $v(x) = \sqrt{1-x}$  and describe the transformation from  $f(x) = \sqrt{x}$ .

**Exercise** Reflect the function  $w(x) = x^2 + x$  across the y-axis and sketch the resulting graph.

**Solutions to Exercises on Reflection and Rotation**

*Solution:* The function  $g(x) = -\frac{1}{x}$  is a reflection of  $f(x) = \frac{1}{x}$  across the x-axis. The negative sign in front of the function inverts it vertically.

*Solution:* The function  $h(x) = \sqrt{-x}$  is a reflection of  $f(x) = \sqrt{x}$  across the y-axis. The negative sign inside the square root reflects the function horizontally.

*Solution:* Reflecting  $p(x) = x^3$  across the x-axis results in  $-x^3$ . This inversion changes the direction of the cubic curve, flipping it vertically.

*Solution:* The function  $q(x) = -\sin(x)$  is the basic sine function reflected across the x-axis. The negative sign reverses the peaks and troughs of the sine wave.

*Solution:* The function  $r(x) = \cos(-x)$  is the same as  $f(x) = \cos(x)$  due to the even property of the cosine function. It represents a rotation of the cosine function by  $\pi$  radians, but the graph remains unchanged.

*Solution:* Reflecting  $s(x) = e^x$  across the y-axis results in  $e^{-x}$ . The graph changes from increasing exponentially to decreasing exponentially as  $x$  increases.

*Solution:* Graphing  $t(x) = -\ln(x)$  shows it is a reflection of  $f(x) = \ln(x)$  across the x-axis. The logarithmic curve is inverted vertically.

*Solution:* The function  $u(x) = \tan(-x)$  is a reflection of the basic tangent function across the y-axis due to the odd property of the tangent function.

*Solution:* The function  $v(x) = \sqrt{1-x}$  is transformed from  $f(x) = \sqrt{x}$  by a horizontal reflection across the y-axis and a horizontal translation 1 unit to the right.

*Solution:* Reflecting  $w(x) = x^2 + x$  across the y-axis results in  $(-x)^2 - x = x^2 - x$ . The quadratic curve is flipped horizontally.

# Chapter 10

## Limits and Continuity - Detail

### 10.1 Introduction to Limits - Detail

#### 10.1.1 Definition of a Limit - Detail

**Definition 14** *The limit of  $f(x)$  as  $x$  approaches  $a$  is  $L$  if for every  $\epsilon > 0$ , there exists a  $\delta > 0$  such that  $|f(x) - L| < \epsilon$  whenever  $0 < |x - a| < \delta$ .*

This definition, known as the  $\epsilon - \delta$  definition of a limit, is fundamental in calculus. It formalizes the concept of approaching a value.

#### Understanding $\epsilon - \delta$ Criteria:

- $\epsilon$  (epsilon) represents how close  $f(x)$  is to  $L$ .
- $\delta$  (delta) represents how close  $x$  is to  $a$ .
- The statement means that  $f(x)$  gets arbitrarily close to  $L$  as  $x$  approaches  $a$ .

**Applications of Limits:** Limits are used in calculus to define derivatives, integrals, and continuity. They are essential in understanding instantaneous rates of change and areas under curves.

**Exercise** Using the  $\epsilon - \delta$  definition, show that the limit of  $f(x) = x^2$  as  $x$  approaches 2 is 4.

**Exercise** Consider  $f(x) = \frac{1}{x}$ . Explain using the  $\epsilon - \delta$  definition why the limit as  $x$  approaches 0 does not exist.

**Exercises on the Definition of a Limit**

**Exercise** Using the  $\epsilon - \delta$  definition, show that the limit of  $f(x) = x^2$  as  $x$  approaches 2 is 4.

**Exercise** Consider  $f(x) = \frac{1}{x}$ . Explain using the  $\epsilon - \delta$  definition why the limit as  $x$  approaches 0 does not exist.

**Exercise** Prove that the limit of  $g(x) = 3x + 2$  as  $x$  approaches 4 is 14 using the  $\epsilon - \delta$  approach.

**Exercise** Use the  $\epsilon - \delta$  definition to demonstrate that the limit of  $h(x) = \sqrt{x}$  as  $x$  approaches 9 is 3.

**Exercise** Verify that the limit of  $p(x) = \frac{x^2-1}{x-1}$  as  $x$  approaches 1 is 2 using the  $\epsilon - \delta$  criteria.

**Exercise** Determine whether the limit of  $q(x) = \frac{\sin(x)}{x}$  as  $x$  approaches 0 exists, and if so, find the limit using the  $\epsilon - \delta$  definition.

**Exercise** Prove that the limit of  $r(x) = \frac{1}{x^2}$  as  $x$  approaches -3 is  $\frac{1}{9}$ , employing the  $\epsilon - \delta$  method.

**Exercise** Using the  $\epsilon - \delta$  definition, establish that the limit of  $s(x) = \frac{x-2}{x^2-4}$  as  $x$  approaches 2 does not exist.

**Exercise** Determine the limit of  $t(x) = x^3$  as  $x$  approaches -1 using the  $\epsilon - \delta$  definition, and verify your result.

**Exercise** Use the  $\epsilon - \delta$  approach to show that the limit of  $u(x) = \frac{1}{\sqrt{x+4}}$  as  $x$  approaches 0 is  $\frac{1}{2}$ .

**Solutions to Exercises on the Definition of a Limit**

*Solution:* To show the limit of  $f(x) = x^2$  as  $x$  approaches 2 is 4, choose  $\delta = \min\left(1, \frac{\epsilon}{5}\right)$ . For  $0 < |x - 2| < \delta$ , we have  $|f(x) - 4| = |x^2 - 4| = |(x - 2)(x + 2)| < 5|x - 2| < \epsilon$ .

*Solution:* For  $f(x) = \frac{1}{x}$ , as  $x$  approaches 0, the function values increase without bound. Therefore, no matter how small  $\delta$  is chosen,  $|f(x) - L|$  cannot be made less than any  $\epsilon > 0$ . Hence, the limit does not exist.



*Solution:* To show the limit of  $g(x) = 3x + 2$  as  $x$  approaches 4 is 14, choose  $\delta = \frac{\epsilon}{3}$ . Then for  $0 < |x - 4| < \delta$ , we have  $|g(x) - 14| = |3x + 2 - 14| = 3|x - 4| < \epsilon$ .

*Solution:* For  $h(x) = \sqrt{x}$  and  $x$  approaching 9, choose  $\delta = \min(1, \epsilon^2)$ . Then  $0 < |x - 9| < \delta$  implies  $|\sqrt{x} - 3| < \epsilon$ .

*Solution:* For  $p(x) = \frac{x^2-1}{x-1}$ , factorize the numerator as  $(x-1)(x+1)$ . The limit as  $x$  approaches 1 is 2, as  $p(x)$  simplifies to  $x+1$  and  $p(1) = 2$ . To show this, choose  $\delta = \epsilon$ .

*Solution:* For  $q(x) = \frac{\sin(x)}{x}$  as  $x$  approaches 0, use the squeeze theorem. Since  $-1 \leq \sin(x) \leq 1$ , then  $-\frac{1}{x} \leq \frac{\sin(x)}{x} \leq \frac{1}{x}$ . As  $x$  approaches 0, both bounds approach 0, so the limit is 0.

*Solution:* To show the limit of  $r(x) = \frac{1}{x^2}$  as  $x$  approaches -3 is  $\frac{1}{9}$ , choose  $\delta = \min(1, 3\epsilon)$ . Then  $0 < |x + 3| < \delta$  implies  $|r(x) - \frac{1}{9}| < \epsilon$ .

*Solution:* For  $s(x) = \frac{x-2}{x^2-4}$ , the function is undefined at  $x = 2$ , and the denominator approaches 0 as  $x$  approaches 2. Thus, the function values become unbounded, and the limit does not exist.

*Solution:* The limit of  $t(x) = x^3$  as  $x$  approaches -1 is -1. Choose  $\delta = \sqrt[3]{\epsilon}$ . Then  $0 < |x + 1| < \delta$  implies  $|x^3 + 1| < \epsilon$ .

*Solution:* To show the limit of  $u(x) = \frac{1}{\sqrt{x+4}}$  as  $x$  approaches 0 is  $\frac{1}{2}$ , choose  $\delta = \min(1, 4\epsilon^2)$ . Then  $0 < |x| < \delta$  implies  $|u(x) - \frac{1}{2}| < \epsilon$ .

### 10.1.2 One-sided Limits

**Definition 15** *The one-sided limits of  $f(x)$  as  $x$  approaches  $a$  from the left (or right) are the values the function approaches as  $x$  approaches  $a$  from the left (or right).*

### 10.1.3 Limits Involving Infinity

**Definition 16** *A limit involving infinity is the value that  $f(x)$  approaches as  $x$  approaches infinity, or as  $f(x)$  approaches infinity for some finite  $x$ .*

## 10.2 Properties of Limits

### 10.2.1 Limit Laws

**Theorem 1** *Standard limit laws include the sum law, product law, quotient law, and power law.*

### 10.2.2 Squeeze Theorem

**Theorem 2** *If  $f(x) \leq g(x) \leq h(x)$  for all  $x$  near  $a$  and the limits of  $f(x)$  and  $h(x)$  as  $x$  approaches  $a$  are equal, then the limit of  $g(x)$  as  $x$  approaches  $a$  is the same.*

### 10.2.3 Limits of Trigonometric Functions

**Theorem 3** *Includes limits such as  $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$ .*

## 10.3 Continuity - Detail

### 10.3.1 Definition and Examples

**Definition 17** *A function is continuous at a point if it is defined at that point, its limit exists at that point, and the limit equals the function value.*

### 10.3.2 Types of Discontinuities

**Definition 18** *Discontinuities can be classified as removable, jump, or infinite.*

### 10.3.3 Intermediate Value Theorem

**Theorem 4** *If a function is continuous on a closed interval  $[a, b]$  and  $N$  is any number between  $f(a)$  and  $f(b)$ , then there exists a number  $c$  in the interval such that  $f(c) = N$ .*

# Chapter 11

## The Derivative - Detail

### 11.1 Understanding Derivatives

#### 11.1.1 Definition and Notation

**Definition 19** *The derivative of a function  $f$  at a point  $a$  is defined as  $\lim_{h \rightarrow 0} \frac{f(a+h)-f(a)}{h}$ , if this limit exists.*

#### 11.1.2 Interpreting Derivatives

**Discussion 1** *The derivative at a point gives the slope of the tangent line to the function at that point. It represents the rate of change of the function with respect to its variable.*

#### 11.1.3 Derivative as a Function

**Definition 20** *The derivative of a function  $f$  is itself a function that assigns to each point  $x$  the slope of the tangent to  $f$  at  $x$ .*

### 11.2 Techniques of Differentiation

#### 11.2.1 Product Rule and Quotient Rule

**Theorem 5** *The product rule states that  $(fg)' = f'g + fg'$  and the quotient rule states that  $\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$ , where  $f$  and  $g$  are functions of  $x$ .*

### 11.2.2 Chain Rule

**Theorem 6** *If a function  $y$  is a function of  $u$  which is a function of  $x$ , then the derivative of  $y$  with respect to  $x$  is given by  $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$ .*

### 11.2.3 Higher-Order Derivatives

**Definition 21** *The  $n$ -th order derivative of  $f$ , denoted  $f^{(n)}$ , is the derivative of  $f^{(n-1)}$ .*

### 11.2.4 Implicit Differentiation

**Method 1** *Implicit differentiation is used when a function is given in an implicit form  $F(x, y) = 0$  rather than the explicit form  $y = f(x)$ .*

## 11.3 Applications of Derivatives

### 11.3.1 Tangent Lines and Normal Lines

**Application 1** *The equation of the tangent line to a curve  $y = f(x)$  at a point  $(a, f(a))$  is  $y - f(a) = f'(a)(x - a)$ .*

### 11.3.2 Increasing and Decreasing Functions

**Theorem 7** *A function  $f$  is increasing (decreasing) on an interval if its derivative  $f'$  is positive (negative) on that interval.*

### 11.3.3 Maxima and Minima

**Method 2** *Critical points, where  $f'(x) = 0$  or  $f'(x)$  does not exist, are potential locations of local maxima or minima.*

### 11.3.4 Concavity and Inflection Points

**Discussion 2** *A function is concave up (down) where its second derivative  $f''$  is positive (negative). Points where  $f''$  changes sign are inflection points.*

### 11.3.5 Optimization Problems

**Application 2** *Optimization involves finding the maximum or minimum values of a function subject to certain constraints, often using derivatives.*

### 11.3.6 Related Rates

**Method 3** *Related rates problems involve finding the rate at which one quantity changes with respect to another using the chain rule.*



# Chapter 12

## Integration

### 12.1 Antiderivatives

#### 12.1.1 Definition and Basic Techniques

**Definition 22** *An antiderivative of a function  $f(x)$  is a function  $F(x)$  such that  $F'(x) = f(x)$ .*

**Example 7:** An antiderivative of  $f(x) = 2x$  is  $F(x) = x^2 + C$ , where  $C$  is a constant.

#### 12.1.2 Indefinite Integrals

**Definition 23** *The indefinite integral of  $f(x)$ , denoted  $\int f(x) dx$ , represents the collection of all its antiderivatives.*

#### 12.1.3 Initial Value Problems

**Problem 1:** Given a differential equation  $f'(x) = g(x)$  and an initial condition  $f(x_0) = y_0$ , find the function  $f(x)$ .

### 12.2 Definite Integrals

#### 12.2.1 Definition and Interpretation

**Definition 24** *The definite integral of  $f(x)$  from  $a$  to  $b$ , denoted  $\int_a^b f(x) dx$ , is the limit of the sum of areas of rectangles under the curve of  $f(x)$  as the width of the rectangles approaches zero.*

### 12.2.2 Properties of Definite Integrals

**Theorem 8** *Properties include linearity, additivity, and the fact that the integral from  $a$  to  $a$  is zero.*

### 12.2.3 Fundamental Theorem of Calculus

**Theorem 9** *If  $F$  is an antiderivative of  $f$  on an interval, then  $\int_a^b f(x) dx = F(b) - F(a)$ .*

### 12.2.4 Numerical Integration Methods

**Method 4** *Methods include the Trapezoidal Rule and Simpson's Rule, which provide approximations of definite integrals.*

## 12.3 Applications of Integrals

### 12.3.1 Area Between Curves

**Application 3** *The area between curves  $f(x)$  and  $g(x)$  from  $a$  to  $b$  is  $\int_a^b |f(x) - g(x)| dx$ .*

### 12.3.2 Volumes of Solids of Revolution

**Application 4** *The volume of a solid obtained by rotating a region about an axis can be found using the Disk or Washer methods.*

### 12.3.3 Arc Length and Surface Area

**Method 5** *Formulas for the arc length of a curve and the surface area of a solid of revolution involve definite integrals.*

### 12.3.4 Work and Fluid Forces

**Application 5** *Work done in moving an object and the force exerted by a fluid can be calculated using integration.*



# Chapter 13

## Advanced Topics in Calculus

### 13.1 Sequences and Series

#### 13.1.1 Convergence and Divergence

**Definition 25** *A sequence  $\{a_n\}$  converges if it approaches a limit as  $n$  goes to infinity. A series  $\sum a_n$  converges if the sequence of its partial sums converges.*

#### 13.1.2 Tests for Convergence

**Method 6** *Tests include the Integral Test, Comparison Test, Ratio Test, and Root Test.*

#### 13.1.3 Power Series and Taylor Series

**Definition 26** *A power series is a series of the form  $\sum_{n=0}^{\infty} a_n(x - c)^n$ . A Taylor series is a power series that represents a function.*

### 13.2 Multivariable Calculus

#### 13.2.1 Functions of Several Variables

**Definition 27** *A function of several variables is a function that takes several inputs and produces a single output.*

### 13.2.2 Partial Derivatives

**Definition 28** *The partial derivative of a function with respect to one of its variables is its derivative with respect to that variable, holding the other variables constant.*

### 13.2.3 Multiple Integrals

**Definition 29** *Multiple integrals involve integration over more than one variable, such as double and triple integrals.*

### 13.2.4 Vector Calculus

**Discussion 3** *Vector calculus involves differentiation and integration of vector fields, including topics like gradient, divergence, and curl.*

## 13.3 Differential Equations

### 13.3.1 First Order Differential Equations

**Method 7** *Methods for solving include separation of variables, integrating factors, and exact equations.*

### 13.3.2 Second Order Linear Differential Equations

**Method 8** *Includes homogeneous and non-homogeneous equations, characteristic equations, and particular solutions.*

### 13.3.3 Systems of Differential Equations

**Discussion 4** *Systems of differential equations involve several interrelated differential equations and can be solved using matrix methods and eigenvalues.*

## 13.4 Conclusion

This concludes the math currently included in CiB. Please fork the LaTeX source code for CiB (available on GitHub) and create your own book that chooses the facts and exercises most relevant to you! Also, starring the CiB project on GitHub would be greatly appreciated! Thanks for reading CiB!

# Appendix I

## Basic GitHub Guide

### A Quick Start to Your GitHub Journey

Welcome to the fascinating world of GitHub, a platform that has revolutionized the way we collaborate on projects, share code, and build software together. Whether you are a programmer, a writer, or a mathematician, GitHub provides a set of powerful tools to help you collaborate with others, manage your projects, and contribute to the vast world of open-source software. In this guide, we will walk you through the foundational steps to get started with GitHub, helping you to navigate, contribute, and make the most out of this incredible platform.

### Creating Your GitHub Account

The first step to joining the GitHub community is to create an account. Here's how you can do it:

1. Visit the [GitHub website](#).
2. Click on the “Sign up” button.
3. Fill in the required information, including your username, email address, and password.
4. Verify your account and complete the sign-up process.

Once you have created your account, take a moment to explore your new GitHub dashboard. Here, you will find a variety of tools and features that will help you manage your projects, collaborate with others, and discover new and interesting repositories.

## Creating Your First Repository

A repository (or “repo”) is a digital directory where you can store your project files. Here’s how you can create your first repository:

1. From your GitHub dashboard, click on the “New” button to create a new repository.
2. Give your repository a name and provide a brief description.
3. Initialize this repository with a README file. (This is an optional step, but it’s a good practice to include a README file in every repository to explain what your project is about.)
4. Click “Create repository.”

Congratulations! You have just created your first GitHub repository. You can now start adding files, collaborating with others, and managing your project right from GitHub.

## Making Changes and Commits

GitHub uses Git, a version control system, to keep track of changes made to your project. Here’s a quick guide on how to make changes and commits:

1. Navigate to your repository on GitHub.
2. Find the file you want to edit, and click on it.
3. Click the pencil icon to start editing.
4. Make your changes and then scroll down to the “Commit changes” section.
5. Provide a commit message that explains the changes you made.
6. Choose whether you want to commit directly to the main branch or create a new branch for your changes.
7. Click “Commit changes.”

Your changes are now saved, and a new commit is created. Every commit has a unique ID, making it easy to track changes, revert to previous versions, and collaborate with others.

## Collaborating with Others

One of the biggest strengths of GitHub is its collaborative nature. Here are some ways you can collaborate with others:

- **Forking:** You can fork a repository, create your own copy, make changes, and then propose those changes back to the original project.
- **Issues:** Use issues to report bugs, request new features, or start a discussion with the community.
- **Pull Requests:** Propose changes to a project by creating a pull request. This allows others to review your changes, discuss them, and eventually merge them into the project.

## Conclusion: Embarking on Your GitHub Adventure

Now that you have a basic understanding of GitHub and how it works, you are ready to embark on your GitHub adventure. Explore repositories, contribute to open-source projects, collaborate with others, and build amazing things together. Remember, the GitHub community is vast and supportive, and there is a wealth of knowledge and resources available to help you along the way. Happy coding!



# Appendix II

## Basic Python and Colab Guide

### Introduction to Python for Calculus

Python's versatility in mathematics, science, engineering, and data analysis stems from its simplicity and extensive library ecosystem. This guide will delve into Python packages essential for math and calculus, showcasing their utility in Google Colab notebooks.

### Setting Up Python and Google Colab

Google Colab is a free, cloud-based platform enabling Python programming in a browser. It offers free computational resources, ideal for Python learning and experimentation.

Visit [Google Colab](#) to start. Create a new notebook, and use code cells to write and execute Python code with Shift+Enter.

### Important Python Packages for Calculus

#### NumPy

NumPy, fundamental for scientific computing, offers support for large, multi-dimensional arrays and matrices, along with various mathematical functions.

#### SymPy

SymPy, a library for symbolic mathematics, allows algebraic manipulations and equation solving symbolically, perfect for calculus operations like differentiation and integration.

## **Matplotlib**

Matplotlib, a Python plotting library, creates static, interactive, and animated visualizations, useful for graphing functions and data in calculus.

## **Pandas**

Pandas provide high-performance, easy-to-use data structures, and data analysis tools, invaluable for handling and analyzing mathematical data.

## **SciPy**

SciPy extends NumPy by adding a collection of algorithms and high-level commands for data manipulation and visualization.

## **Examples and Applications**

### **Calculating Derivatives and Integrals with SymPy**

Illustrate using SymPy to compute derivatives and integrals of functions.

### **Data Visualization with Matplotlib and Pandas**

Demonstrate graphing functions and datasets, highlighting calculus concepts.

### **Solving Equations with SciPy**

Show how to solve equations that commonly appear in calculus.

### **Numerical Methods in Python**

Discuss Python's capabilities in numerical differentiation and integration, useful in calculus.

### **Using Python for Limits and Continuity**

Examples showcasing how Python can help in understanding limits and continuity in functions.

## **Interactive Learning with Google Colab**

Highlight the benefits of using Colab notebooks for interactive calculus learning, including real-time collaboration and easy sharing.



## Creating a Colab Notebook for Practice Problems

In this section, we will guide you through creating a Google Colab notebook to solve calculus practice problems using Python.

### Setting Up Your Colab Notebook

To start solving calculus problems with Python:

1. Open [Google Colab](#).
2. Choose 'New Notebook' to create a blank notebook.
3. Rename the notebook to something descriptive, like 'Calculus Practice'.

### Installing Necessary Libraries

At the beginning of your notebook, install any necessary Python libraries. For these exercises, ensure NumPy, SymPy, and Matplotlib are available:

Remove # if the following packages are not installed:

```
# !pip install numpy sympy matplotlib
```

### Solving Exercise 1: Graphing a Linear Function

Let's solve the first exercise, which involves graphing a linear function.

```
import numpy as np
import matplotlib.pyplot as plt

# Define the function
def f(x):
    return 3*x - 2

# Generate x values
x = np.linspace(-10, 10, 400)

# Plot the function
plt.plot(x, f(x))
plt.xlabel('x')
plt.ylabel('f(x)')
plt.title('Graph of f(x) = 3x - 2')
plt.grid(True)
```

```
plt.show()

# Slope and y-intercept
print("Slope: 3")
print("Y-intercept: -2")
```

### Solving Exercise 2: Identifying Undefined Points in a Function

Now, let's address the second exercise, which requires identifying for what values the function  $g(x) = \frac{1}{x}$  is undefined.

```
import sympy as sp

x = sp.symbols('x')
g = 1/x

# Display the function
sp.plot(g, (x, -10, 10), title='Graph of g(x) = 1/x',
        xlabel='x', ylabel='g(x)', ylim=(-10, 10))

# Identifying undefined points
print("g(x) is undefined for x = 0")
```

### Accessing the Completed Colab Notebook

The Colab notebook we've discussed is available for viewing and interaction. You can access it by clicking on the following link: [Finished Colab Notebook on Graphing and Analysis](https://colab.research.google.com/drive/1HF-cmwqfIZ803i1i-CFyR7ssWsDt7WvV). This link will take you directly to the notebook hosted on Google Colab, where you can view the complete code and run it yourself.

<https://colab.research.google.com/drive/1HF-cmwqfIZ803i1i-CFyR7ssWsDt7WvV>

### Adding the Notebook to Your GitHub Repository

If you have downloaded the Colab notebook to your local machine and want to add it to your Git repository, follow these terminal commands on your Ubuntu machine:

```
# Navigate to your local Git repository directory
cd path/to/your/repo
```

```
# Add the Colab notebook file to the repository
git add name_of_the_notebook.ipynb

# Commit the addition with a descriptive message
git commit -m "Add Colab notebook for calculus exercises"

# Push the changes to the remote GitHub repository
git push origin main
```

### Using Colab Notebooks for Problem Solving

These examples demonstrate how you can use Google Colab and Python to solve and visualize calculus problems. You can use similar steps to tackle other exercises, explore different functions, and deepen your understanding of calculus concepts.

### Conclusion: Interactive Learning with Colab Notebooks

Google Colab notebooks offer an interactive and accessible way to explore calculus using Python. By integrating theoretical concepts with computational examples, students can gain a deeper understanding of calculus. We encourage you to use these notebooks to solve exercises, visualize mathematical concepts, and explore the vast possibilities that Python and Colab offer.

### Conclusion: Python and Colab in Calculus

Python, with its comprehensive libraries, offers a powerful toolset for calculus exploration. Combined with Google Colab, it provides an accessible, interactive platform for learning and experimentation. Embrace Python and Colab to enhance your understanding of calculus and to explore mathematical problems creatively and efficiently.



# Appendix III

## Basic L<sup>A</sup>T<sub>E</sub>X Guide

### A Quick Start to Your L<sup>A</sup>T<sub>E</sub>X Journey

Welcome to the immersive world of L<sup>A</sup>T<sub>E</sub>X, a typesetting system widely used for creating scientific and professional documents due to its powerful handling of formulas and bibliographies. This guide is designed to offer you the foundational steps to grasp the basics of L<sup>A</sup>T<sub>E</sub>X, enabling you to craft documents of high typographic quality akin to this book.

### Setting Up Your L<sup>A</sup>T<sub>E</sub>X Environment

Before you can start creating documents with L<sup>A</sup>T<sub>E</sub>X, you need to set up a working L<sup>A</sup>T<sub>E</sub>X environment on your computer. Here's how you can do it:

1. Download and install a T<sub>E</sub>X distribution, which includes L<sup>A</sup>T<sub>E</sub>X. For Windows, MiKTeX is a popular choice, while Mac users might prefer MacTeX, and TeX Live is widely used on Linux.
2. Install a L<sup>A</sup>T<sub>E</sub>X editor. Some popular options include TeXShop (for Mac), TeXworks (cross-platform), and Overleaf (an online L<sup>A</sup>T<sub>E</sub>X editor).
3. Ensure that your T<sub>E</sub>X distribution and L<sup>A</sup>T<sub>E</sub>X editor are properly configured and integrated.

### Creating Your First L<sup>A</sup>T<sub>E</sub>X Document

Once your L<sup>A</sup>T<sub>E</sub>X environment is set up, you are ready to create your first L<sup>A</sup>T<sub>E</sub>X document. Follow these steps:

1. Open your L<sup>A</sup>T<sub>E</sub>X editor and create a new document.
2. Insert the following code to set up a basic L<sup>A</sup>T<sub>E</sub>X document:

```
\documentclass{article}
\begin{document}
Hello, \LaTeX\ world!
\end{document}
```

3. Save your document with a .tex file extension.
4. Compile your document using your L<sup>A</sup>T<sub>E</sub>X editor. This process converts your .tex file into a PDF document.
5. View the output PDF and admire your first L<sup>A</sup>T<sub>E</sub>X creation.

## Understanding L<sup>A</sup>T<sub>E</sub>X Commands and Environments

L<sup>A</sup>T<sub>E</sub>X documents are created using a series of commands and environments. Commands typically start with a backslash `\` and are used to format text, insert special characters, or execute functions. Environments are used to define specific sections of your document that require special formatting.

- **Commands:** For example, `\{italics}` will render the word "italics" in italic font.
- **Environments:** To create a bulleted list, you would use the *itemize* environment:

```
\begin{itemize}
  \item First item
  \item Second item
\end{itemize}
```

## Adding Structure to Your Document

L<sup>A</sup>T<sub>E</sub>X makes it easy to structure your documents with sections, subsections, and chapters. Here's how you can add structure:

```
\section{Introduction}
This is the introduction of your document.
\subsection{Background}
This subsection provides background information.
\subsubsection{Details}
This is a subsubsection for more detailed information.
```

## Including Mathematical Formulas

L<sup>A</sup>T<sub>E</sub>X excels at typesetting mathematical formulas. Use the *equation* environment or the \$ sign for inline formulas. For example:

The quadratic formula is  $\left( x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \right)$ .

## Adding Images and Tables

You can also include images and tables in your L<sup>A</sup>T<sub>E</sub>X documents:

- **Images:** Use the *graphicx* package and the *includegraphics* command.
- **Tables:** Use the *tabular* environment to create tables.

## Compiling Your Document

L<sup>A</sup>T<sub>E</sub>X documents need to be compiled to produce a PDF. This can be done through your L<sup>A</sup>T<sub>E</sub>X editor. If your document includes bibliographies or cross-references, you may need to compile multiple times.

## Conclusion: Embracing the Power of L<sup>A</sup>T<sub>E</sub>X

Congratulations! You have taken your first steps into the world of L<sup>A</sup>T<sub>E</sub>X. With practice, you will discover that L<sup>A</sup>T<sub>E</sub>X is a powerful tool for creating professional-quality documents, from simple articles to complex books. Embrace the learning curve, explore the vast array of packages available, and join the community of L<sup>A</sup>T<sub>E</sub>X users who are ready to help you on your journey. Happy typesetting!





# Bibliography

- [MT77] F. Mosteller and J. W. Tukey. *Data Analysis and Regression: A Second Course in Statistics*. Addison-Wesley Pub Co, Reading, MA, 1977.