

Darada Property

WmD?[®]

Your medication is called: Amlodipine

It is to help:



High blood pressure



Angina (chest pain on exertion)


How to take, use, give:



Take with a drink



Tablets can be crushed if needed



Swallow whole

You may also: (tell a doctor or carer if any of these side effects worry you)




Feel sick, be sick or get tummy pain



Get a headache



Feel dizzy



Get a red face




Feel sleepy



Find it harder to get to sleep or stay asleep



Get swollen feet, hands or ankles



Get a swollen face



Get heart problems

You may also: (tell a doctor or carer straight away if you)



Get a heartbeat problem. You could feel your heart is beating slower



Get changes in mood



Feel anxious



Feel low or depressed, or feel you might want to harm yourself



Get an allergy. You could get a skin rash or itching



Get swollen lips or mouth (may be an allergy)



Get double vision

What you also need to know to keep you safe while on this medicine: (check with a health professional if you are worried).



It is not usually given if you have angina or have had a heart attack



Do not drink or eat grapefruit while taking this medicine



This medicine may affect your eyes. If it does, you **MUST** take extra care when driving, or not drive

What you need to know

The medicine may be given to you if you have a heart condition. It is important to tell your doctor if you have a heart condition or if you have had a heart attack. The medicine may be given to you if you have a heart condition or if you have had a heart attack. It is important to tell your doctor if you have a heart condition or if you have had a heart attack.

What you need to know

The medicine may be given to you if you have a heart condition. It is important to tell your doctor if you have a heart condition or if you have had a heart attack. The medicine may be given to you if you have a heart condition or if you have had a heart attack. It is important to tell your doctor if you have a heart condition or if you have had a heart attack.

Check the product information leaflet for a full list of side effects and how to use this medicine



Do not stop taking this medicine suddenly



Do not drive or operate machines if this medicine makes you feel faint or dizzy



Do not take this medicine if
you are pregnant or
breastfeeding