



WmD?[®]

It is to help:

A cartoon illustration of a young boy with brown hair, wearing a yellow shirt and a blue bow tie. He is sitting at a table, holding a fork in his right hand and a knife in his left hand, ready to eat. On the table in front of him is a plate with a piece of meat, some vegetables, and a glass of milk.

1 of 3



Feel less hungry



Feel faint, sweaty or dizzy
(your blood sugar levels may
be going down)



Feel weak or tired. You may
have anaemia


You may also: (tell a doctor or carer straight away if you)



Feel weak, sick, or have
breathing problems (could be
lactic acidosis)



Become deeply unconscious (a
Coma)



Get lung and breathing
problems

What you also need to know to keep you safe while on this
medicine: (check with a health professional if you are worried).



If you have diabetes, the
closer you stick to your diet
the better this medicine works



Stop this medicine if you have
an iodine injection before an X-
Ray or scan



Do not drink alcohol while on
this medicine



Do not take this medicine if
you are pregnant or
breastfeeding



Stop taking this if you get diarrhoea or are sick (vomit)



This does not mix well with many other medicines. You must not take aspirin or other non-steroidal anti-inflammatory drugs (NSAIDs) such as ibuprofen. Always check with your health professional



Check the product information leaflet for a full list of side effects and how to use this medicine



You must get advice from your healthcare professional if you want to get pregnant or breastfeed