

Programme Evaluation of Maternal and Child Cash Transfer (MCCT) Programme in Kayah State, Myanmar

Draft Report

11 October 2019

Contents

Acknowledgements	7
Acronyms and abbreviations	8
1 Background	9
2 Study aims and objectives	9
2.1 Research Questions	9
2.2 Study Output	10
3 Methods	10
3.1 Study design	10
3.1.1 Survey of selected households, villages, and townships	10
3.1.2 Quasi-experimental cohort study	11
3.2 Data collection	13
3.2.1 Questionnaire design:	13
3.3 Training of enumerators	14
3.3.1 Anthropometric Training	14
3.3.2 Interviewer and Supervisor Training	15
3.3.3 Additional Supervisor Training	15
3.4 Survey management and supervision	15
3.5 Quality control	16
3.6 Data processing and analysis	17
3.6.1 Tools and methods	17
3.6.2 Indicator estimation	17
3.6.3 Regression discontinuity	26
4 Results	27
4.1 Population-representative survey	27
4.1.1 Poverty Probability Index	27
4.1.2 Food consumption score	28
4.1.3 Food consumption score - nutrition	30
4.1.4 Coping strategies index	35
4.1.5 Household food expenditure share	39
4.1.6 Water, sanitation and hygiene	42
4.1.7 Child health	48
4.1.8 Period prevalence of childhood illnesses	56
4.1.9 Treatment-seeking behaviour	58
4.1.10 Child nutrition	79
4.1.11 Maternal health	83

4.1.12	Antenatal care	88
4.1.13	Birth/delivery	97
4.1.14	Postnatal care	100
4.1.15	Newborn care	104
4.2	Maternal nutrition	110
4.2.1	Minimum dietary diversity for women	110
4.3	Cohort study	111
4.3.1	Child anthropometric indicators	111

Annex 1: Study questionnaires	115
--------------------------------------	------------

Annex 2: Anthropometric standardisation results	116
--	------------

Annex 3: Anthropometric data quality check	117
---	------------

List of Tables

2	FCS food groups and weights	19
3	FCS classification thresholds	20
4	Food groups classified by nutrients-of-interest for FCS-N	20
5	FCS-N categories based on nutrient-rich food groups consumption	20
6	Reduced CSI strategies/behaviours and their corresponding severity weights	21
7	Commonly used HFES thresholds	22
8	Myanmar EPI Schedule 2016	23
9	MUAC cut-offs for maternal acute undernutrition	25
10	Food consumption score	28
11	Food consumption score - nutrition	31
12	Consumption-based Coping Strategies Index	35
13	Livelihoods-based coping strategies index	37
14	Household food expenditure share	40
15	Water service ladders	43
16	Sanitation service ladders	45
17	Handwashing service ladders	46
18	Immunisation coverage	49
19	Immunisation coverage per vaccine type	51
20	Coverage of immunisation-associated services	55
21	Period prevalence of childhood illnesses	56
22	Treatment-seeking for diarrhoea	59
23	Where/who treatment is sought from	62
24	Payment for treatment	64
25	Treatment-seeking for acute respiratory infection	66
26	Where/who treatment is sought from	69

27	Payment for treatment	71
28	Treatment-seeking for fever	73
29	Where/who treatment is sought from	76
30	Payment for treatment	78
31	Child stunting/stuntedness	79
32	Child underweight	80
33	Child wasting by weight-for-height z-score	80
34	Child wasting by MUAC	80
35	Infant and young child feeding	81
36	Months to wait until next pregnancy	84
37	Use of contraceptive methods	85
38	Where contraception services received	86
39	Information on family planning/contraception	87
40	Antenatal care coverage	89
41	Antenatal care counselling	92
42	Antenatal care supplementation	94
43	Antenatal care testing	96
44	Gestational age at birth (weeks)	97
45	Location of birth/delivery	97
46	Reason for choosing birth/delivery location	98
47	Who assisted in birth/delivery	99
48	Birth/delivery method	100
49	Costs incurred for birth/delivery	100
50	Postnatal care coverage	101
51	Vitamin B1 supplementation	102
52	Costs of postnatal care	103
53	Newborn care coverage	105
54	Costs of newborn care	106
55	Fed colostrum to newborn	107
56	Knowledge of danger signs	108
57	Care of cord and delivery wound	109
58	Minimum dietary diversity for women	110

List of Figures

1	Poverty probability index by location type	27
2	Food consumption score classification by location type	29
3	Food consumption score classification by wealth quintiles	29
4	Vitamin A food consumption score classification by location type	32
5	Vitamin A food consumption score classification by wealth quintiles	33
6	Protein food consumption score classification by location type	34

7	Protein food consumption score classification by wealth quintiles	35
8	Consumption-based coping strategies index by location type	36
9	Consumption-based coping strategies index by wealth quintiles	36
10	Livelihoods-based coping strategies index by location type	37
11	Livelihoods-based coping strategies index by wealth quintiles	38
12	Household food expenditure share by location type	39
13	Household food expenditure share by wealth quintiles	40
14	Household food expenditure share by location type	41
15	Household food expenditure share by wealth quintiles	42
16	Water services ladder by location type	44
17	Water services ladder by wealth quintiles	44
18	Sanitation services ladder by location type	45
19	Water services ladder by wealth quintiles	46
20	Handwashing services ladder by location type	47
21	Handwashing services ladder by wealth quintiles	48
22	Immunisation coverage by location type	50
23	Immunisation coverage by wealth quintiles	50
24	Immunisation coverage per vaccine by location type	53
25	Immunisation coverage per vaccine by wealth quintiles	54
26	Immunisation-associated services coverage by location type	55
27	Immunisation-associated services coverage by wealth quintiles	56
28	Period prevalence of childhood illnesses by location type	57
29	Period prevalence of childhood illnesses by wealth quintiles	57
30	Treatment-seeking for diarrhoea by location type	60
31	Treatment-seeking for diarrhoea by wealth quintiles	60
32	Reasons for not seeking treatment by location type	61
33	Where/who treatment is sought from by location type	63
34	Costs incurred for treatment	65
35	Treatment-seeking for acute respiratory infection by location type	67
36	Treatment-seeking for acute respiratory infection by wealth quintiles	67
37	Reasons for not seeking treatment by location type	68
38	Where/who treatment is sought from by location type	70
39	Costs incurred for treatment by location type	72
40	Treatment-seeking for fever by location type	74
41	Treatment-seeking for fever by wealth quintiles	74
42	Reasons for not seeking treatment by location type	75
43	Where/who treatment is sought from by location type	77
44	Costs incurred for treatment in Kayah state	79
45	Infant and young child feeding by location type	82
46	Infant and young child feeding by wealth quintiles	83

47	Antenatal care coverage by location type	90
48	Antenatal care coverage by wealth quintiles	90
49	Weight-for-age z-score regression discontinuity	II1
50	Height-for-age z-score regression discontinuity	II2
51	Weight-for-height z-score regression discontinuity	II3
52	MUAC regression discontinuity	II4

Acknowledgements

Acronyms and abbreviations

Acronym	Definition
CPI	Community Partners International
CSI	Coping strategies index
CCSI	Consumption-based coping strategies index
DSW	Department of Social Welfare
FCS	Food consumption score
FCS-N	Food consumption score - nutrition
GAD	General Administration Department
GAM	Global acute malnutrition
HAZ	Height-for-age z-score
HFES	Household food expenditure share
IYCF	Infant and young child feeding
LCSI	Livelihoods-based coping strategies index
MAM	Global acute malnutrition
MCCT	Mother and Child Cash Transfer
MMK	Myanmar Kyatt
MSWRR	Ministry of Social Welfare, Relief and Resettlement
MS-NPAN	Multi-Sector National Plan of Action for Nutrition
MUAC	Middle upper arm circumference
NNC	National Nutrition Council
NSPSP	National Social Protection Strategic Plan
ODK	Open Data Kit
PPI	Poverty probability index
RCT	Randomised controlled trial
RD	Regression discontinuity
SAM	Severe acute malnutrition
SBCC	Social Behavioural Change Communication
TEM	Technical error of measurement
WASH	Water, sanitation and hygiene
WFP	World Food Programme

1 Background

Kayah State remains one of the less developed areas of Myanmar and is home to some of the most remote and isolated communities in the country with decades long armed conflicts between Ethnic Armed Organizations and Myanmar Tatmadaw. As confirmed by Myanmar Demographic and Health Survey conducted in 2015, children in Kayah State are more likely to be malnourished than the average child in Myanmar, with the prevalence of stunting being particularly high [[Ministry of Health and Sports - MoHS/Myanmar and ICF, 2017](#)]. Moreover, certain maternal and child health indicators are among the lowest in Myanmar, specifically concerning contraception prevalence, antenatal care visits as well as immunization rates amongst children 12 and 23 months of age.

Since 2017, the Ministry of Social Welfare, Relief and Resettlement (MSWRR) started rolling out a Maternal and Child Cash Transfer (MCCT) program in Chin, Rakhine, Kayah, and Kayin States through the Department of Social Welfare, in line with the National Social Protection Strategic Plan (NSPSP) and the Multi-Sector National Plan of Action for Nutrition (MS-NPAN). The program aims to improve nutritional outcomes for all mothers and children during the first 1000 days of life by ensuring that pregnant women and mothers have improved practices on nutrition, infant, and young child feeding, and health-seeking behaviors. The program aims to provide universal coverage to social behavior change communication (SBCC) and a maternal and child cash transfer (MCCT) of 15,000 MMK per month for all pregnant women and mothers with children under 2 years of age.

2 Study aims and objectives

To provide a basis for measuring and evaluating the outcomes and impact of the MCCT program over time by estimating current levels of indicators related to:

1. Nutrition based on anthropometric measurements of maternal and child weight, height, and mid-upper arm circumference (MUAC) in Kayah State;
2. Behaviors related to nutrition, infant and young child feeding (IYCF), and health-seeking among mothers in Kayah State; and,
3. Knowledge related to health, nutrition, and hygiene among pregnant women and mothers with under five children in Kayah state.

2.1 Research Questions

1. What are the current levels of indicators related to nutrition, infant and young child feeding (IYCF), and health-seeking behaviors of mothers and under 5 children in Kayah State?
2. What is the impact of the MCCT program on child stunting in Kayah State?
3. What is the impact of the MCCT program on maternal nutritional status in Kayah State?

2.2 Study Output

1. Estimates of the current levels of indicators related to nutrition, IYCF, and health-seeking behaviors among mothers and their children under 5 in Kayah State
2. Estimate of prevalence of childhood stunting in the study cohort of children exposed to the MCCT program and those who are not in Kayah State
3. Estimate of prevalence of maternal nutritional status in the study cohort of mothers exposed to the MCCT program and those who are not in Kayah State

3 Methods

3.1 Study design

The study included two primary components:

3.1.1 Survey of selected households, villages, and townships

This was composed of three surveys.

1. **Representative household survey** - The baseline survey was designed for the measurement and evaluation of outcomes over time by estimating the current levels of indicators related to nutrition, IYCF, and health-seeking behaviors. The baseline study was conducted during the roll-out of MCCT program in Kayah State.
2. **Village profiles** - Village level demographics, assets, and conditions were assessed in each of the selected villages in Kayah State.
3. **Township profiles** - Township profiles were developed from the township-level data collected from all townships in Kayah State from which sampled villages were from.

This component of the study was designed to address research question 1 (Section 2.1). The survey results were used to provide context to the results of the second study component (see Section 3.1.2).

3.1.1.1 Sampling frame

To assess the various outcome indicators for the assessment¹, a two-stage cluster sample survey was implemented. The **first stage sample** was a sample of about 75 villages out of the total villages in Kayah State stratified into 1) *urban*, 2) *rural* and 3) *hard-to-reach* areas. The **second stage sample**, also called the within-community sample, was a sample of households from the selected villages with mothers with children less than 5 years old. The second stage sample was selected using the list of household with children under 5 years of age from each village as identified by township General Administrative Unit (GAD) whenever possible.

A total of about 14 households with pregnant mothers and mothers with children less than 5 years were sampled in each village. If a small community was selected that was likely to have fewer than the required

¹Based on the earlier Chin assessment and based on feedback from various stakeholders.

number of eligible respondents, all eligible households and persons in that community are sampled by moving door-to-door. If a household had more than one mother with children less than 5 years of age, mother with children less than 2 years of age was prioritized for data collection. If there were more than one mother with children less than 2 years of age, one of these mothers was selected randomly. Only the under 5 children of the index mother were included in the anthropometric measurements if there were more than one mother with under 5 children in the same household. In addition, all pregnant mothers present in the household of the respondents were measured for middle upper arm circumference (MUAC).

3.1.1.2 Sample size estimation

Using childhood stunting as index indicator for sample size calculations, the sample size needed to detect a change of mean HAZ from 1.432 in Kayah State² to -0.932 with a calculated standard deviation 1.14406³ was calculated. This is an equivalent drop in stunting prevalence of about 10%. This equates to a sample size of 137 children less than 5 years old. Assuming a design effect for a cluster survey of 2, this sample size was inflated to 274. A sample size of 274 respondents was needed for each strata (*urban, rural and hard-to-reach*) in Kayah State giving a total of 822 respondents in total. Assuming a non-response rate of 20% overall, a total of 330 respondents was the target sample for each strata in Kayah State.

As described in the sampling frame above, a total of 25 villages per strata and 14 respondent mothers with children less than 5 years old were the target number of samples and a total of 350 respondents in each strata was the overall target in order to achieve at least 274. In Kayah state, a total of 1050 respondents was the overall target. This is anticipated to be more than enough sample size as per calculations above. For the selected respondents who were not present at the time of data collection, two revisits were conducted with the respondent classified as a non-responder after two unsuccessful attempts.

3.1.2 Quasi-experimental cohort study

As part of a longitudinal, quasi-experimental evaluation design, the baseline survey included a cohort of mothers and their children selected through specific study eligibility criteria based on eligibility and receipt of the services and benefits of the MCCT program. Once recruited, this cohort was assessed on appropriate indicators related to nutrition, IYCF and health-seeking behaviours at baseline. This same cohort will then be assessed at endline on the same set of appropriate and relevant indicators to detect any difference between those receiving benefits from the MCCT from those who are not based on a regression discontinuity analytical approach (see Section 3.6.3).

This component of the study provides a statistically robust comparison between those exposed to the MCCT program and those who are not hence detecting the possible impact that the programme has on a specific set of impact indicators.

Given the universal nature of the MCCT program and the need to create effective treatment and comparison groups to detect program impact, this study component used a similar regression discontinuity (RD) design as was developed and used for the MCCT evaluation in Chin State, but with distinct changes that

²Based on 2015 Myanmar Demographic and Health Survey [Ministry of Health and Sports - MoHS/Myanmar and ICF, 2017]

³Based on 2015 Myanmar Demographic and Health Survey [Ministry of Health and Sports - MoHS/Myanmar and ICF, 2017]

address limitations of the latter [Ministry of Social Welfare, Relief and Resettlement and Livelihoods and Food Security Trust Fund, 2018]. This design was made possible by the fact that there is a specific cut-off point for programme eligibility. Ultimately, the programme (treatment) effect can be detected as a discontinuity in the regression line around this cut-off which in this case is a specific date that determines eligibility. The design is *quasi-experimental*, since treatment and comparison groups were not selected at random but based on predefined characteristics.

3.1.2.1 Eligibility criteria

Eligibility criteria depended on three factors:

- Date of start of recruitment into the MCCT program (1st October 2018)
- Date when the recruitment for study participants will occur (6th August 2019)
- Receipt of MCCT benefits

The optimal bandwidth of ages of the children to include in the study needed to be as close to the cut-off date as possible such that the ages of the children in the control and intervention groups will be roughly similarly distributed. This meant up to one month before and after the recruitment date of 1st October 2018. Given this, the following eligibility criteria were applied:

Control group

Following are the criteria for eligibility to the control group of the study:

1. Women who gave birth on/after 1 September 2018 but before 1 Oct 2018 (i.e. 1 month before the MCCT registration cut-off)
2. Women who were pregnant/gave birth after 1 Oct 2018 but have not been recruited into program

Intervention group

The criteria for eligibility to the intervention group of the study was women who were pregnant on or after 1 Oct 2018 and have been recruited into the program.

3.1.2.2 Sample size and selection

The RD design effect was factored in for sample size calculations. This design effect was used to inflate standard randomized controlled trials (RCT) sample sizes in order to account for the RD design. The RD design effect was primarily influenced by the distribution of characteristics of the cut-off variable (i.e. running variable) [Bor et al., 2014]. Previous literature advise that if the running variable is normally distributed around the cut-off, then the design effect is approximately 2.75. If the running variable is uniformly distributed around the cut-off, the design effect is 4, and if it is bimodal in distribution, then the design effect can go up to 5 [Schochet, 2009].

Given that the running variable for this study is the start date of registration in relation to pregnancy, it was expected that there would be various peaks in number of pregnancies at specific time points following a

general seasonal pattern not necessarily related to the cut-off date. The distribution is therefore likely multi-modal. So an assumption of at least 5 design effect for this RD design was taken into account.

Using prevalence of stunting as the index indicator for calculating sample size and using mean change in height-for-age z-score (HAZ) as the specific variable to use for assessing change, the same sample size calculation as in the survey was arrived at which was 137 for controls and 137 for intervention (total of 274) for each state. However, given the very narrow bandwidth of inclusion into the study used (one month before and after the programme start), it is likely that the universe population from which to select the cohort is quite small and that the total sample size of 274 is likely requiring an exhaustive sampling of all children born within the specified periods above. In this case, the inflation factor to account for design effect was not considered to be relevant given that an exhaustive sample will be drawn to begin with. Hence the 274 sample size for Kayah State was set as the target sample size.

3.1.2.3 Sample recruitment

Sample recruitment utilised a full list or registry of births one month before and after the 1st of October 2018. As an exhaustive sample was needed, all efforts were undertaken to identify and recruit into the study including engagement with the DSW in listing out and finding these births across Kayah state.

3.2 Data collection

Data was collected using an electronic data entry system based on the **Open Data Kit (ODK)** standard that runs on the Android operating software (OS) platform for mobile devices. The study instrument (see Annex 1) was encoded into the electronic data entry system platform and was served out of a secure aggregate server hosted by ONA⁴. Each data collector was provided with a tablet running on Android OS that was configured with the ODK application that receives the electronic data form. All measurements and answers by respondents were then recorded on the tablets and transmitted to the remote server whenever there was a mobile and/or wireless internet signal. This data collection system was not reliant on internet connection as the data collection could be performed offline, with the completed e-questionnaires saved on the interviewers' tablets. An internet connection was needed only when finalized encoded forms were ready to submit to the remote server, and this submission was done at regular intervals and timed when there was available internet connection for the data collectors. The survey teams aimed to submit forms on a daily basis.

3.2.1 Questionnaire design:

Structured questionnaires for the household survey, village profiles, and township profiles were in English and translated into local languages. For the household survey, all questions and responses were translated into written Myanmar, Pwo Karen, and Sgaw Karen. The data collector had the option to switch from one language to another on the tablet. The household survey also included anthropometric measurements for weight, height and MUAC for all pregnant women, mothers who have recently given birth, and every child up to five years of age in selected households. A separate form was developed for the anthropometric measurements.

⁴See <https://ona.io>

The village profile questionnaire assessed various village level characteristics relevant to the study such as number of and distances to nearby markets, education, and health facilities; presence of community groups (e.g., Village Health Committees) and their functionality; presence of community volunteers (e.g., auxiliary midwives and community health workers); access and quality of water and sanitation facilities; and general agricultural practices.

The township profile questionnaire collected data on the township's population, geography, and governance, with summaries of key nutrition indicators.

All approved questionnaires were back-translated into English to ensure that appropriate translation was done with discrepancies between language versions discussed and resolved between the study team and translators. Prior to the start of data collection, the household survey questionnaire was pilot tested to understand how well the wording and content, context effects, and interface design of the instruments performed in an actual survey setting and were revised as necessary. Respondents for field testing were purposively sampled from the same population as the baseline assessment (e.g., mothers from the study area, village leaders). The recommended minimum of 32 respondents were recruited for field testing of the household questionnaire. Field testing was iterative with changes to the data collection instruments implemented immediately and the revised instrument was re-tested. Field testing involved feedback from the following three sources:

- Respondents: During field testing of the questionnaire, respondents were asked first to respond to the questions and then to re-interpret each question in their own words to learn how respondents understood the question and how they came up with their answers. The study team members probed for potential sources of ambiguity/confusion including word choice and translation, as well as the relevance and comprehensiveness of responses options to each question.
- Interviewers: The study team members provided feedback on the questionnaire content to confirm that questions and responses were clear in the context of data collection. Field testing also examined how well the e-questionnaire and tablet functioned in the field. Senior study team members collected data from enumerators for the purpose of field testing using a standardized debriefing questionnaire.
- Design and content experts: Members of the study team reviewed the data collected during the field testing phase to check whether the responses correspond to the meaning of individual questions and the overall goals of the study.

3.3 Training of enumerators

Different trainings were conducted based on the skill that was being trained on and the type of personnel that was being trained.

3.3.1 Anthropometric Training

Recruited anthropometrists received an intensive 5-day training from an expert anthropometric measurement trainer with supervision by National Nutrition Center (NNC) and nutrition advisor(s). The training covered an introduction to the purpose of the study, nutritional indicators, anthropometry, accuracy and validity of measurements, utilization and equipment of the measurement tools (e.g., MUAC tape, weighing

scales, measuring board), and electronic data entry. The training emphasized practice and standardization of the anthropometric measurements to be used during the study according to methods adapted from the [WHO's Multicentre Growth Reference Study \[de Onis et al., 2004\]](#) and [INTERGROWTH-21 Project \[Is-mail et al., 2013\]](#) anthropometric training protocols.

During the training, women of reproductive age and children under five years of age were invited to participate as subjects, and each subject was measured twice by the expert trainer serving as gold standard and twice by each trainee. To assess inter- and intra-rater precision, the technical error of measurement (TEM) was calculated, with an acceptable TEM for an individual trainee defined as no more than twice that of the expert [Ulijaszek and Kerr, 1999]. Accuracy was assessed by calculating mean differences between the gold standard trainer and the trainee. Only participants who meet the required standard for accuracy and precision were selected for anthropometric data collection during the baseline assessment. The training was conducted in Myanmar language.

3.3.2 Interviewer and Supervisor Training

All supervisors and enumerators received an intensive 5-day training from study team members that covered the purposes of the study, informed consent procedures, the questionnaire guides and responses, electronic data entry, referral protocols for severe acute malnutrition, roles and responsibilities of all field staff, data quality assurance procedures, problem-solving, and conflict-sensitive approaches to data collection. The performance of trainees were assessed post-training using a written assessment based on didactic lessons, as well as performance during practice sessions. Separate trainings were conducted for the supervisors and interviewers in Kayah State.

3.3.3 Additional Supervisor Training

All supervisors received an additional 3 days of training from study team members that covered screening, random sampling method to identify respondents, quality control in the field, quality control of data entry and uploading, team management, and logistics.

3.4 Survey management and supervision

During data collection, 1 field supervisor supervised 1 team each composed of 4 enumerators and 2 anthropometrists. Each team was expected to complete 14 household interviews and one village profile per day depending on feasibility. Community Partners International worked with the Central NNC and State/Region Nutrition Teams and nutrition experts in developing a referral guideline and standard operating procedures to ensure that any women or children who were measured during the study and suspected of having severe acute malnutrition (SAM) were immediately referred to an appropriate health provider. Survey teams were provided with contact information of the nearest health providers to ensure timely referral. Alerts to refer women and children were programmed into the e-questionnaire on the ODK platform, based on WHO criteria for SAM.

3.5 Quality control

Quality control procedures throughout the assessment design, implementation, and analysis included:

- Translation, back-translation, and pilot testing as described above;
- Development of standard operating procedures and training manuals. CPI developed and distributed guidelines on data collection procedures, including the questionnaire and anthropometry methods, to all members of the data collection team;
- Training of data collectors where supervisors, enumerators, and anthropometrists practiced administering the questionnaire and checking and correcting questionnaires for accuracy and completeness. The performance of anthropometrists and data collectors were assessed in terms of confidence, independence, and reliability during practice sessions during the trainings, and based on results of an evaluation at the end of the training period;
- Anthropometric devices were calibrated and supervisors re-assessed some of the anthropometric measurements at intervals. Maintenance and quality check of all the other devices were done regularly as per the manufacturer's instructions. Data collection was coordinated and continuously supervised by field supervisors and senior researchers.
- The data collection tool was designed in such a way that any potential errors related to data entry were minimized by programmatically adding data entry checks that raise appropriate prompts to the data collector based on the potential error detected. The data collector was not allowed by the system to continue to the next field until the error has been corrected. For the anthropometric measurements, a test for the feasibility of the measurements taken based on the child's age and sex was implemented. If the measurement was above or below a certain threshold of feasible values, the system prompted the data collector to check their recording of the measurement, and to perform the measurement again. This approach was expected to pick up at least 90% of possible errors due to data entry and measurement errors. To complement this, an algorithm was added to randomly ask the data collector to repeat the measurement, even if the result of the first measurement was plausible. This approach was expected to pick up at least 90% of possible errors due to data entry and measurement errors.
- The Data Manager reviewed the data in real-time as they were uploaded to the server. This was done using a purpose-built application called `myanmarMCCTchecks`⁵ that allows for routine data checks automatically. In most cases, the error was detected and corrected immediately due to the data quality assurance measures built into the ODK platform, but if necessary, the interviewer was requested to revisit the respondent. In this case, the corrected questionnaire was uploaded and the wrong one deleted, so that only correct, checked records were stored in the database.

⁵See <https://validmeasures.io/myanmarMCCTchecks>

3.6 Data processing and analysis

3.6.1 Tools and methods

Data handling, processing, analysis and reporting were done using **version 3.6.1** of the [R Language for Statistical Computing](#)⁶ [R Core Team, 2019]. The R package **myanmarMCCTdata** was developed to support the handling, processing and analysis of data collected for the study. The package includes functions to retrieve data from the [ONA](#) server database, to appropriately structure datasets, to check and clean data, to recode data to respective indicator sets, to estimate indicators and to perform appropriate comparative analysis specified by the analysis⁷ [Guevarra and Goza, 2019]. The package uses several other R packages that provide additional functions for data processing, analysis and reporting⁸.

The main motivation for using R was to achieve transparent and reproducible research between the study collaborators and with external reviewers and/or readers. Developing the **myanmarMCCTdata** R package is the first step towards this aim of transparency and reproducibility as any collaborator and/or reviewer can easily retrieve the same codebase for data handling, processing and analysis used by the primary researchers. The second step was writing reports (including this report) using R itself using a seamless integration between data analysis and reporting. This was achieved using R Markdown [Xie et al., 2018] for producing reports which included code chunks of how the data processed using **myanmarMCCTdata** functions were analysed to produce the results reported here. This report and the analysis herein, can be reviewed and reproduced by collaborators and reviewers by following instructions given [here](#).

3.6.2 Indicator estimation

Data was processed and recoded based on standard and established indicator definitions whenever available. For some indicators, modifications to standard definitions were made based on information requirements by stakeholders or study-specific definitions were established in order to achieve analysis requirements. Several recode functions were developed in R through the **myanmarMCCTdata** package for this purpose. The following indicator sets were assessed:

3.6.2.1 Poverty probability index (PPI)

The [Poverty Probability Index \(PPI\)](#) is a poverty measurement tool that is statistically-sound, yet simple to use. The answers to 10 country-specific questions about a household's characteristics and asset ownership are scored to compute the likelihood that the household is living below a certain poverty line or definition [Innovations for Poverty Action, 2019]. PPI is used to identify households who are most likely to be poor or vulnerable to poverty.

The Myanmar MCCT survey collected answers to the 10 Myanmar-specific questions on household characteristics and asset ownership responses which were used to determine the household's PPI score based on

⁶Can be downloaded from the [Comprehensive R Archive Network](#) where guidance on installing R is also available.

⁷See <https://validmeasures.org/myanmarMCCTdata> to learn more about the **myanmarMCCTdata** package and how data is handled and processed using this package.

⁸For a full list of other R packages that **myanmarMCCTdata** depends on, see <https://validmeasures.org/myanmarMCCTdata>

the Myanmar national poverty line. The household's PPI score was then matched to poverty line/definition-specific lookup tables to determine the household's poverty probability. The matching of the PPI score to lookup tables was facilitated using the PPI lookup tables for Myanmar from the `ppitables` package [Guevarra, 2019a] for R.

3.6.2.2 Water, sanitation and hygiene

Recoding for indicators for water, sanitation and hygiene (WASH) were based on standard indicator definitions provided by WHO and UNICEF [WHO/UNICEF Joint Monitoring Programme for Water Supply and Sanitation, 2018, World Health Organization, 2006].

The following drinking water access indicators were assessed:

- Access to improved drinking water sources (regardless of collection time⁹)
- Access to unimproved drinking water sources only (unprotected dug well and unprotected spring)
- Drinking water from surface water (directly from a river, dam, lake, pond, stream, canal/irrigation canal)
- Probably safe drinking water based on appropriate water treatment method applied to water

These drinking water access indicators have been assessed for each of the three seasons (summer, rainy and winter season) experienced in Kayin and Kayah states.

In addition to drinking water access, the following drinking water use and storage behaviour was also assessed:

- Use of a water container/pot that is clean, covered and with a cup and handle

The following indicators on access to sanitation facilities were assessed:

- Access to improved sanitation facilities that are not shared with other households¹⁰
- Access to improved sanitation facilities that are shared with other households¹¹
- Access to unimproved sanitation facilities
- Use of open defecation

The following indicators on access to handwashing facilities were assessed:

- Access to handwashing facility on premises with soap and water¹²
- Access to handwashing facilities on premises without soap and water¹³

⁹Time to collect water from reported water source was not collected in baseline survey. The access to improved drinking water sources indicator does not take time into account. This indicator can be reframed as at least limited access to improved drinking water sources taking into account the definitions for limited and basic drinking water service.

¹⁰This is termed as access to **basic** sanitation facilities as per current indicator definitions.

¹¹This is termed as access to **limited** sanitation facilities as per current indicator definitions.

¹²This is termed as access to **basic** handwashing facilities as per current indicator definitions.

¹³This is termed as access to **limited** handwashing facilities as per current indicator definitions.

- No handwashing facilities

In addition, handwashing practices and behaviours were also assessed.

3.6.2.3 Food consumption score (FCS)

The **Food Consumption Score (FCS)** is an index developed by the World Food Programme (WFP) in 1996 [Vulnerability Assessment and Mapping World Food Programme, 2008]. The **FCS** is a household level indicator that aggregates food group diversity and frequency over the past 7 days. These food groups are then weighted according to their relative nutritional value. This means that food groups that are nutritionally-dense such as animal products are given greater weight than those containing less nutritionally dense foods such as tubers. The weights are then added up to come up with a household score which are then used to classify households into either poor, borderline, or acceptable food consumption. The **FCS** is a measure of quantity of caloric intake.

A brief questionnaire was used to ask respondents about the frequency of their households' consumption of eight different food groups over the previous seven days. The eight food groups are: main staples; pulses; vegetables; fruit; meat/fish; milk; sugar; and, oil. The frequency of consumption of these different food groups consumed by a household during the 7 days before the survey was then used to calculate a weighted diet diversity score for each household using the food group weights (Table 2) by multiplying the reported 7-day frequencies with the food group weights.

Table 2: FCS food groups and weights

	Food items	Food groups	Weight
1	Maize , maize porridge, rice, sorghum, millet pasta, bread and other cereals Cassava, potatoes and sweet potatoes, other tubers, plantains	Main staples	2
2	Beans, peas, groundnuts and cashew nuts	Pulses	3
3	Vegetables, leaves	Vegetables	1
4	Fruits	Fruits	1
5	Beef, goat, poultry, pork, eggs and fish	Meat and fish	4
6	Milk yoghurt and other diary	Milk	4
7	Sugar and sugar products, honey	Sugar	0.5
8	Oils, fats and butter	Oil	0.5
9	Spices, tea, coffee, salt, fishpowder, small amounts of milk for tea.	Condiments	0

All the weighted consumption frequencies were then summed up per household to arrive at the FCS for the specific household. Then, Using the appropriate thresholds (Table 3), each household is classified accordingly based on their FCS.

Table 3: FCS classification thresholds

FCS	Profiles
0 - 21	Poor
21.5 - 35	Borderline
> 35	Acceptable

3.6.2.4 Food consumption score nutrition quality analysis

In addition to the standard FCS, WFP has devised a set of nutritional adequacy indicators that assess the nutrition quality of the household diet using the data provided by a slightly modified FCS questionnaire that disaggregates some of the food groups into sub-food groups known to contain vitamin A, protein and heme-iron. This is called the Food Consumption Score Nutrition Quality Analysis or FCS-N [World Food Programme, 2015]. The FCS-N focuses on three nutrients - vitamin A, iron and protein - to assess the nutrition adequacy of a household's diet. The FCS-N was calculated by first aggregating the individual food groups into nutrient-rich food groups shown in Table 4.

Table 4: Food groups classified by nutrients-of-interest for FCS-N

Nutrients	Food Groups
Vitamin A-rich foods	Dairy, organ meat, eggs, orange vegetables, green leafy vegetables and orange fruits
Protein-rich foods	Pulses, dairy, flesh meat, organ meat, fish and eggs
Heme iron-rich foods	Flesh meat, organ meat, fish

Households were then classified by their frequency of consumption of the three nutrients of interest.

Table 5: FCS-N categories based on nutrient-rich food groups consumption

Category	No. of days of consumption
Never	0 days
Sometimes	1-6 days
At least daily	7 or more days

3.6.2.5 Coping strategy index

The coping strategy index or CSI is a simple indicator that reveals how households manage or cope with shortfalls in food consumption. It is based on a list of possible behaviours that households may use as strategies for coping with conditions of having little or no food to eat [Maxwell and Caldwell, 2008]. The CSI combines the frequency by which a behaviour is utilised and the severity of each strategy used. There are

two variations of the CSI. One assesses food consumption behaviours or strategies employed by the household in response to food insecurity (consumption-based CSI) while the other looks into the longer term, livelihoods-based strategies employed by households (livelihoods-based CSI).

The consumption-based coping strategy index (CCSI) used in this study was based on the reduced CSI questionnaire in order to allow for comparability between results for states and over time. The reduced CSI is based on 5 behaviours and corresponding severity weights for each as shown below.

Table 6: Reduced CSI strategies/behaviours and their corresponding severity weights

Behaviours/Strategies	Severity Weights
Rely on less preferred and less expensive food	1
Borrow food or rely on help from a relative or friend	2
Limit portion size at mealtimes	1
Restrict consumption by adults in order for small children to eat	3
Reduce number of meals eaten in a day	1

The CCSI was calculated by multiplying the reported frequency of utilising a coping behaviour by the corresponding severity weight shown in Table 6 and then taking the aggregate of these weighted scores. The CCSI is a continuous variable and the mean CCSI is used to describe the level of utilisation of coping strategies by households.

The livelihoods based coping strategies index or LCSi was used to better understand a household's longer term coping capacity to food insecurity. The respondent was asked to recall whether in the past 30 days their household has engaged in any of a set of 10 coping behaviours/strategies due to a lack of food or lack of money to buy food. These behaviours/categories were categorised into *stress* strategies, *crisis* strategies and *emergency* strategies. Based on the respondent's responses, their household was classified into the most severe strategy reported.

3.6.2.6 Household food expenditure share

Taking household expenditure as a proxy of income, the proportion of this expenditure spent on food is an indicator of household food security. A household that is more food insecure spends a bigger proportion of income on food. This indicator was calculated from the various monetary values of household spending grouped into food and non-food items. The share of household expenditure on food was estimated as follows:

$$\text{Household food expenditure share} = \frac{\sum \text{Household expenditure on food}}{\sum \text{Total household expenditure}} \times 100$$

The following (see Table 7) commonly used thresholds for classifying household food insecurity based on its food expenditure share were used:

Table 7: Commonly used HFES thresholds

HFES	Food insecurity classification
> 75%	Very vulnerable
65-75%	High food insecurity
50-65%	Medium food insecurity
< 50%	Low food insecurity

3.6.2.7 Child health

The child health indicators of the Myanmar MCCT Programme Evaluation include the following:

- Immunisation coverage;
- Period prevalence of childhood illnesses;
- Treatment-seeking behaviour for childhood illnesses; and,
- Prevalence of low birthweight

Immunisation coverage

Coverage of the expanded programme on immunisation (EPI) in Myanmar was based on the 2016 revised routine vaccination schedule when the pneumococcal and inactivated polio vaccine injection was introduced [[Central Expanded Programme on Immunization, Ministry of Health, Republic of the Union of Myanmar, 2016](#)] (see Table 8).

Table 8: Myanmar EPI Schedule 2016

Dose/Age	Vaccine
At birth	BCG
	Hepatitis B
First Dose (2nd month)	BCG
	DPT, Hepatitis B, HiB
	Pentavalent 1
	Pneumococcal Conjugate Vaccine 1
	Oral Polio Vaccine 1
	DPT, Hepatitis B, HiB
Second dose (4th month)	Pentavalent 2
	Pneumococcal Conjugate Vaccine 2
	Oral Polio Vaccine 2
	Inactivated Polio Vaccine
Third dose (6th month)	DPT, Hepatitis, HiB
	Pentavalent 3
	Pneumococcal Conjugate Vaccine 3
	Oral Polio Vaccine 3
Fourth dose (9th month)	Measles - Rubella
Fifth dose (18th Month)	Measles

Period prevalence of childhood illnesses and treatment-seeking

Mothers' report of their children's illness specifically diarrhoea, fever¹⁴ or cough¹⁵ in the past 2 weeks provides the data for assessing the period prevalence of these childhood illnesses. Then, for those whose children has been ill, mothers' reported on whether treatment was sought, where it was sought and whether payment for treatment was required are used to assess treatment-seeking.

Child birthweight

Child birthweight was assessed based on either mother's report of child's birthweight or by recorded birthweight on child's health card. However, for purposes of accuracy, only the recorded birthweight on a health card was used to determine whether child was born with low birth weight. Children with birthweight of less than 2500 grams were classified as being born with low birthweight [Woertman et al., 1993, Kelly et al., 1997].

3.6.2.8 Child nutrition

¹⁴Fever is used as a proxy for illnesses due to an infection such as malaria

¹⁵Cough is used as a proxy for an acute respiratory infection (ARI)

The Myanmar MCCT Programme evaluation assessed the following child nutrition indicators:

Childhood undernutrition

Respective childhood undernutrition indices for childhood stunting, childhood wasting and childhood underweight were assessed. The height-for-age z-scores (HAZ), weight-for-height z-scores (WAZ) and weight-for-age z-scores of each 6-59 month old child in the sample were calculated using the using the R package **zscorer**¹⁶ [Myatt and Guevarra, 2019]. Once z-score values were calculated, implausible z-score values were flagged using the WHO Growth Standards flagging criteria [World Health Organization (WHO), 2006]. This was facilitated through the **nutrheckr** R package¹⁷ [Guevarra, 2019b]. Further anthropometric data quality checks were performed using **nipnTK**¹⁸ [Myatt, 2019] R package which is based on an anthropometric data quality check toolkit developed and produced by the [National Information Platforms for Nutrition \(NIPN\)](#) and funded by the European Union (see Annex 3 for the anthropometric data quality report).

Infant and young child feeding

Infant and young child feeding indicators were assessed based on standard WHO and UNICEF definitions [World Health Organization, 2008]. The following indicators were assessed:

- Exclusive breastfeeding
- Early initiation of breastfeeding
- Minimum meal frequency
- Minimum dietary diversity
- Minimum acceptable diet

3.6.2.9 Maternal health

A standard set of maternal health indicators covering *family planning, antenatal care, birth/delivery, postnatal care* and *newborn care* were assessed.

3.6.2.10 Maternal nutrition

The following maternal nutrition indicators were assessed:

Maternal undernutrition

Maternal undernutrition was assessed based on MUAC. No consensus on MUAC cut-offs to determine maternal acute undernutrition have been identified and currently used cut-offs are specifically for pregnant and lactating women (PLW). Current research suggest that a cut-off of 23 cms as most appropriate for PLW [Ververs et al., 2013] instead of the commonly used 18.5 and 21 cms cutoffs in supplementation programmes

¹⁶See <https://nutriverse.io/zscorer> for details on how the z-scores are calculated

¹⁷See <https://nutriverse.io/nutrheckr> for details on how the flagging criteria is applied

¹⁸See <https://nutriverse.io/nipnTK> for details on how anthropometric data quality check is performed using the **nipnTK** R package

for PLW. Given that the Myanmar MCCT dataset has data for women of reproductive age and not only PLW, the 18.5 and 21 cms cutoffs were used to assess levels of acute undernutrition in mothers (see Table 9).

Table 9: MUAC cut-offs for maternal acute undernutrition

MUAC cut-offs	Classification
< 21	Global acute undernutrition
18.5 to 21	Moderate acute undernutrition
< 18.5	Severe acute undernutrition

Minimum dietary diversity for women

Minimum dietary diversity for women or MDD-W is a dichotomous indicator of whether or not women 15–49 years of age have consumed at least five out of ten defined food groups the previous day or night. The proportion of women 15–49 years of age who reach this minimum in a population can be used as a proxy indicator for higher micronutrient adequacy, one important dimension of diet quality.

The indicator is calculated as follows:

$$\text{MDD-W} = \frac{\text{Women 15-49 years of age who consumed 5 out of 10 food groups in the previous day or night}}{\text{Women 15-49 years of age}}$$

The ten food groups are:

1. Grains, white roots and tubers, and plantains
2. Pulses (beans, peas and lentils)
3. Nuts and seeds
4. Dairy
5. Meat, poultry and fish
6. Eggs
7. Dark green leafy vegetables
8. Other vitamin A-rich fruits and vegetables
9. Other vegetables
10. Other fruits

A fully reproducible data processing report can be reviewed [here](#).

Using the the recoded data for indicators, population-level estimates with 95% confidence limits of these indicators were calculated using standard classical estimation techniques. Proportion-type indicators were estimated by getting the proportion of those having the indicator of interest out of the total sample. The 95% confidence limits for a proportion-type indicator was estimated using the following formula:

$$95\% \text{ confidence limits} = p \pm 1.96 \times \sqrt{\frac{p \times (1 - p)}{n^2}}$$

where :

p = proportion estimate

n = sample size

Indicators requiring the calculation of the population mean were estimated by calculating the mean of the sample for the indicator of interest. The 95% confidence limits for a population mean indicator was estimated using the following formula:

$$95\% \text{ confidence limits} = \sigma \pm 1.96 \times \frac{SD}{\sqrt{n}}$$

where :

σ = sample mean

SD = standard deviation

n = sample size

Indicator estimation was performed on the dataset for respondents for the population-based survey component of the study.

3.6.3 Regression discontinuity

Using data from respondents who qualify for eligibility to the second component of the study, outcome nutrition indicators for children and mothers were plotted by date of birth of the index child to demonstrate whether a regression discontinuity on the indicators of interest were demonstrable between those who are not benefitting from the MCCT programme (on the basis of being born up to a month before the start of the programme) and those who are (born up to a month after start of the programme). Discontinuity was detected visually through the plot and statistically by testing any difference between the regression line of those in the control group to the regression line of those in the treatment group.

4 Results

4.1 Population-representative survey

4.1.1 Poverty Probability Index




Figure 1: Poverty probability index by location type

Poverty probability of households across different locations in Kayah state show high variability ranging from as low as 1.4% to as high as a little over 95%. This distribution of poverty is roughly similar across urban, rural and hard-to-reach areas as seen in Figure 1 above.

The poverty probabilities was primarily used to classify households into wealth quintiles which was subsequently used for wealth stratification of indicator results.

4.1.2 Food consumption score

Table 10: Food consumption score

	Mean food consumption score	Poor (%)	Borderline (%)	Acceptable (%)
Kayah				
<i>Geographic</i>				
Rural	66.0	0.0	2.8	97.2
Urban	75.6	0.0	0.6	99.4
Hard-to-reach	43.2	4.9	28.1	67.0
<i>Wealth</i>				
Wealthiest	77.1	0.0	0.0	100.0
Wealthy	72.0	0.0	0.0	100.0
Medium	63.0	0.0	6.7	93.3
Poor	52.4	1.5	16.8	81.6
Poorest	40.1	6.3	32.7	61.0

Table 10 presents FCS results by geographic location and by wealth quintiles. There is a clear geographic and wealth gradient with FCS as shown in Figure 2 and Figure ???. Poor FCS is at 4.90463215258856% in hard-to-reach areas whilst it is only 0% and 0% in urban and rural areas respectively. In the poor and poorest households, poor FCS is at 1.53061224489796% and 6.34146341463415% respectively compared to no households with poor FCS in medium, wealthy and wealthiest households.




Figure 2: Food consumption score classification by location type




Figure 3: Food consumption score classification by wealth quintiles

4.1.3 Food consumption score - nutrition

Table II: Food consumption score - nutrition

	Vitamin A-rich foods			Protein-rich foods			Heme iron-rich foods		
	Never (%)	Sometimes (%)	At least daily (%)	Never (%)	Sometimes (%)	At least daily (%)	Never (%)	Sometimes (%)	At least daily (%)
Kayah									
<i>Geographic</i>									
Rural	0.0	4.1	95.9	0.3	15.0	84.6	8.5	61.1	30.4
Urban	0.0	5.8	94.2	0.3	4.1	95.6	0.6	49.7	49.7
Hard-to-reach	3.0	31.1	65.9	10.9	53.1	36.0	20.4	69.5	10.1
<i>Wealth</i>									
Wealthiest	0.0	5.2	94.8	0.0	3.0	97.0	0.4	47.6	51.9
Wealthy	0.0	3.2	96.8	0.0	8.6	91.4	1.1	54.8	44.1
Medium	0.5	11.5	88.0	1.0	19.1	79.9	6.2	67.9	25.8
Poor	1.5	21.4	77.0	4.1	38.3	57.7	12.8	71.9	15.3
Poorest	2.4	30.7	66.8	14.6	58.0	27.3	29.8	61.5	8.8




Figure 4: Vitamin A food consumption score classification by location type




Figure 5: Vitamin A food consumption score classification by wealth quintiles



Figure 6: Protein food consumption score classification by location type




Figure 7: Protein food consumption score classification by wealth quintiles

4.1.4 Coping strategies index

4.1.4.1 Consumption-based coping strategies index

Table 12: Consumption-based Coping Strategies Index

Mean CSI	
Kayah	
<i>Geographic</i>	
Rural	4.8
Urban	3.6
Hard-to-reach	4.0
<i>Wealth</i>	
Wealthiest	2.2
Wealthy	4.2
Medium	4.8
Poor	5.2
Poorest	4.6




Figure 8: Consumption-based coping strategies index by location type




Figure 9: Consumption-based coping strategies index by wealth quintiles

4.I.4.2 Livelihood-based coping strategies index

Table 13: Livelihoods-based coping strategies index

	Secure (%)	Stress (%)	Crisis (%)	Emergency (%)
Kayah				
<i>Geographic</i>				
Rural	40.8	38.9	17.2	3.1
Urban	49.4	37.8	9.9	2.9
Hard-to-reach	44.4	46.9	3.5	5.2
<i>Wealth</i>				
Wealthiest	49.8	36.8	10.4	3.0
Wealthy	43.0	40.9	12.4	3.8
Medium	43.5	40.2	14.4	1.9
Poor	41.3	44.9	7.7	6.1
Poorest	45.9	45.4	4.4	4.4




Figure 10: Livelihoods-based coping strategies index by location type




Figure 11: Livelihoods-based coping strategies index by wealth quintiles

4.1.5 Household food expenditure share




Figure 12: Household food expenditure share by location type




Figure 13: Household food expenditure share by wealth quintiles

Table 14: Household food expenditure share

		Food Insecurity by HFES			
		Vulnerable (%)	High (%)	Medium (%)	Low (%)
Kayah					
<i>Geographic</i>					
	Rural	0.3	0.0	0.0	99.7
	Urban	0.0	0.0	0.3	99.7
	Hard-to-reach	6.3	0.5	1.4	91.8
<i>Wealth</i>					
	Wealthiest	0.0	0.0	0.4	99.6
	Wealthy	0.5	0.0	0.0	99.5
	Medium	0.5	0.0	0.0	99.5
	Poor	3.6	0.5	0.0	95.9
	Poorest	7.3	0.5	2.4	89.8




Figure 14: Household food expenditure share by location type




Figure 15: Household food expenditure share by wealth quintiles

4.1.6 Water, sanitation and hygiene

4.1.6.1 Water services ladder

Table 15: Water service ladders

	Summer Season			Rainy Season			Winter Season		
	At least limited (%)	Unimproved (%)	Surface water (%)	At least limited (%)	Unimproved (%)	Surface water (%)	At least limited (%)	Unimproved (%)	Surface water (%)
Kayah									
<i>Geographic</i>									
Rural	74.6	7.8	17.6	80.3	10.7	9.1	74.9	7.5	17.6
Urban	89.5	4.9	5.5	89.0	5.5	5.5	89.8	4.9	5.2
Hard-to-reach	63.8	17.0	19.2	68.8	17.0	14.2	64.1	16.7	19.2
<i>Wealth</i>									
Wealthiest	90.0	3.0	6.9	92.6	2.2	5.2	90.5	2.6	6.9
Wealthy	84.9	4.3	10.8	83.9	6.5	9.7	85.5	3.8	10.8
Medium	67.5	11.5	21.1	73.7	12.4	13.9	67.0	12.0	21.1
Poor	64.8	12.8	22.4	71.9	14.8	13.3	65.8	12.8	21.4
Poorest	70.2	19.5	10.2	71.7	21.0	7.3	70.2	19.0	10.7




Figure 16: Water services ladder by location type




Figure 17: Water services ladder by wealth quintiles

4.1.6.2 Sanitation services ladder

Table 16: Sanitation service ladders

	Basic (%)	Limited (%)	Unimproved (%)	Open Defecation (%)
Kayah				
<i>Geographic</i>				
Rural	7.5	42.9	49.2	0.3
Urban	26.2	47.4	25.6	0.3
Hard-to-reach	4.1	27.4	47.7	20.0
<i>Wealth</i>				
Wealthiest	26.8	56.3	16.0	0.0
Wealthy	17.2	46.2	36.6	0.0
Medium	8.6	34.9	53.6	2.9
Poor	5.1	31.1	56.6	6.6
Poorest	3.4	23.9	44.4	27.3




Figure 18: Sanitation services ladder by location type




Figure 19: Water services ladder by wealth quintiles

4.1.6.3 Handwashing service ladders

Table 17: Handwashing service ladders

	Basic (%)	Limited (%)	No Facility (%)
Kayah			
<i>Geographic</i>			
Rural	86.2	0.0	4.7
Urban	89.8	0.0	2.9
Hard-to-reach	38.4	2.7	10.6
<i>Wealth</i>			
Wealthiest	90.9	0.0	4.3
Wealthy	91.4	0.0	3.2
Medium	81.3	1.0	2.9
Poor	58.7	1.0	7.1
Poorest	28.8	2.9	13.7




Figure 20: Handwashing services ladder by location type




Figure 21: Handwashing services ladder by wealth quintiles

4.1.7 Child health

4.1.7.1 Immunisation coverage

Table 18: Immunisation coverage

	Ever vaccinated (%)	Vaccination card retention rate (%)	Immunisation access (%)	Immunisation utilisation (%)	Full immunisation coverage (%)	Hepatitis B immunisation given within first 24 hours (%)
Kayah						
<i>Geographic</i>						
Rural	100.0	89.7	97.1	83.8	32.4	56.8
Urban	100.0	92.3	93.4	87.9	34.1	72.8
Hard-to-reach	89.7	49.0	86.3	32.5	8.5	21.3
<i>Wealth</i>						
Wealthiest	100.0	93.2	93.2	90.9	36.4	72.3
Wealthy	100.0	92.2	95.3	85.9	29.7	58.8
Medium	100.0	60.0	98.0	52.0	24.0	50.4
Poor	89.8	72.7	82.0	56.0	12.0	23.7
Poorest	87.9	51.0	86.2	31.0	10.3	21.2




Figure 22: Immunisation coverage by location type




Figure 23: Immunisation coverage by wealth quintiles

Table 19: Immunisation coverage per vaccine type

	BCG (%)	Hepatitis B (%)	Penta 1 (%)	Penta 2 (%)	Penta 3 (%)	OPV 1 (%)	OPV 2 (%)	OPV 3 (%)	IPV (%)	Measles 1 (%)	Measles 2 (%)	Rubella (%)	Pneumococcal (%)
Kayah													
<i>Geographic</i>													
Rural	17.1	11.9	15.2	15.8	15.2	15.5	15.8	14.7	15.2	15.5	7.2	14.5	11.1
Urban	20.6	16.2	20.1	20.1	19.6	20.1	19.1	18.6	19.9	17.9	10.9	14.5	15.3
Hard-to-reach	18.6	6.1	7.9	7.7	7.4	8.1	7.6	6.5	7.7	6.6	4.8	8.5	5.7
<i>Wealth</i>													
Wealthiest	17.7	14.7	17.7	17.7	17.3	17.7	16.5	15.6	17.3	16.0	9.1	13.9	12.6
Wealthy	22.8	17.2	21.3	22.1	21.3	22.1	21.7	20.2	22.1	20.6	12.0	16.1	17.2
Medium	16.4	7.4	9.7	9.7	9.0	9.0	9.0	8.7	9.0	8.7	5.4	9.0	7.0
Poor	17.2	7.9	12.1	12.1	11.7	12.6	13.0	11.7	12.1	10.9	5.4	10.9	6.3
Poorest	18.8	6.4	7.9	7.5	7.5	8.3	7.1	6.4	7.9	7.1	5.3	9.8	7.5




Figure 24: Immunisation coverage per vaccine by location type



Figure 25: Immunisation coverage per vaccine by wealth quintiles

Table 20: Coverage of immunisation-associated services

		Vitamin A supplementation (%)	Deworming (%)
Kayah			
<i>Geographic</i>			
	Rural	72.6	28.1
	Urban	71.7	27.2
	Hard-to-reach	47.3	16.7
<i>Wealth</i>			
	Wealthiest	68.3	21.8
	Wealthy	71.1	34.2
	Medium	67.3	16.8
	Poor	52.7	22.6
	Poorest	45.8	20.3




Figure 26: Immunisation-associated services coverage by location type




Figure 27: Immunisation-associated services coverage by wealth quintiles

4.1.8 Period prevalence of childhood illnesses

Table 21: Period prevalence of childhood illnesses

	Diarrhoea (%)	ARI (%)	Fever (%)
Kayah			
<i>Geographic</i>			
Rural	1.5	16.6	15.7
Urban	1.9	10.1	21.5
Hard-to-reach	6.5	15.6	15.6
<i>Wealth</i>			
Wealthiest	1.9	12.2	17.8
Wealthy	0.9	13.7	22.6
Medium	3.8	15.2	18.6
Poor	2.5	14.6	13.6
Poorest	9.2	15.7	15.3




Figure 28: Period prevalence of childhood illnesses by location type




Figure 29: Period prevalence of childhood illnesses by wealth quintiles

4.1.9 Treatment-seeking behaviour

4.1.9.1 Diarrhoea

Table 22: Treatment-seeking for diarrhoea

			Reasons for not seeking treatment						
	Sought treatment (%)	Time to treatment (days)	No facility (%)	Facility inaccessible (%)	Expensive (%)	Not necessary (%)	Advised not to (%)	Alternative treatment (%)	Do not know treatment (%)
Kayah									
<i>Geographic</i>									
Rural	100.0	1.4	0.0	0.0	0.0	0.0	0	0.0	0
Urban	57.1	0.9	0.0	0.0	0.0	0.0	0	33.3	0
Hard-to-reach	66.7	3.7	11.8	41.2	17.6	5.9	0	0.0	0
<i>Wealth</i>									
Wealthiest	75.0	1.7	0.0	0.0	0.0	0.0	0	0.0	0
Wealthy	50.0	0.7	0.0	0.0	0.0	0.0	0	33.3	0
Medium	80.0	1.1	0.0	33.3	0.0	0.0	0	0.0	0
Poor	40.0	2.4	25.0	25.0	0.0	0.0	0	0.0	0
Poorest	71.4	4.2	8.3	41.7	25.0	8.3	0	0.0	0




Figure 30: Treatment-seeking for diarrhoea by location type




Figure 31: Treatment-seeking for diarrhoea by wealth quintiles




Figure 32: Reasons for not seeking treatment by location type

Table 23: Where/who treatment is sought from

		Where/who treatment is sought from												
		Township hospital (%)	Station hospital (%)	RHC/ health assistant (%)	SRHC/ midwife (%)	Private clinic/ doctor (%)	Community health worker (%)	Traditional healer (%)	Untrained health worker (%)	Drug from shop (%)	EHO clinic/ volunteer (%)	Family member (%)	NGOs/ clinic (%)	Auxilliary midwife (%)
Kayah <i>Geographic</i>														
	Rural	17.6	5.9	11.8	41.2	0.0	0.0	0	0	17.6	0.0	0	5.9	0.0
	Urban	12.5	0.0	0.0	0.0	25.0	0.0	0	0	25.0	12.5	0	0.0	0.0
	Hard-to-reach	0.0	0.0	0.0	36.1	0.0	8.3	0	0	8.3	19.4	0	11.1	8.3
Wealth														
	Wealthiest	28.6	0.0	14.3	42.9	0.0	0.0	0	0	0.0	0.0	0	0.0	0.0
	Wealthy	0.0	0.0	0.0	14.3	14.3	0.0	0	0	28.6	14.3	0	14.3	0.0
	Medium	9.1	9.1	9.1	36.4	9.1	0.0	0	0	18.2	0.0	0	0.0	0.0
	Poor	10.0	0.0	0.0	20.0	0.0	10.0	0	0	10.0	0.0	0	10.0	20.0
	Poorest	0.0	0.0	0.0	38.5	0.0	7.7	0	0	11.5	26.9	0	11.5	3.8




Figure 33: Where/who treatment is sought from by location type

Table 24: Payment for treatment

		Payment for						
	Payment for service (MMK)	Transportation (%)	Registration (%)	Medicine (%)	Laboratory fees (%)	Provider fees (%)	Gifts (%)	Took loan (%)
Kayah								
Geographic								
Rural	10823.5	0	0	100	0	0	0	50.0
Urban	5637.5	0	0	100	0	0	0	28.6
Hard-to-reach	458.3	0	0	100	0	0	0	0.0
Wealth								
Wealthiest	13157.1	NaN	NaN	NaN	NaN	NaN	NaN	40.0
Wealthy	5928.6	0	0	100	0	0	0	33.3
Medium	8227.3	0	0	100	0	0	0	42.9
Poor	950.0	0	0	100	0	0	0	25.0
Poorest	461.5	0	0	100	0	0	0	0.0




Figure 34: Costs incurred for treatment

4.1.9.2 Acute respiratory infection

Table 25: Treatment-seeking for acute respiratory infection

			Reasons for not seeking treatment						
	Sought treatment (%)	Time to treatment (days)	No facility (%)	Facility inaccessible (%)	Expensive (%)	Not necessary (%)	Advised not to (%)	Alternative treatment (%)	Do not know treatment (%)
Kayah									
<i>Geographic</i>									
Rural	88.9	1.1	0.0	0.0	6.2	0.0	0	0.0	0.0
Urban	71.1	1.2	0.0	0.0	15.0	0.0	0	15.0	0.0
Hard-to-reach	63.9	5.9	12.5	42.5	12.5	2.5	0	2.5	2.5
<i>Wealth</i>									
Wealthiest	69.2	1.2	0.0	0.0	0.0	0.0	0	22.2	0.0
Wealthy	83.9	1.1	0.0	0.0	0.0	0.0	0	8.3	0.0
Medium	72.5	1.5	0.0	5.9	23.5	0.0	0	0.0	0.0
Poor	79.3	2.6	7.7	69.2	23.1	0.0	0	0.0	0.0
Poorest	63.9	10.0	17.4	30.4	8.7	4.3	0	4.3	4.3




Figure 35: Treatment-seeking for acute respiratory infection by location type




Figure 36: Treatment-seeking for acute respiratory infection by wealth quintiles




Figure 37: Reasons for not seeking treatment by location type

Table 26: Where/who treatment is sought from

	Where/who treatment is sought from												
	Township hospital (%)	Station hospital (%)	RHC/ health assistant (%)	SRHC/ midwife (%)	Private clinic/ doctor (%)	Community health worker (%)	Traditional healer (%)	Untrained health worker (%)	Drug from shop (%)	EHO clinic/ volunteer (%)	Family member (%)	NGOs/ clinic (%)	Auxilliary midwife (%)
Kayah													
<i>Geographic</i>													
Rural	3.6	7.1	11.9	32.1	17.9	1.2	0.0	0	14.3	0.0	2.4	2.4	1.2
Urban	23.4	8.5	0.0	4.3	48.9	0.0	0.0	0	6.4	0.0	4.3	0.0	0.0
Hard-to-reach	2.1	5.3	5.3	36.2	1.1	3.2	1.1	0	2.1	21.3	1.1	2.1	13.8
<i>Wealth</i>													
Wealthiest	11.5	7.7	11.5	7.7	46.2	0.0	0.0	0	0.0	0.0	3.8	0.0	0.0
Wealthy	9.3	5.6	5.6	29.6	22.2	0.0	0.0	0	16.7	0.0	3.7	1.9	1.9
Medium	11.3	11.3	3.8	32.1	18.9	0.0	0.0	0	9.4	3.8	0.0	1.9	3.8
Poor	2.1	2.1	10.4	33.3	2.1	4.2	0.0	0	4.2	16.7	4.2	2.1	8.3
Poorest	2.4	4.9	2.4	29.3	7.3	4.9	2.4	0	2.4	24.4	0.0	2.4	17.1




Figure 38: Where/who treatment is sought from by location type

Table 27: Payment for treatment

	Payment for service (MMK)	Payment for						Took loan (%)
		Transportation (%)	Registration (%)	Medicine (%)	Laboratory fees (%)	Provider fees (%)	Gifts (%)	
Kayah								
<i>Geographic</i>								
Rural	5264.3	0.0	2.5	77.5	0	2.5	2.5	21.9
Urban	8755.3	0.0	0.0	81.8	0	0.0	0.0	6.8
Hard-to-reach	22613.8	6.5	0.0	67.7	0	0.0	0.0	31.4
<i>Wealth</i>								
Wealthiest	8403.8	0.0	0.0	75.0	0	0.0	0.0	0.0
Wealthy	5733.3	0.0	3.7	81.5	0	3.7	0.0	22.0
Medium	4162.3	0.0	0.0	64.0	0	0.0	4.0	23.7
Poor	45097.9	5.9	0.0	76.5	0	0.0	0.0	14.3
Poorest	1317.1	9.1	0.0	81.8	0	0.0	0.0	46.7




Figure 39: Costs incurred for treatment by location type

4.1.9.3 Fever

Table 28: Treatment-seeking for fever

			Reasons for not seeking treatment						
	Sought treatment (%)	Time to treatment (days)	No facility (%)	Facility inaccessible (%)	Expensive (%)	Not necessary (%)	Advised not to (%)	Alternative treatment (%)	Do not know treatment (%)
Kayah									
<i>Geographic</i>									
Rural	76.5	1.4	0.0	0.0	5.0	5.0	0	0.0	0
Urban	85.2	1.1	0.0	0.0	14.3	0.0	0	19.0	0
Hard-to-reach	65.3	2.0	13.5	48.6	2.7	0.0	0	5.4	0
<i>Wealth</i>									
Wealthiest	76.3	1.2	0.0	0.0	0.0	0.0	0	9.1	0
Wealthy	84.3	1.1	0.0	0.0	0.0	6.7	0	6.7	0
Medium	73.5	1.6	0.0	31.2	6.2	0.0	0	18.8	0
Poor	70.4	1.8	17.6	35.3	23.5	0.0	0	5.9	0
Poorest	68.6	2.0	11.8	41.2	0.0	0.0	0	0.0	0




Figure 40: Treatment-seeking for fever by location type




Figure 41: Treatment-seeking for fever by wealth quintiles




Figure 42: Reasons for not seeking treatment by location type

Table 29: Where/who treatment is sought from

	Where/who treatment is sought from												
	Township hospital (%)	Station hospital (%)	RHC/ health assistant (%)	SRHC/ midwife (%)	Private clinic/ doctor (%)	Community health worker (%)	Traditional healer (%)	Untrained health worker (%)	Drug from shop (%)	EHO clinic/ volunteer (%)	Family member (%)	NGOs/ clinic (%)	Auxilliary midwife (%)
Kayah													
<i>Geographic</i>													
Rural	8.3	1.2	15.5	26.2	28.6	1.2	0	0.0	9.5	1.2	2.4	1.2	0.0
Urban	32.2	3.3	0.0	3.3	45.6	0.0	0	2.2	3.3	2.2	2.2	0.0	0.0
Hard-to-reach	2.1	2.1	11.6	36.8	0.0	1.1	0	3.2	3.2	12.6	4.2	3.2	14.7
<i>Wealth</i>													
Wealthiest	22.5	2.5	2.5	7.5	55.0	0.0	0	0.0	0.0	2.5	5.0	0.0	0.0
Wealthy	16.7	1.4	8.3	23.6	33.3	1.4	0	0.0	8.3	1.4	0.0	0.0	0.0
Medium	20.0	4.6	15.4	24.6	16.9	0.0	0	0.0	4.6	1.5	3.1	1.5	4.6
Poor	4.3	2.2	8.7	23.9	6.5	0.0	0	8.7	2.2	6.5	4.3	2.2	15.2
Poorest	2.4	0.0	7.3	29.3	4.9	2.4	0	2.4	9.8	22.0	4.9	4.9	9.8




Figure 43: Where/who treatment is sought from by location type

Table 30: Payment for treatment

	Payment for service (MMK)	Payment for						Took loan (%)
		Transportation (%)	Registration (%)	Medicine (%)	Laboratory fees (%)	Provider fees (%)	Gifts (%)	
Kayah								
<i>Geographic</i>								
Rural	8614.3	2.9	0	82.9	0	0.0	2.9	30.8
Urban	9549.4	0.0	0	78.1	0	3.1	0.0	16.2
Hard-to-reach	1434.7	0.0	0	77.1	0	0.0	0.0	12.8
<i>Wealth</i>								
Wealthiest	8989.5	0.0	0	66.7	0	0.0	0.0	18.2
Wealthy	10522.8	3.2	0	80.6	0	3.2	0.0	17.5
Medium	6789.2	0.0	0	74.2	0	0.0	3.2	27.5
Poor	1663.0	0.0	0	89.5	0	0.0	0.0	22.7
Poorest	1251.2	0.0	0	78.6	0	0.0	0.0	18.8




Figure 44: Costs incurred for treatment in Kayah state

4.1.10 Child nutrition

4.1.10.1 Prevalence of childhood stunting

<<<<<<< HEAD

Table 31: Child stunting/stuntedness

	Height-for-age z-score	Global stunting/ stuntedness (%)	Moderate stunting/ stuntedness (%)	Severe stunting/ stuntedness (%)
Kayah				
<i>Geographic</i>				
Rural	-1.4	31.5	25.7	5.9
Urban	-1.1	17.4	14.9	2.6
Hard-to-reach	-1.6	47.9	30.1	17.9

4.1.10.2 Prevalence of childhood underweight

Table 32: Child underweight

	Weight-for-age z-score	Global underweight (%)	Moderate underweight (%)	Severe underweight (%)
Kayah				
<i>Geographic</i>				
Rural	-1.1	14.5	11.4	3.1
Urban	-0.9	14.6	12.6	2.0
Hard-to-reach	-1.1	24.8	20.4	4.3

4.1.10.3 Prevalence of childhood wasting by weight-for-height z-score

Table 33: Child wasting by weight-for-height z-score

	Weight-for-height z-score	Global acute malnutrition (%)	Moderate acute malnutrition (%)	Severe acute malnutrition (%)
Kayah				
<i>Geographic</i>				
Rural	-0.4	5.9	4.6	1.2
Urban	-0.5	3.8	2.6	1.3
Hard-to-reach	-0.3	4.0	2.5	1.5

4.1.10.4 Prevalence of childhood wasting by MUAC

Table 34: Child wasting by MUAC

	MUAC (cm)	Global acute malnutrition (%)	Moderate acute malnutrition (%)	Severe acute malnutrition (%)
Kayah				
<i>Geographic</i>				
Rural	15.2	2.7	2.2	0.5
Urban	15.1	2.5	1.0	1.5
Hard-to-reach	14.5	4.7	3.2	1.5

4.1.10.5 Infant and young child feeding

Table 35: Infant and young child feeding

	Early initiation of breastfeeding (%)	Exclusive breastfeeding (%)	Minimum meal frequency (%)	Minimum dietary diversity (%)	Minimum acceptable diet (%)
Kayah					
<i>Geographic</i>					
Rural	87.8	12.3	55.4	43.2	24.1
Urban	91.5	8.1	50.4	54.3	30.8
Hard-to-reach	76.3	4.2	27.3	14.0	9.3
<i>Wealth</i>					
Wealthiest	89.2	10.4	51.5	50.0	39.4
Wealthy	91.2	9.2	50.0	51.4	21.1
Medium	78.2	8.1	44.6	33.3	18.1
Poor	86.6	5.9	37.6	25.2	15.1
Poorest	76.6	5.0	26.1	5.9	5.8




Figure 45: Infant and young child feeding by location type




Figure 46: Infant and young child feeding by wealth quintiles

4.1.11 Maternal health

4.1.11.1 Family planning

Table 36: Months to wait until next pregnancy

	Time to wait for next child (months)	Time to wait for next child for currently pregnant (months)	Time to wait for next child according to husband (months)
Kayah			
<i>Geographic</i>			
Rural	47.1	41.2	48.2
Urban	46.1	28.8	43.6
Hard-to-reach	39.9	23.6	37.8
<i>Wealth</i>			
Wealthiest	48.9	41.5	45.9
Wealthy	43.7	31.6	42.9
Medium	44.0	33.3	44.4
Poor	41.6	26.5	39.6
Poorest	40.5	22.8	39.0

Table 37: Use of contraceptive methods

		Contraceptive methods														
	Using contraceptives now (%)	Used contraceptives before (%)	Female sterilisation (%)	Male sterilisation (%)	IUD (%)	Implant (%)	Injectable (%)	Pills (%)	Male condom (%)	Female condom (%)	Lactational amenorrhea (%)	Rhythm method (%)	Withdrawal (%)	Emergency contraception (%)	Diaphragm (%)	Foam/jelly (%)
Kayah																
<i>Geographic</i>																
Rural	59.1	54.2	7.4	0.6	7.4	4.3	63.2	15.3	0.0	0	0	1.2	0.6	0	0	0
Urban	71.6	64.6	6.4	0.0	5.9	15.4	56.9	14.4	1.1	0	0	0.0	0.0	0	0	0
Hard-to-reach	31.3	31.3	0.8	0.0	0.8	18.3	62.7	16.7	0.0	0	0	0.8	0.0	0	0	0
<i>Wealth</i>																
Wealthiest	68.7	64.1	5.0	0.0	7.9	11.4	57.1	15.7	1.4	0	0	0.7	0.7	0	0	0
Wealthy	69.5	62.7	8.4	0.0	6.3	10.5	60.0	14.7	0.0	0	0	0.0	0.0	0	0	0
Medium	55.2	51.2	4.5	0.0	3.4	8.0	69.3	13.6	0.0	0	0	1.1	0.0	0	0	0
Poor	46.9	39.0	4.4	1.1	3.3	14.4	58.9	16.7	0.0	0	0	1.1	0.0	0	0	0
Poorest	27.3	30.3	3.1	0.0	1.6	20.3	59.4	15.6	0.0	0	0	0.0	0.0	0	0	0

Table 38: Where contraception services received

	Government hospital (%)	Government health centre (%)	Government village health worker (%)	UHC/MCH centre (%)	Private hospital (%)	Private doctor (%)	Pharmacy (%)	NGO (%)	EHO clinic (%)	Auxilliary midwife (%)	Midwife (%)
Kayah											
<i>Geographic</i>											
Rural	21.2	8.3	1.0	3.6	1.6	5.7	3.6	1.0	5.2	6.2	42.5
Urban	42.2	4.9	0.0	17.2	6.4	7.8	7.8	0.0	2.0	1.5	10.3
Hard-to-reach	8.0	4.5	7.1	3.6	0.0	1.8	0.0	4.5	20.5	17.0	33.0
<i>Wealth</i>											
Wealthiest	36.8	7.6	0.0	11.8	6.2	6.2	9.7	0.0	2.8	0.7	18.1
Wealthy	30.2	4.3	0.0	9.5	5.2	8.6	6.9	0.9	2.6	5.2	26.7
Medium	22.6	7.5	0.9	10.4	0.9	5.7	0.0	1.9	5.7	5.7	38.7
Poor	21.2	4.7	5.9	3.5	0.0	4.7	1.2	2.4	10.6	12.9	32.9
Poorest	10.3	5.2	6.9	6.9	0.0	0.0	0.0	3.4	25.9	17.2	24.1

Table 39: Information on family planning/contraception

	Received family planning information (%)	Source of family planning information											Appropriate family planning knowledge (%)
		Government hospital (%)	Government health centre (%)	Government village health worker (%)	UHC/MCH centre (%)	Private hospital (%)	Private doctor (%)	Pharmacy (%)	NGO (%)	EHO clinic (%)	Auxilliary midwife (%)	Midwife (%)	
Kayah													
<i>Geographic</i>													
Rural	84.1	18.5	7.1	2.4	6.0	0.0	3.6	0	0.6	3.6	5.4	53.0	89.7
Urban	93.0	38.9	4.0	2.0	18.8	4.0	5.4	0	1.3	2.0	0.7	22.8	97.6
Hard-to-reach	64.6	5.8	2.2	5.8	2.2	0.0	0.0	0	2.9	23.7	7.2	50.4	84.8
<i>Wealth</i>													
Wealthiest	93.5	33.7	5.9	1.0	19.8	3.0	5.0	0	1.0	4.0	1.0	25.7	94.5
Wealthy	88.1	23.3	3.3	3.3	8.9	2.2	6.7	0	2.2	3.3	3.3	43.3	94.9
Medium	82.6	22.8	7.9	4.0	5.0	1.0	1.0	0	1.0	4.0	4.0	49.5	92.0
Poor	71.2	11.5	3.8	3.8	7.7	0.0	2.6	0	1.3	11.5	7.7	50.0	88.1
Poorest	64.1	11.6	1.2	4.7	2.3	0.0	0.0	0	2.3	25.6	7.0	45.3	83.3

4.1.12 Antenatal care

4.1.12.1 Antenatal care coverage

Table 40: Antenatal care coverage

Costs incurred for ANC services												
	One ANC visit (%)	Four ANC visits (%)	Incur cost for ANC? (%)	Cost amount (MMK)	Transport (%)	Registration (%)	Medicine (%)	Laboratory (%)	Provider (%)	Gifts (%)	Took a loan? (%)	
Kayah <i>Geographic</i>												
	Rural	28.4	28.4	51.9	53937.5	40.0	5.0	47.5	37.5	47.5	5.0	22.5
	Urban	31.4	28.0	63.5	95016.4	34.4	4.9	77.0	47.5	78.7	1.6	14.8
	Hard-to-reach	26.7	23.0	5.6	140500.0	66.7	16.7	66.7	16.7	16.7	0.0	16.7
Wealth												
	Wealthiest	27.4	25.0	77.8	103285.7	31.4	5.7	68.6	57.1	77.1	0.0	8.6
	Wealthy	31.9	29.7	60.4	59781.2	31.2	3.1	71.9	43.8	75.0	6.2	21.9
	Medium	27.9	23.8	36.7	68888.9	50.0	0.0	61.1	38.9	55.6	5.6	16.7
	Poor	28.7	27.9	22.8	49192.3	46.2	7.7	53.8	15.4	7.7	0.0	38.5
	Poorest	28.7	25.0	4.3	103000.0	66.7	0.0	33.3	0.0	0.0	0.0	0.0




Figure 47: Antenatal care coverage by location type




Figure 48: Antenatal care coverage by wealth quintiles

4.1.12.2 Antenatal care counselling

Table 41: Antenatal care counselling

		Restricted eating...							
		Attended ANC counselling (%)	Restricted eating certain foods? (%)	Vegetables (%)	Fruits (%)	Grains (%)	Meat (%)	Fish (%)	Dairy (%)
Kayah <i>Geographic</i>									
	Rural	93.5	18.3	13.3	13.3	0.0	6.7	20.0	0.0
	Urban	79.2	19.8	10.5	21.1	5.3	15.8	15.8	5.3
	Hard-to-reach	72.2	19.7	14.3	25.0	3.6	32.1	10.7	3.6
<i>Wealth</i>									
	Wealthiest	93.3	17.8	37.5	25.0	0.0	0.0	25.0	0.0
	Wealthy	81.1	26.4	7.1	14.3	7.1	7.1	14.3	0.0
	Medium	71.4	13.0	14.3	14.3	0.0	0.0	0.0	0.0
	Poor	82.5	17.6	16.7	33.3	0.0	16.7	25.0	0.0
	Poorest	74.3	19.4	5.6	11.1	5.6	44.4	5.6	5.6

4.1.12.3 Antenatal care supplementation

Table 42: Antenatal care supplementation

						Source of iron-folic acid tablets				
						Government hospital (%)	EHO clinic (%)	Private doctor/ clinic (%)	SRHC/ RHC (%)	Routine ANC location in village/ward (%)
						Vitamin B1 (%)	Iron folic acid (%)	No. of days IFA taken (days)	Cost of IFA (MMK)	
Kayah										
Geographic										
Rural		98.8	96.3	176.5	1124.9	6.3	0.0	6.3	73.4	16.5
Urban		85.4	95.8	164.9	5610.8	28.3	1.1	12.0	17.4	17.4
Hard-to-reach		74.6	69.7	113.2	671.2	2.0	23.2	0.0	45.5	34.3
Wealth										
Wealthiest		88.9	100.0	166.1	2383.8	15.6	0.0	17.8	26.7	11.1
Wealthy		96.2	94.3	175.1	1701.8	24.0	2.0	10.0	38.0	16.0
Medium		88.9	90.7	152.9	9.2	18.4	4.1	2.0	61.2	22.4
Poor		82.4	80.9	160.3	186.3	7.3	16.4	0.0	45.5	27.3
Poorest		73.1	68.8	112.4	1031.2	0.0	18.8	0.0	46.9	37.5

4.1.12.4 Antenatal care testing

Table 43: Antenatal care testing

	Tests done? (%)	Type of test				Location test was done						Incur costs? (%)	Cost amount (MMK)
		Hepatitis B (%)	Hepatitis C (%)	HIV/AIDS (%)	Syphillis (%)	Government hospital (%)	EHO clinic (%)	Private doctor/ clinic (%)	SRHC/ RHC (%)	Routine ANC location in village/ward (%)			
Kayah													
<i>Geographic</i>													
Rural	35.3	8.8	0	20.6	0.0	0.0	0.0	8.8	26.5	8.8	40.0	4091.7	
Urban	37.0	0.0	0	25.9	3.7	7.4	3.7	3.7	11.1	3.7	54.4	7580.8	
Hard-to-reach	21.5	0.0	0	0.8	13.1	2.3	6.2	0.0	7.7	7.7	7.5	1886.1	
<i>Wealth</i>													
Wealthiest	37.5	0.0	0	37.5	0.0	12.5	0.0	0.0	25.0	0.0	57.5	9486.0	
Wealthy	50.0	9.1	0	27.3	0.0	4.5	0.0	13.6	31.8	9.1	50.0	5558.2	
Medium	30.4	4.3	0	8.7	13.0	0.0	0.0	4.3	17.4	8.7	34.2	3118.4	
Poor	17.6	0.0	0	3.9	7.8	2.0	3.9	0.0	9.8	3.9	19.2	653.8	
Poorest	22.4	0.0	0	1.2	12.9	2.4	7.1	0.0	4.7	9.4	11.1	2794.2	

4.I.I3 Birth/delivery

Table 44: Gestational age at birth (weeks)

	Gestational age (weeks)
Kayah	
<i>Geographic</i>	
Rural	35.8
Urban	36.5
Hard-to-reach	36.6
<i>Wealth</i>	
Wealthiest	36.4
Wealthy	36.3
Medium	35.7
Poor	36.6
Poorest	36.6

Table 45: Location of birth/delivery

	Home (%)	Government hospital (%)	Private doctor/ clinic (%)	SRHC/RHC (%)	Routine ANC location in village (%)	EHO clinic (%)
Kayah						
<i>Geographic</i>						
Rural	46.2	41.0	0.0	12.8	0	0
Urban	16.5	82.4	1.1	0.0	0	0
Hard-to-reach	90.1	7.0	0.0	2.8	0	0
<i>Wealth</i>						
Wealthiest	14.3	78.6	2.4	4.8	0	0
Wealthy	29.4	62.7	0.0	7.8	0	0
Medium	41.2	52.9	0.0	5.9	0	0
Poor	76.5	17.6	0.0	5.9	0	0
Poorest	91.4	7.5	0.0	1.1	0	0

Table 46: Reason for choosing birth/delivery location

	Convenience (%)	Tradition (%)	Close distance (%)	Safety for mother/baby (%)	Affordable cost (%)
Kayah					
<i>Geographic</i>					
Rural	11.0	1.4	5.5	50.7	12.3
Urban	11.8	0.0	4.4	69.1	1.5
Hard-to-reach	26.0	34.6	0.0	9.4	22.0
<i>Wealth</i>					
Wealthiest	8.8	0.0	2.9	76.5	0.0
Wealthy	14.6	0.0	2.4	56.1	4.9
Medium	14.6	8.3	8.3	43.8	10.4
Poor	27.6	20.7	0.0	24.1	17.2
Poorest	21.0	35.8	1.2	7.4	25.9

Table 47: Who assisted in birth/delivery

	Doctor (%)	Nurse (%)	Lady health visitor (%)	Midwife (%)	Auxilliary midwife (%)	Traditional birth attendant (%)	On my own (%)	Relatives (%)	EHO cadres (%)
Kayah									
<i>Geographic</i>									
Rural	37.8	0.0	2.4	37.8	4.9	7.3	2.4	0.0	1.2
Urban	69.8	3.1	1.0	20.8	2.1	2.1	0.0	0.0	0.0
Hard-to-reach	5.6	0.7	0.0	11.3	6.3	34.5	14.1	21.8	2.1
<i>Wealth</i>									
Wealthiest	64.4	4.4	2.2	26.7	0.0	0.0	0.0	0.0	0.0
Wealthy	54.7	1.9	0.0	34.0	1.9	3.8	0.0	0.0	1.9
Medium	46.3	1.9	1.9	22.2	7.4	5.6	3.7	9.3	0.0
Poor	14.7	0.0	1.5	22.1	1.5	30.9	5.9	13.2	1.5
Poorest	6.5	0.0	0.0	10.8	9.7	33.3	17.2	18.3	2.2

Table 48: Birth/delivery method

	Normal (%)	Caesarian (%)	Vacuum (%)	Forceps (%)
Kayah				
<i>Geographic</i>				
Rural	81.7	18.3	0	0
Urban	54.2	45.8	0	0
Hard-to-reach	95.8	4.2	0	0
<i>Wealth</i>				
Wealthiest	48.9	51.1	0	0
Wealthy	75.5	24.5	0	0
Medium	70.4	29.6	0	0
Poor	94.1	5.9	0	0
Poorest	95.7	4.3	0	0

Table 49: Costs incurred for birth/delivery

	Incurred costs? (%)	Delivery cost (MMK)	Took a loan? (%)
Kayah			
<i>Geographic</i>			
Rural	84.1	9786765	39.1
Urban	93.8	17384444	27.0
Hard-to-reach	50.7	8976056	18.1
<i>Wealth</i>			
Wealthiest	95.6	17255814	20.9
Wealthy	92.5	14514286	33.3
Medium	81.5	15482954	43.2
Poor	63.2	6460714	25.6
Poorest	48.4	7922727	15.6

4.1.14 Postnatal care

Table 50: Postnatal care coverage

			Provider of postnatal care							
	Received postnatal care (%)	Time to care (days)	Doctor (%)	Nurse (%)	Lady health visitor (%)	Midwife (%)	Auxilliary midwife (%)	Traditional birth attendant (%)	Relatives (%)	EHO cadres (%)
Kayah										
<i>Geographic</i>										
Rural	75.6	84.6	30.4	4.3	2.2	60.9	0.0	0	0.0	2.2
Urban	63.5	119.0	72.7	9.1	2.3	15.9	0.0	0	0.0	0.0
Hard-to-reach	26.1	450.2	8.3	13.9	0.0	38.9	16.7	0	2.8	19.4
<i>Wealth</i>										
Wealthiest	68.9	185.1	56.0	8.0	4.0	32.0	0.0	0	0.0	0.0
Wealthy	58.5	95.0	52.0	8.0	0.0	36.0	0.0	0	0.0	4.0
Medium	61.1	123.9	40.0	16.0	0.0	44.0	0.0	0	0.0	0.0
Poor	47.1	396.3	25.0	8.3	4.2	41.7	4.2	0	0.0	16.7
Poorest	29.0	69.5	16.7	0.0	0.0	45.8	20.8	0	4.2	12.5

Table 51: Vitamin B1 supplementation

	Vitamin B1 supplementation duration (weeks)
Kayah	
<i>Geographic</i>	
Rural	148.2
Urban	102.1
Hard-to-reach	55.6
<i>Wealth</i>	
Wealthiest	101.6
Wealthy	97.8
Medium	117.0
Poor	116.9
Poorest	66.2

Table 52: Costs of postnatal care

	Costs incurred? (%)	Reason for costs					
		Transportation (%)	Registration (%)	Medicine (%)	Laboratory fees (%)	Provider fees (%)	Gifts (%)
Kayah							
Geographic							
Rural	0.5	0.1	0	75.0	0.0	0	12.5
Urban	0.5	0.6	0	28.6	0.0	0	14.3
Hard-to-reach	0.1	0.7	0	0.0	33.3	0	0.0
Wealth							
Wealthiest	0.5	0.3	0	66.7	0.0	0	0.0
Wealthy	0.5	0.0	0	33.3	0.0	0	66.7
Medium	0.5	0.6	0	20.0	20.0	0	0.0
Poor	0.4	0.3	0	66.7	0.0	0	0.0
Poorest	0.1	0.5	0	50.0	0.0	0	0.0

4.1.15 Newborn care

Table 53: Newborn care coverage

			Provider of newborn care								
	Newborn care within 24 hours (%)	Newborn care within 48 hours (%)	Doctor (%)	Nurse (%)	Lady health visitor (%)	Midwife (%)	Auxilliary midwife (%)	Traditional birth attendant (%)	Relatives (%)	EHO cadres (%)	
Kayah											
Geographic											
Rural	87.8	91.5	32.7	2.0	2.0	53.1	6.1	0.0	0.0	2.0	
Urban	90.6	93.8	70.8	6.2	2.1	18.8	2.1	0.0	0.0	0.0	
Hard-to-reach	21.8	28.9	8.3	5.6	0.0	44.4	25.0	5.6	2.8	8.3	
Wealth											
Wealthiest	95.6	97.8	61.5	3.8	3.8	30.8	0.0	0.0	0.0	0.0	
Wealthy	94.3	94.3	50.0	3.3	0.0	40.0	3.3	0.0	0.0	3.3	
Medium	74.1	79.6	46.4	7.1	0.0	35.7	10.7	0.0	0.0	0.0	
Poor	39.7	45.6	23.8	4.8	4.8	47.6	4.8	4.8	0.0	4.8	
Poorest	24.7	33.3	11.5	0.0	0.0	42.3	30.8	3.8	3.8	7.7	

Table 54: Costs of newborn care

	Costs incurred? (%)	Cost amount (MMK)	Reason for costs						Took a loan? (%)
			Transportation (%)	Registration (%)	Medicine (%)	Laboratory fees (%)	Provider fees (%)	Gifts (%)	
Kayah									
<i>Geographic</i>									
Rural	24.0	10147059	0	0	40.0	0	0.0	0.0	61.1
Urban	23.6	10916667	0	0	22.2	0	11.1	11.1	38.1
Hard-to-reach	2.4	NaN	0	0	0.0	0	0.0	0.0	100.0
<i>Wealth</i>									
Wealthiest	20.9	8277778	0	0	0.0	0	20.0	20.0	22.2
Wealthy	20.0	9494444	0	0	0.0	0	0.0	0.0	80.0
Medium	20.9	11700000	0	0	75.0	0	0.0	0.0	22.2
Poor	22.6	6250000	0	0	33.3	0	0.0	0.0	71.4
Poorest	6.5	3250000	0	0	0.0	0	0.0	0.0	50.0

Table 55: Fed colostrum to newborn

	Fed colostrum to newboarn (%)
Kayah	
<i>Geographic</i>	
Rural	96.3
Urban	89.6
Hard-to-reach	85.9
<i>Wealth</i>	
Wealthiest	95.6
Wealthy	88.7
Medium	92.6
Poor	89.7
Poorest	86.0

Table 56: Knowledge of danger signs

	Danger signs							Does not know any danger signs (%)
	Feeding less (%)	Convulsion (%)	High/low temperature (%)	Local infection (%)	No less movement (%)	Fast/difficulty breathing (%)	Yellow skin (%)	
Kayah								
<i>Geographic</i>								
Rural	0.0	5.9	3.9	0.0	2	3.9	33.3	51.0
Urban	6.5	0.0	10.9	0.0	0	4.3	41.3	37.0
Hard-to-reach	5.4	0.8	3.9	0.8	0	1.6	6.2	81.4
<i>Wealth</i>								
Wealthiest	8.0	0.0	8.0	0.0	0	0.0	52.0	32.0
Wealthy	3.2	9.7	3.2	0.0	0	9.7	32.3	41.9
Medium	3.0	3.0	6.1	0.0	3	0.0	30.3	54.5
Poor	3.8	0.0	7.5	0.0	0	3.8	5.7	79.2
Poorest	4.9	0.0	3.7	1.2	0	1.2	7.3	81.7

Table 57: Care of cord and delivery wound

	Inappropriate cord care (%)	Inappropriate delivery wound care (%)
Kayah		
<i>Geographic</i>		
Rural	49.3	36.8
Urban	53.5	26.5
Hard-to-reach	52.3	75.9
<i>Wealth</i>		
Wealthiest	50.0	26.3
Wealthy	43.8	26.5
Medium	52.0	43.5
Poor	53.2	67.8
Poorest	57.5	73.6

4.2 Maternal nutrition

4.2.1 Minimum dietary diversity for women

Table 58: Minimum dietary diversity for women

	Minimum dietary diversity for women (%)
Kayah	
<i>Geographic</i>	
Rural	89.7
Urban	89.2
Hard-to-reach	83.7
<i>Wealth</i>	
Wealthiest	89.6
Wealthy	87.6
Medium	92.8
Poor	80.6
Poorest	86.3

4.3 Cohort study

4.3.1 Child anthropometric indicators




Figure 49: Weight-for-age z-score regression discontinuity




Figure 50: Height-for-age z-score regression discontinuity




Figure 51: Weight-for-height z-score regression discontinuity




Figure 52: MUAC regression discontinuity

Annex 1: Study questionnaires

This goes here. What else?

Annex 2: Anthropometric standardisation results

Standardisation results here

Annex 3: Anthropometric data quality check

References

- Jacob Bor, Ellen Moscoe, Portia Mutevedzi, Marie Louise Newell, and Till Bärnighausen. Regression discontinuity designs in epidemiology: Causal inference without randomized trials. *Epidemiology*, 25(5): 729–737, 2014. ISSN 15315487. doi: 10.1097/EDE.000000000000138.
- Central Expanded Programme on Immunization, Ministry of Health, Republic of the Union of Myanmar. Expanded Program on Immunization Multi Year Plan 2017-2021 Revised January 2016. Technical report, 2016.
- Mercedes de Onis, Adelheid W Onyango, Jan Van den Broeck, W C Chumlea, and Reynaldo Martorell. Measurement and standardization protocols for anthropometry used in the construction of a new international growth reference. *Food and Nutrition Bulletin*, 25(supplement 1), 2004.
- Ernest Guevarra. *ppitables: Lookup Tables to Generate Poverty Likelihoods and Rates using the Poverty Probability Index (PPI)*, 2019a. URL <https://CRAN.R-project.org/package=ppitables>. R package version 0.5.1.
- Ernest Guevarra. *nutricheckr: Tools for Nutrition Assessment and Analysis*, 2019b. R package version 0.1.0.
- Ernest Guevarra and Abdoul-Aziz Goza. *myanmarMCCTdata: Myanmar Mother and Child Cash Transfer (MCCT) Programme Evaluation Data Handler*, 2019. URL <https://github.com/validmeasures/myanmarMCCTdata>. R package version 0.1.0.
- Innovations for Poverty Action. Myanmar 2015 PPI User Guide. Technical Report August, 2019.
- L Cheikh Ismail, H E Knight, E O Ohuma, L Hoch, and W C Chumlea. Anthropometric standardisation and quality control protocols for the construction of new , international , fetal and newborn growth standards : the INTERGROWTH-21 st Project. pages 48–55, 2013. doi: 10.1111/1471-0528.12127.
- A Kelly, J Kevany, M deOnis, and P M Shah. A WHO collaborative study of maternal anthropometry and pregnancy outcomes (Reprinted from International Journal of Gynecology and Obstetrics, vol 53, pg 219-233, 1996). *International journal of gynaecology and obstetrics: the official organ of the International Federation of Gynaecology and Obstetrics*, 57(1):1–15, April 1997.
- Daniel Maxwell and Richard Caldwell. *The Coping Strategies Index: A tool for rapid measurement of household food security and the impact of food aid in programs in humanitarian emergencies Field Methods Manual*. CARE USA, January 2008.
- Ministry of Health and Sports - MoHS/Myanmar and ICF. Myanmar Demographic and Health Survey 2015-16, 2017. URL <http://dhsprogram.com/pubs/pdf/FR324/FR324.pdf>.
- Ministry of Social Welfare, Relief and Resettlement and Livelihoods and Food Security Trust Fund. Maternal and child cash transfer programme in Chin State (MCCT) Baseline Survey Report, 2018.
- Mark Myatt. *nipnTK: NiPN Anthropometric Data Toolkit*, 2019. URL <https://github.com/nutriverse/nipnTK>. R package version 0.1.0.

- Mark Myatt and Ernest Guevarra. *zscorer: Child Anthropometry z-Score Calculator*, 2019. URL <https://CRAN.R-project.org/package=zscorer>. R package version 0.3.0.
- R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2019. URL <https://www.R-project.org/>.
- Peter Z Schochet. Statistical Power for Regression Discontinuity Designs in Education Evaluations. 34(2): 238–266, 2009. doi: 10.3102/1076998609332748. URL <https://www.jstor.org/stable/40263528>.
- Stanley J. Ulijaszek and Deborah A. Kerr. Anthropometric measurement error and the assessment of nutritional status. *British Journal of Nutrition*, 82(3):165–177, 1999. doi: 10.1017/S0007114599001348.
- Mija-tesse Ververs, Annick Antierens, Anita Sackl, Nelly Staderini, and Valerie Captier. Which Anthropometric Indicators Identify a Pregnant Woman as Acutely Malnourished and Predict Adverse Birth Outcomes in the Humanitarian Context? *PLoS Currents*, 2013.
- Vulnerability Assessment and Mapping World Food Programme. Food consumption analysis: Calculation and use of the food consumption score in food security analysis. Technical report, February 2008.
- WHO/UNICEF Joint Monitoring Programme for Water Supply and Sanitation. Core questions on water, sanitation and hygiene for household surveys 2018 Update. Technical report, Geneva, 2018.
- Willem Woertman, Esther de Hoop, Mirjam Moerbeek, Sytse U Zuidema, Debby L Gerritsen, and Steven Teerenstra. Use of a simple anthropometric measurement to predict birth weight. WHO Collaborative Study of Birth Weight Surrogates. *Bulletin of the World Health Organization*, 71(2):157–163, 1993.
- World Food Programme. Food Consumption Score Nutritional Quality Analysis (FCS-N). Technical report, Rome, August 2015.
- World Health Organization. Core Questions on Drinking-water and Sanitation for Household Surveys. World Health Organization, Geneva, May 2006.
- World Health Organization. *Indicators for assessing infant and young child feeding practices: conclusions of a consensus meeting held 6–8 November 2007*. World Health Organization, Washington, DC, 2008.
- World Health Organization (WHO). *WHO Child Growth Standards: Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: Methods and development*. World Health Organization, Geneva, 2006. ISBN 9789241547185. doi: 10.1037/e569412006-008. URL <http://hps.kbsplit.hr/hps-2008/pdf/doko3.pdf>.
- Yihui Xie, J.J. Allaire, and Garrett Grolemond. *R Markdown: The Definitive Guide*. Chapman and Hall/CRC, Boca Raton, Florida, 2018. URL <https://bookdown.org/yihui/rmarkdown>. ISBN 9781138359338.