

# intro\_project

April 8, 2022

```
[1]: # Using ML to classify LAEs in the NEP Field
import tables as tb
import numpy as np

import matplotlib.pyplot as plt
from matplotlib.colors import LogNorm

from astropy import constants as const
from astropy.table import Table, column, join
from astropy.io import fits
import astropy.units as u
from astropy.coordinates import SkyCoord
from astropy.visualization import ZScaleInterval

from regions import CircleSkyRegion, CirclePixelRegion

from hetdex_api.survey import Survey, FiberIndex
from hetdex_api.config import HDRconfig
from hetdex_api.detections import Detections
from hetdex_api.elixer_widget_cls import ElixerWidget

from hetdex_tools.get_spec import get_spectra

import pandas as pd
import seaborn as sb
```

```
[2]: det_object = Detections('hdr2.1', loadtable = False)
```

```
[3]: # Once it has loaded you want to filter out the data by selecting those that
      ↪are in the NEP field
      # to do this I will give you the verticies of a box that will encompass all the
      ↪NEP field - Oscar

      # The center of the NEP field is given by:
      # NEP Central Coordinates:
      # R.A. = 18hours00minutes00seconds, decl. = 66 degree 33minute 38.552 arcmin
```

```

# Then make a radius of 3.5 degrees centered above and find all the RA and DEC
↪coordinates
# in the DF that are within this circle

# creating the circle region in the sky (NEP field)
ra = '18h00m00s'
dec = '+66d33m38.552s'
center_sky_coords = SkyCoord(ra, dec, frame = 'icrs')

maskregion = det_object.query_by_coords(center_sky_coords, 3.5 * u.deg)
detects_in_NEP = det_object[maskregion]      # Sources within the NEP footprint

print('det_object: ', end = "")
print(np.size(detects_in_NEP.detectid))

```

det\_object: 69799

```

[ ]: ra = detects_in_NEP.ra * u.deg
     dec = detects_in_NEP.dec * u.deg
     input_coords = SkyCoord(ra, dec)

```

```

[ ]: sources = get_spectra(input_coords[10])

```

```

[ ]: sources

```

```

[ ]: det_object.__dict__.keys()

```

```

[ ]: print(det_object.version)

```

```

[ ]: # Once you have selected sources within the NEP footprint we can then go ahead
     ↪and find some
     # spectra from these sources - Oscar
     spectra = detects_in_NEP.hdf.file.root.Spectra

```

center\_sky.separations(skycoords entire) return indices return distance return 3d

separate by dist mask

main goal of algorithm want it to distinguish lae vs o2 emitter. wouldn't impose cuts unless training.

cut = filter cut = signal to noise could do plya

might need cuts for training for confident lya and o2

increase confidence by visually inspecting

could visually inspect to increase confidence.

plotting histograms to look for outliers

hetdex isn't perfect and it catches emission lines that aren't real. visual inspections helps

no need for dataframes if i found another way

save as csv with astropy table

csv into get\_spec()

has nice documentation

get\_spectrum good for ids \*\*\*\*\*USE\*\*\* returns all fiber spec with corresponding weights.

try to see if can get LAE samples. 02 samples. and ambiguous samples. Clasify some of them. Sig

detection object filter by fields. turn to astropy table and then filter.

Want psf weighted.

[ ]: