

Graduate Seminar

An introduction to stochastic calculus with applications to finance and SPDEs

Nicholas Eisenberg

March 29, 2022

Calculus

In undergraduate Calculus, we focus only on real valued measurable on $\mathbb{R}^d$:

$$f : D \subseteq \mathbb{R} \rightarrow \mathbb{R},$$

and the measure space that is implied is the following:

$$(\mathbb{R}, \mathcal{B}(\mathbb{R}), \lambda),$$

where λ is Lebesgue measure. Recall the average value of f on (a, b) :

$$\frac{1}{\lambda[(a, b)]} \int_a^b f(x) \, dx. = \frac{1}{b - a} \int_a^b f(x) \, dx.$$

Stochastic Calculus

When studying Probability Theory, we consider measurable functions (random variables),

$$X : \Omega \rightarrow \mathbb{R},$$

and we consider the measure spaces

$$(\Omega, \mathcal{F}, \mathbb{P}) \quad \text{and} \quad (\mathbb{R}, \mathcal{B}(\mathbb{R}), \lambda),$$

where Ω and $\mathcal{F}$ are referred to as the sample space and set of possible events and

$$\mathbb{P}(\Omega) = 1.$$

The average or expected value of X is

$$\mathbb{E}(X) = \int_{\Omega} X(\omega) \, d\mathbb{P}(\omega) = \int_{\mathbb{R}} x \, dF(x) = \int_{\mathbb{R}} xf(x) \, dx.$$

Stochastic Processes

Definition (Stochastic Process)

A stochastic process is a random variable that is indexed by set, T , which usually represents time:

$$X_t(\omega), \quad t \in T, \omega \in \Omega.$$

One can think a stochastic processes as a two parameter function

$$X_t(\omega) = X(t, \omega) : T \times \Omega \rightarrow \mathbb{R}.$$


Definition (Sample Path)

The variable ω represents the idea of a single possible outcome. The following function is then referred to as a *sample path*:

$$t \mapsto X_t(\omega).$$


Random Walk Example

2 sample paths of a random walk
jump size = 1.0 and step size = 1.0
number of steps = 5




Random Walk Example

2 sample paths of a random walk
jump size = 0.707 and step size = 0.5
number of steps = 10




Random Walk Example

2 sample paths of a random walk
jump size = 0.5 and step size = 0.25
number of steps = 20




Random Walk Example

2 sample paths of a random walk
jump size = 0.354 and step size = 0.12
number of steps = 40



Random Walk Example



Brownian Motion.

Theorem

Consider a random walk $X = \{X_0 = 0, X_{t_1}, \dots, X_{t_n}\}$. Suppose that for $k \in \{1, \dots, n\}$ that

$$\mathbb{P}[X_{t_k} = -h] = \mathbb{P}[X_{t_k} = h] = 1/2.$$

and suppose that the time increments are all constant:

$$t_k - t_{k-1} = d \quad \forall k \in \{1, \dots, n\}.$$

Suppose that $h^2 = d$. Now define

$$B_n(t_k) = \sum_{i=0}^k X_{t_i}.$$

Then B_n converges to a Brownian motion.

The Ito integral (special case)

Suppose that f is a nice continuous function. Then we can define:

$$\int_0^t f(s)dB_s(\omega) := \lim_{\|P\| \rightarrow 0} \sum_{i \in P} f(t_{i-1})(B_{t_i}(\omega) - B_{t_{i-1}}(\omega)),$$


where $P = t_0 = 0, t_1, t_2, \dots, t_n = t$ is a partition of $[0, t]$. In other words, the Ito integral in this case the left point Riemann sum.

In general, this formula is not true because the increments

$$dB_{t_i}(\omega) := B_{t_i}(\omega) - B_{t_{i-1}}(\omega)$$


are very bad.

White Noise



Increments of a smooth function for reference

Increments, $df(x_i) = f(x_i) - f(x_{i-1})$, for $f(x) = \sin(x^2)$



A 'Fundamental Theorem of Ito Calculus'

If f is twice continuously differentiable then we have the following "Fundamental Theorem of Ito Calculus"


$$\int_a^t \frac{d}{ds} f(B_s) dB_s = f(B_t) - f(B_a) - \frac{1}{2} \int_a^t \frac{d^2}{ds^2} f(B_s) ds.$$

Example

If we apply the above to $f(x) = x^2/2$ with $a = 0$ then we have

$$\int_0^t B_s dB_s = \frac{B_t^2}{2} - \frac{t}{2}.$$

Two sample paths



One more example

For a nice function $f(t, x)$ we can generalize Itos formula to

$$f(t, B_t) - f(a, B_a) = \int_a^t \frac{\partial f}{\partial x}(s, B_s) dB_s + \int_a^t \left(\frac{\partial f}{\partial t}(s, B_s) + \frac{\partial^2 f}{\partial x^2}(s, B_s) \right) ds.$$

Example


Now consider the function $f(t, x) = tx$. Then

$$\frac{\partial f}{\partial t}(t, x) = x, \quad \frac{\partial f}{\partial x}(t, x) = t, \quad \frac{\partial^2 f}{\partial x^2}(t, x) = 0$$

Plugging this in above for $a = 0$ gives us

$$tB_t = \int_0^t s \, dB_s + \int_0^t B_s \, ds, \quad \implies \quad \int_0^t B_s \, ds = tB_t - \int_0^t s \, dB_s.$$

Smoothness?!



Why do we care about Brownian Motion?



An application to finance.

Consider a stock that has value S_t at time t . Consider $t = 0$ to be the point of entry into the investment. We can define a return rate as follows:

$$S_t = S_0(1 + R_t), \quad R_t := \frac{S_t - S_0}{S_0} \text{ is the return rate over the period } [0, t].$$

We want to make predictions of our return. In a very simple model, we can assume that

$$R_t \sim \text{Normal}(\mu_t, \sigma^2), \quad \mu \in \mathbb{R}, \sigma \in (0, \infty).$$

In terms of Stochastic Calculus, we express this as the following stochastic differential equation:

$$R_t = \frac{dS_t}{S_t} = \mu dt + \sigma dB_t, \quad \text{where } B_t \text{ is a Brownian motion.}$$

Modeling a stock price with geometric brownian motion

The notation

$$dS_t = \mu S_t dt + \sigma S_t dB_t, \quad \text{where } B_t \text{ is a Brownian motion}$$


is only symbolic as dB_t does not exist. The above can be legitimately viewed as the following integral equation:

$$S_t - S_0 = \int_0^t \mu S_k dk + \int_0^t \sigma S_k dB_k.$$

The stochastic processes S_t that solves the above equation is called a Geometric Brownian Motion. Notice that we have come up with a random variable that models the stock price:

$$S_t(\omega) = S_0 + \int_0^t \mu S_k(\omega) dk + \int_0^t \sigma S_k(\omega) dB_k(\omega).$$

Geometric Brownian Motion: $dX_t = \sigma X_t dW_t + \mu X_t dt$
 $(a, b) = (0, 1)$, $X_a = 5$, $\sigma = 0.01$, and $\mu = 0.001$
we subdivide the domain $n = 500$ times



Black-Scholes option pricing model for European calls

Suppose at time $t = 0$ you buy a contract from a seller, allowing you to purchase 1 share of a stock at time $t = T$ and a set price of K , regardless if what the stock price at time T is.

$$\begin{aligned}\text{Payout and time } T &= \begin{cases} S_T - K & \text{if } S_T > k \\ 0 & \text{otherwise} \end{cases} \\ &= (S_T - K)^+\end{aligned}$$

What is a fair price for the this contract?

For this we need a to create a self-financing strategy.

We assume that the stocks price follows a geometric Brownian motion,

$$dS_t = \mu S_t dt + \sigma S_t dB_t.$$

Let a_{t_i} be the amount of shares owned over the period $t \in (t_{i-1}, t_i]$. Then at any time $t \in [0, T]$, the amount of shares owned is

$$a_t = \sum_{i=1}^n a_{t_i} \mathbf{1}_{(t_{i-1}, t_i]}(t)$$

and the realized over $[0, T]$

$$\begin{aligned} a_T S_T - a_0 S_0 &= \sum_{i=1}^n a_{t_i} (S_{t_i} - S_{t_{i-1}}) = \int_0^T a_t dS_t \\ &= \mu \int_0^T a_t S_t dt + \sigma \int_0^T a_t S_t dB_t \end{aligned}$$

Summarizing the last slide...

$$\text{Our stock profit on time } [0, T] = \underbrace{a_T S_T - a_0 S_0 = \int_0^T a_t \, dS_t}$$

Now suppose we also invest in a bond with interest rate, r , that has price β_t at time t . Similarly...

$$\text{Our bond profit on time } [0, T] = \underbrace{b_T S_T - b_0 S_0 = \int_0^T b_t \, d\beta_t}$$

Definition (Self Financing Strategy)

A pair of stocks and bonds, (a, b) , is self financing if

$$a_T S_T + b_T S_T = a_0 S_0 + b_0 S_0 + \int_0^T a_t \, dS_t + \int_0^T b_t \, d\beta_t$$

Fair price for a European call option

A fair price for a European call option that allows the buyer to buy 1 share at price K at time T regardless of the price of the stock S_T at time T is the amount one would need to invest in a self financing strategy,

$$\underbrace{a_T S_T + b_T S_T}_{V_T} = \underbrace{a_0 S_0 + b_0 S_0}_{V_0} + \int_0^T a_t dS_t + \int_0^T b_t d\beta_t,$$

such that the value of the strategy at time T is equal to $(S_T - K)^+$:

$$V_T = (S_T - K)^+$$

Let $f(S_t, T - t) = V_t$. So we see that

$$f(S_T, 0) = V_T = (S_T - K)^+ \quad \text{and} \quad f(S_0, T) = V_0$$

So we need to find f . Moreover, $f(S_0, T)$ is the fair price to pay.

It can be shown that f solves the following partial differential equation

$$\begin{cases} \frac{\partial f}{\partial s}(s, x) = \frac{1}{2}\sigma^2 x^2 \frac{\partial^2 f}{\partial x^2}(s, x) + rx \frac{\partial f}{\partial x}(s, x) - rf(s, x) & (s, x) \in [0, T] \times (0, \infty) \\ f(0, x) = (x - K)^+ & x \in (0, \infty). \end{cases}$$

The solution is given by

$$f(t, x) = x\Phi(g(t, x)) - Ke^{-rt}\Phi(h(t, x)),$$

where

$$g(t, x) = \left[\ln\left(\frac{x}{K}\right) + \left(r + \frac{1}{2}\sigma^2\right)t \right] / (\sigma\sqrt{t}),$$

and

$$h(t, x) = g(t, x) - \sigma/\sqrt{t},$$

and

$$\Phi(t) = \frac{1}{\sqrt{2\pi}} \int_0^t \exp\left(-\frac{|x|^2}{2}\right) dx$$

Potential for unlimited profit

In practice, option contracts are priced by "the market". This means that the cost of a contract is literally what the buyer is willing to pay or what the seller is willing to sell at.

This gives the educated buyer (or seller) a chance to make profit.


Suppose the fair price for a call contract with strike K is $\$q$ but we find a buyer, who does not know about the Black Scholes model, who wants to buy the contract for $\$p > \q . We can do the following:

- ▶ sell a contract for $\$p$,
- ▶ invest $\$q$ in stock and bond at time $t = 0$ in a self financing strategy (HARD TO DO),
- ▶ keep the remaining $p - q$ dollars.
- ▶ pay the buyer $(S_T - K)^+$ at expiration.

No matter what, we make $p - q$ dollars. If we sell 100 contracts then we make $100(p - q)$...


Sample path of a Brownian Sheet, $W(t, x)$

Brownian Sheet




Sample path of a Brownian Sheet, $W(t, x)$

Brownian Sheet



Sample path of a Brownian Sheet, $W(t, x)$

Brownian Sheet



Stochastic Heat Equation

$$\begin{cases} \frac{\partial}{\partial t} u(t, x) - \frac{\partial^2}{\partial x^2} u(t, x) = \lambda u(t, x) \frac{\partial^2}{\partial t \partial x} W(t, x) & t \geq 0, x \in \mathbb{R} \\ u(0, x) = f(x) & x \in \mathbb{R} \end{cases}$$

The solution to the stochastic heat equation is given through the following integral equation:

$$u(t, x) = \int_{\mathbb{R}} G(t, x - y) f(y) dy + \lambda \int_0^t \int_{\mathbb{R}} G(t - s, x - y) u(s, y) W(ds, dy),$$


where

$$G(t, x) = \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{|x|^2}{2t}\right)$$

A finite difference method simulation of

$$u_t(t, x) - \frac{1}{2}u_{xx}(t, x) = 0u(t, x)W(t, x)$$


$$u(x, 0) = \delta_0(x)$$



A finite difference method simulation of

$$u_t(t, x) - \frac{1}{2}u_{xx}(t, x) = 1u(t, x)W(t, x)$$


$$u(x, 0) = \delta_0(x)$$



A finite difference method simulation of

$$u_t(t, x) - \frac{1}{2}u_{xx}(t, x) = 2u(t, x)W(t, x)$$


$$u(x, 0) = \delta_0(x)$$



A finite difference method simulation of

$$u_t(t, x) - \frac{1}{2}u_{xx}(t, x) = 2.15u(t, x)\dot{W}(t, x)$$

$$u(x, 0) = \delta_0(x)$$



Growth of Moments

We have a quantity that describes the growth of the p^{th} moments of the solutions to the stochastic heat question:

$$\lim_{t \rightarrow \infty} \frac{\log [\mathbb{E}(|u(t, x)|^p)]}{t}.$$

What does this limit say about the growth or decay of the moments?

Growth of Moments for the Stochastic Heat Equation

$$\begin{cases} \frac{\partial}{\partial t} u(t, x) - \frac{1}{2} \frac{\partial^2}{\partial x^2} u(t, x) = \sqrt{\lambda} u(t, x) \frac{\partial}{\partial x} W(x) & t \geq 0, x \in \mathbb{R} \\ u(0, x) = 1 & x \in \mathbb{R} \end{cases}$$

Theorem (Hu 2002)

For the above equation, we can prove that

$$\lim_{t \rightarrow \infty} \frac{\log [\mathbb{E}(|u(t, x)|^p)]}{t^3} = \frac{p(p-1)\lambda^2}{24}$$

which means that

$$t \mapsto \mathbb{E}(|u(t, x)|^p) = O \left[\exp \left(\frac{p(p-1)\lambda^2}{24} t^3 \right) \right], \quad \text{as } t \rightarrow \infty.$$

So the moments grow exponentially in time.

References

 K. Chung and R. Williams (1990)

Introduction to Stochastic Integration

Probability and Its Applications, Birkhäuser

 Hu, Y. 2002

Chaos expansion of heat equations with white noise potentials.

Potential Analysis **16**, no. 1, 45-66.

 H. Kuo (2000)

Introduction to Stochastic Integration

Universitext, Springer