

1 Questions

Nick's Question 1.1
In the Wave equation paper you showed that for the Greens function, we have that

$$\mathcal{F}G(t, \cdot)(\xi) = \frac{\sin(t|\xi|)}{|\xi|}$$

and so we see that

$$\mathcal{F}G(t, \cdot)(c\xi) = \frac{1}{c} \mathcal{F}(ct, \cdot)(\xi)$$

However in the Fractional SPDE paper the greens function is more complicated and we have that

$$\begin{aligned} \mathcal{F}Y_{\alpha, \beta, \gamma, d}(t, \cdot)(\xi) &= t^{\beta+\gamma-1} E_{\beta, \beta+\gamma}(-2^{-1} \nu t^\beta |\xi|^\alpha) \\ &= t^{\beta+\gamma-1} H_{1,2}^{1,1} \left(2^{-1} \nu t^\beta |\xi|^\alpha \middle| \begin{matrix} (0, 1) \\ (0, 1) \end{matrix} \right. (1 - \beta - \gamma, \beta) \Big) \\ &= t^{\beta+\gamma-1} \frac{1}{2\pi i} \int_L \mathcal{H}_1^{1,1}(s) \left(2^{-1} \nu t^\beta |\xi|^\alpha \right)^{-s} ds \end{aligned}$$

And so we see that

$$\mathcal{F}Y_{\alpha, \beta, \gamma, d}(t, \cdot)(c\xi) = t^{\beta+\gamma-1} \frac{1}{2\pi i} \int_L \mathcal{H}_1^{1,1}(s) \left(2^{-1} \nu t^\beta |c\xi|^\alpha \right)^{-s} ds$$

However I am unsure how to handle the constant, c , on the inside so we can get a nice scaling property like we do in the Wave Equation paper.

Also, instead of using the Fox-H function we can just say

$$\begin{aligned} \mathcal{F}Y_{\alpha, \beta, \gamma, d}(t, \cdot)(\xi) &= t^{\beta+\gamma-1} E_{\beta, \beta+\gamma}(-2^{-1} \nu t^\beta |\xi|^\alpha) \\ &= t^{\beta+\gamma-1} \sum_{k=0}^{\infty} \frac{(-2^{-1} \nu t^\beta |\xi|^\alpha)^k}{\Gamma(\beta k + (\beta + \gamma))} \end{aligned}$$

and so

$$\mathcal{F}Y_{\alpha, \beta, \gamma, d}(t, \cdot)(c\xi) = t^{\beta+\gamma-1} \sum_{k=0}^{\infty} \frac{(-2^{-1} \nu t^\beta |c\xi|^\alpha)^k}{\Gamma(\beta k + (\beta + \gamma))}$$

but I still don't see how to handle the c .

Here is how one should proceed:

$$\begin{aligned} \mathcal{F}Y_{\alpha, \beta, \gamma, d}(t, \cdot)(c\xi) &= t^{\beta+\gamma-1} \sum_{k=0}^{\infty} \frac{(-2^{-1} \nu t^\beta |c\xi|^\alpha)^k}{\Gamma(\beta k + (\beta + \gamma))} \\ &= t^{\beta+\gamma-1} \sum_{k=0}^{\infty} \frac{(-2^{-1} \nu t^\beta c^\alpha |\xi|^\alpha)^k}{\Gamma(\beta k + (\beta + \gamma))} \\ &= t^{\beta+\gamma-1} \sum_{k=0}^{\infty} \frac{(-2^{-1} \nu (tc^{\frac{\alpha}{\beta}})^\beta |\xi|^\alpha)^k}{\Gamma(\beta k + (\beta + \gamma))} \\ &= c^{\frac{\alpha}{\beta}(1-\gamma-\beta)} (tc^{\frac{\alpha}{\beta}})^{\beta+\gamma-1} \sum_{k=0}^{\infty} \frac{(-2^{-1} \nu (tc^{\frac{\alpha}{\beta}})^\beta |\xi|^\alpha)^k}{\Gamma(\beta k + (\beta + \gamma))} \end{aligned}$$

$$= c^{\frac{\alpha}{\beta}(1-\gamma-\beta)} \mathcal{F}Y_{\alpha,\beta,\gamma,d} \left(c^{\frac{\alpha}{\beta}} t, \cdot \right) (c\xi)$$

Please check if this scaling reduces to heat and wave equations correctly.

Nick's Question 1.2 In the wave equation paper with Xia Chen you said that that fundamental solution to the SPDE

$$\begin{cases} \frac{\partial^2 u}{\partial t^2}(t, x) - \frac{1}{2} \Delta u(t, x) = \sqrt{\Theta} u(t, x) \dot{W}(x) & x \in \mathbb{R}^d, t > 0 \\ u(0, \cdot) = 1 \\ u_t(0, \cdot) = 0 \end{cases} \quad (1.1)$$

wspde

and the mild solution is defined to be

$$u(t, x) = 1 + \sqrt{\Theta} \int_0^t \left(\int_{\mathbb{R}^d} G(t-s, x-y) u(s, y) W(\delta y) \right) ds \quad (1.2)$$

weqs

but for general initial conditions for the wave equation we have the solution to be

$$\begin{aligned} u(t, x) = & \frac{\partial}{\partial t} \int_{\mathbb{R}^d} G(t-s, x-y) u(0, y) dy + \int_{\mathbb{R}^d} G(t-s, x-y) u_t(0, y) dy \\ & + \sqrt{\Theta} \int_0^t \left(\int_{\mathbb{R}^d} G(t-s, x-y) u(s, y) W(\delta y) \right) ds \end{aligned} \quad (1.3)$$

gweqs

But when the initial conditions are such as in 1.1 then the first two integral in 1.3 are 1 and 0 respectively and so we arrive at 1.2.

In the fractional SPDE

$$\begin{cases} (\partial^\beta + \frac{\nu}{2}(-\Delta)^{\alpha/2}) u(t, x) = I_t^\gamma \left[\rho(u(t, x)) \dot{W}(x) \right] & x \in \mathbb{R}^d, t > 0 \\ u(0, \cdot) = \mu & \beta \in (0, 1] \\ u(0, \cdot) = \mu_0, \quad u_t(0, \cdot) = \mu_1 & \beta \in (1, 2] \end{cases} \quad (1.4)$$

fspde

and we define

$$J_0(t, x) = \begin{cases} (Z(t, \cdot) * \mu)(x) & \beta \in (0, 1] \\ (Z^*(t, \cdot) * \mu_0)(x) + (Z(t, \cdot) * \mu_1)(x) & \beta \in (1, 2] \end{cases}$$

and

$$I(t, x) = \iint_{[0,t] \times \mathbb{R}} Y(t-s, x-y) \rho(u(s, y)) W(\delta y) ds$$

and the general solution for 1.4 is

$$u(t, x) = J_0(t, x) + I(t, x) \quad (1.5)$$

feqs

My question is that if we consider the following fractional spde

$$\begin{cases} (\partial^\beta + \frac{\nu}{2}(-\Delta)^{\alpha/2}) u(t, x) = I_t^\gamma \left[\rho(u(t, x)) \dot{W}(x) \right] & x \in \mathbb{R}^d, t > 0 \\ u(0, \cdot) = 1 & \beta \in (0, 1] \\ u(0, \cdot) = 1, \quad u_t(0, \cdot) = 0 & \beta \in (1, 2] \end{cases} \quad (1.6)$$

fspde1

then can we calculate $J_0(t, x)$? More specifically is $J_0(t, x) = 1$ in this case? The wave equation paper worked out nicely because in that paper $J_0(t, x) = 1$ and so the solution was able to be written as 1.2. The 1 allows us to easily do the Piccard iteration to get equation (2.3) from the Wave equation paper with Xia Chen.

So if our solution can not be written as

$$u(t, x) = 1 + I(t, x)$$

I am not sure at the moment how we can come up with an expression for the f'_n s in the Weiner Expansion like you did in equation (2.3) in the paper with Xia Chen.

Hi, Nick, here is the answer:

Notice that, by Theorem 4.1 of my paper,

$$\begin{aligned} (Z_{\alpha, \beta, \gamma, d}(t, \cdot) * 1)(x) &= \mathcal{F}Z_{\alpha, \beta, \gamma, d}(t, \cdot)(0) \\ &= t^{\lceil \beta \rceil - 1} E_{\beta, \lceil \beta \rceil}(0) \\ &= \frac{t^{\lceil \beta \rceil - 1}}{\Gamma(\lceil \beta \rceil)}. \end{aligned}$$

Hence, when $\beta \in (0, 1]$, we see that $(Z_{\alpha, \beta, \gamma, d}(t, \cdot) * 1)(x) = 1$. When $\beta \in (1, 2]$,

$$\begin{aligned} (Z_{\alpha, \beta, \gamma, d}^*(t, \cdot) * 1)(x) &= \mathcal{F}Z_{\alpha, \beta, \gamma, d}^*(t, \cdot)(0) \\ &= E_{\beta}(0) \\ &= \frac{1}{\Gamma(1)} = 1. \end{aligned}$$

Hence, under the initial conditions given by (1.6), $J_0(t, x) \equiv 1$.

Nick's Question 1.3 Guess for the constant (1.20) from the Wave Equation Paper In the wave equation paper, the constant (1.20) was

$$C = \begin{cases} \frac{4 - \Delta}{3 - \Delta} & \text{under Assumption A} \\ \frac{4 - d}{3 - d} & \text{under Assumption B or C} \end{cases} \quad (1.7) \quad \boxed{\text{const}}$$

I believe in the fractional paper the constant will be

$$\begin{cases} \frac{2(\beta + \gamma - 1) + 2 - \beta\Delta/\alpha}{2(\beta + \gamma - 1) + 1 - \beta\Delta/\alpha} & \text{under Assumption A} \\ \frac{2(\beta + \gamma - 1) + 2 - \beta d/\alpha}{2(\beta + \gamma - 1) + 1 - \beta d/\alpha} & \text{under Assumption B or C} \end{cases}$$

I think this because in the wave equation paper where we use $(4 - \Delta)$ this gets replaced by $(2(\beta + \gamma - 1) + 2 - \beta\Delta/\alpha)$ and thus we can guess that

$$\begin{cases} 4 - \Delta & \rightarrow & 2(\beta + \gamma - 1) + 2 - \beta\Delta/\alpha \\ 3 - \Delta & \rightarrow & 2(\beta + \gamma - 1) + 1 - \beta\Delta/\alpha \end{cases}$$

Note that when $\alpha = \beta = 2$ and when $\gamma = 0$ then the guess reduces to the wave equation.

Nick's Question 1.4 Lemma 2.3 for the fractional equation

1. In the outline you sent me, you use

$$\mathcal{F}_a = \left\{ g \in H^a(\mathbb{R}^d) : \|g\|_{L^2(\mathbb{R}^d)} = 1, \mathcal{E}_a(g, g) < \infty \right\}$$

Where $H^a(\mathbb{R}^d)$ is the fractional Sobelov Space where $H^2(\mathbb{R}^d) = W^{1,2}(\mathbb{R}^d)$. However, I saw on line that $H^a(\mathbb{R}^d) = W^{a,2}(\mathbb{R}^d)$ and so $H^2(\mathbb{R}^d) = W^{2,2}(\mathbb{R}^d)$. Is there a mistake here?

2. Where is $\mathcal{M}$ defined with the sup over $\mathcal{F}_d$ and not $\mathcal{F}_a$? Is this a typo?
3. When defining $\mathcal{E}_a(g, g)$ why does the $(2\pi)^{-d}$ appear here but not in the wave paper?
4. How do we generalize ρ from the wave paper to the fractional equation? For the wave equation we have that

$$\rho = (\mathcal{M}(\gamma, 2))^{(4-\alpha)/2}$$

but why does it generalize to this in the fractional sense

$$\rho = (\mathcal{M}_a(\gamma, 2))^{(2a-\alpha)/a} = 2^{-a/\alpha} (\mathcal{M}_a(\gamma, 1))^{(2a-\alpha)/a}$$

which is what you put in the outline you sent me.

2 Proof Sketches

Nick's Question 2.1 A mild solution for

$$\begin{cases} (\partial^\beta + \frac{\nu}{2}(-\Delta)^{\alpha/2}) u(t, x) = I_t^\gamma [\rho(u(t, x)) \dot{W}(x)] & x \in \mathbb{R}^d, t > 0 \\ u(0, \cdot) = 1 & \beta \in (0, 1] \\ u(0, \cdot) = 1, \quad u_t(0, \cdot) = 0 & \beta \in (1, 2] \end{cases}$$

must satisfy the following integral equation

$$u(t, x) = 1 + \int_0^t \left(\int_{\mathbb{R}^d} u(t-s, x-y) Y(s, y) W(\delta y) \right) ds$$

an through a Piccard Iteration Scheme we can see that the solution should be given by

$$u(t, x) = 1 + \sum_{k=1}^{\infty} \theta^{k/2} I_k(f_k(\cdot, x, t)) \quad (2.1) \quad \text{sol}$$

where

$$f_k(x_1, \dots, x_k, x, t) = \int_0^t \int_0^{t_k} \cdots \int_0^{t_2} Y(t-t_k, x-x_k) \cdots Y(t_2-t_1, x_2-x_1) dt_1 \cdots dt_n \quad (2.2) \quad \text{solker}$$

$$= \int_0^t \int_0^{t_k} \cdots \int_0^{t_2} Y(t_1, x-x_k) \cdots Y(t_k-t_{k-1}, x_2-x_1) dt_1 \cdots dt_n \quad (2.3)$$

and also

$$\mathcal{F}f_k(\cdot, x, t)(\xi_1, \dots, \xi_n) = e^{-i(\sum_{j=1}^n \xi_j) \cdot x} \int_{[0, t]^{n <}} \prod_{k=1}^n \overline{\mathcal{F}Y(t_{t_k+1} - t_k, \cdot)} \left(\sum_{j=1}^k \xi_j \right) dt \quad (2.4) \quad \boxed{\text{fsolker}}$$

where $t_{n+1} = t$

Nick's Question 2.2 Greens Function Scaling Property

For and $c > 0$ we have

$$\mathcal{F}Y(ct, \cdot)(\xi) = c^{b+r-1} \mathcal{F}Y(t, \cdot)(t^{b/a} \xi). \quad (2.5) \quad \boxed{\text{fscale}}$$

which is shown above.

Nick's Question 2.3 $\mathcal{H}^{\otimes n}$ Norm Scaling Property

Suppose that $\mu(d\xi) = \varphi(\xi)d\xi$ and φ satisfies $\varphi(c\xi) = c^{-(d-\alpha)}$ for some $\alpha \in (0, d)$ and for any $c > 0$. Also recall that

$$\mathcal{F}Y(ct, \cdot)(\xi) = c^{b+r-1} \mathcal{F}Y(t, \cdot)(t^{b/a} \xi).$$

Then

$$\left\| \tilde{f}_n(\cdot, 0, t) \right\|_{H^{\otimes n}}^2 = t^{[2(b+r)-b\alpha/a]n} \left\| \tilde{f}_n(\cdot, 0, 1) \right\|_{H^{\otimes n}}^2 \quad (2.6) \quad \boxed{\text{nscale}}$$

Proof:

$$\left\| \tilde{f}_n(\cdot, 0, t) \right\|_{H^{\otimes n}}^2 = \frac{1}{n!} \sum_{\rho \in \Sigma_n} \int_{\mathbb{R}^{nd}} \mathcal{F}f_n(\cdot, 0, t)^2(\vec{\xi}) \varphi(\vec{\xi}) d\vec{\xi}$$

where $\varphi(\vec{\xi}) = \varphi(\xi_1) \cdots \varphi(\xi_n)$

$$\begin{aligned} \int_{\mathbb{R}^{nd}} \mathcal{F}f_n(\cdot, 0, t)^2 \varphi(\vec{\xi}) d\vec{\xi} &= \int_{\mathbb{R}^{nd}} \left(\int_{[0, t]^{n <}} \overline{\mathcal{F}Y(t - t_n, \cdot)(\xi_1 + \cdots + \xi_n)} \cdots \overline{\mathcal{F}Y(t_2 - t_1, \cdot)(\xi_1)} \right)^2 \varphi(\vec{\xi}) d\vec{\xi} \\ &= t^{2n(b+r-1)} \int_{\mathbb{R}^{nd}} \left(\int_{[0, t]^{n <}} \overline{\mathcal{F}Y\left(1 - \frac{t_n}{t}, \cdot\right)(t^{b/a}(\xi_1 + \cdots + \xi_n))} \cdots \overline{\mathcal{F}Y\left(\frac{t_2}{t} - \frac{t_1}{t}, \cdot\right)(t^{b/a}\xi_1)} \right)^2 \varphi(\vec{\xi}) d\vec{\xi} \\ &= t^{2n(b+r-1)+2n} \int_{\mathbb{R}^{nd}} \left(\int_{[0, 1]^{n <}} \overline{\mathcal{F}Y(1 - t_n, \cdot)(t^{b/a}(\xi_1 + \cdots + \xi_n))} \cdots \overline{\mathcal{F}Y(t_2 - t_1, \cdot)(t^{b/a}\xi_1)} \right)^2 \varphi(\vec{\xi}) d\vec{\xi} \\ &= t^{2n(b+r-1)+2n} \int_{\mathbb{R}^{nd}} \left(\int_{[0, 1]^{n <}} \overline{\mathcal{F}Y(1 - t_n, \cdot)(\xi_1 + \cdots + \xi_n)} \cdots \overline{\mathcal{F}Y(t_2 - t_1, \cdot)(\xi_1)} \right)^2 \varphi(t^{-b/a}\vec{\xi}) t^{-ndb/a} d\vec{\xi} \\ &= t^{2n(b+r-1)+2n-nb\alpha/a} \int_{\mathbb{R}^{nd}} \left(\int_{[0, 1]^{n <}} \overline{\mathcal{F}Y(1 - t_n, \cdot)(\xi_1 + \cdots + \xi_n)} \cdots \overline{\mathcal{F}Y(t_2 - t_1, \cdot)(\xi_1)} \right)^2 \varphi(\vec{\xi}) d\vec{\xi} \\ &= t^{2n(b+r-1)+2n-nb\alpha/a} \int_{\mathbb{R}^{nd}} \mathcal{F}f_n(\cdot, 0, 1)(\vec{\xi})^2 \varphi(\vec{\xi}) d\vec{\xi} \\ &= t^{[2(b+r-1)+2-nb\alpha/a]n} \left\| \tilde{f}_n(\cdot, 0, 1) \right\|_{H^{\otimes n}}^2 \\ &= t^{[2(b+r)-b\alpha/a]n} \left\| \tilde{f}_n(\cdot, 0, 1) \right\|_{H^{\otimes n}}^2 \end{aligned}$$

Nick's Question 2.4 Lemma 2.3 We define

$$\mathcal{F}_a = \left\{ g \in H^a(\mathbb{R}^d) : \|g\|_{L^2(\mathbb{R}^d)} = 1, \mathcal{E}_a(g, g) < \infty \right\}$$

and then we also define

$$\mathcal{M}_a(f, \theta) = \sup_{g \in \mathcal{F}_a} \left\{ \langle g^2 * f, g^2 \rangle^{1/2} - \frac{\theta}{2} \int_{\mathbb{R}^d} |\xi|^a |\mathcal{F}g(\xi)|^2 d\xi \right\} \quad (2.7) \quad \boxed{\text{var}}$$

Now we define $\tilde{g}(x) = B^{-d/2}g(x/B)$. Its easy to show that $\mathcal{F}\tilde{g}(\xi) = B^{d/2}\mathcal{F}g(B\xi)$. By a change of variables, we can then show that

$$\int_{\mathbb{R}^d} |\xi|^a |\mathcal{F}\tilde{g}(\xi)|^2 d\xi = B^{-a} \int_{\mathbb{R}^d} |\xi|^a |\mathcal{F}g(\xi)|^2 d\xi$$

Next, by applying the scaling property of f , $f(cx) = c^{-\alpha}f(x)$, we can show by again using a change of variables that

$$\begin{aligned} (\tilde{g}^2 * f)(x) &= B^{-d} \int_{\mathbb{R}^d} \tilde{g}^2(y/B) f(x-y) dy \\ &= \int_{\mathbb{R}^d} g^2(u) f(x-uB) du \\ &= B^{-\alpha} \int_{\mathbb{R}^d} g^2(u) f\left(\frac{x}{B} - u\right) du \\ &= B^{-\alpha} (g^2 * f)(x/B) \end{aligned}$$

This can be used to show that

$$\langle \tilde{g}^2 * f, \tilde{g}^2 \rangle = B^{-\alpha} \langle g^2 * f, g^2 \rangle$$

Now we can rewrite (2.7) as the following

$$\mathcal{M}_a(\Theta f, \theta) = \frac{1}{\theta} \sup_{g \in \mathcal{F}_a} \left\{ \frac{\sqrt{\Theta}}{\theta} B^{\alpha/2} \langle \tilde{g}^2 * f, \tilde{g}^2 \rangle^{1/2} - \frac{B^a}{2} \int_{\mathbb{R}^d} |\xi|^a |\mathcal{F}\tilde{g}(\xi)|^2 d\xi \right\}$$

and as before, in order to preserve a scaling property, we want to choose B such that

$$\frac{\sqrt{\Theta}}{\theta} B^{\alpha/2} = B^a$$

and solving for B we get

$$B = \left(\frac{\sqrt{\Theta}}{\theta} \right)^{\frac{2}{2a-\alpha}}$$

and after plugging this B in we get that

$$\mathcal{M}_a(\Theta f, \theta) = \Theta^{\frac{a}{2a-\alpha}} \theta^{-\frac{\alpha}{2a-\alpha}} \mathcal{M}_a(f, 1)$$

Some definitions for solvability

$$L_n(y_1, \dots, y_n) = n! \int_0^\infty e^{-t} \tilde{f}_n(y_1, \dots, y_n, 0, t) dt$$

$$= \sum_{\sigma \in \Sigma_n} \int_0^\infty e^{-t} \int_{[0,t]^n} \prod_{k=1}^n Y(s_k - s_{k-1}, y_{\sigma(k)} - y_{\sigma(k-1)}) ds dt$$

where $s_0 = y_{\sigma(0)} = 0$.

$$\begin{aligned} H_n(t, \mathbf{x}) &= n! \int_{(\mathbb{R}^d)^n} \prod_{k=1}^n K(x_k - y_k) \tilde{f}_n(y_1, \dots, y_n, 0, t) d\mathbf{y} \\ &= \sum_{\sigma \in \Sigma_n} \int_{[0,t]^n} \int_{(\mathbb{R}^d)^n} \prod_{k=1}^n K(x_k - y_k) \prod_{k=1}^n Y(s_k - s_{k-1}, y_{\sigma(k)} - y_{\sigma(k-1)}) d\mathbf{y} ds \end{aligned}$$

$$L(x) = \int_0^\infty e^{-t} Y(t, x) dt \quad \text{and} \quad \mathcal{F}L(\xi) = \int_0^\infty e^{-t} \mathcal{F}Y(t, \cdot)(\xi) dt = \frac{1}{1 + \frac{\nu}{2} |\xi|^a}$$

$$L_n(y_1, \dots, y_n) = \sum_{\sigma \in \Sigma_n} L(y_{\sigma(1)}) L(y_{\sigma(2)} - y_{\sigma(1)}) \cdots L(y_{\sigma(n)} - y_{\sigma(n-1)})$$

$$\mathcal{F}L_n(\xi_1, \dots, \xi_n) = \sum_{\sigma \in \Sigma_n} \prod_{k=1}^n \frac{1}{1 + \frac{\nu}{2} \left| \sum_{j=k}^n \xi_j \right|^a}$$

$$T_n = \int_{(\mathbb{R}^d)^n} \left[\sum_{\sigma \in \Sigma_n} \prod_{k=1}^n \frac{1}{1 + \frac{\nu}{2} \left| \sum_{j=k}^n \xi_j \right|^a} \right] \mu(d\xi_1) \cdots \mu(d\xi_n)$$

$$C'_\mu = \int_{(\mathbb{R}^d)^n} \left(\frac{1}{1 + |\xi|^a} \right)^2 \mu(d\xi)$$

Nick's Question 2.5 Lemma 3.3, 3.4, and 3.5 Same proof as in wave paper. Have all details in my notebook.

Nick's Question 2.6 Lemma 3.6 We have that

$$C'_\mu < \infty \quad \text{and} \quad T_n < (n!)^2 (C'_\mu)^n$$

proof: By the same spherical substitution in the wave paper we see that

$$C'_\mu = \left(\int_{S^{d-1}} \varphi(\gamma) d\sigma(\gamma) \right) \left(\int_0^\infty \frac{\rho^{\alpha-1}}{(1 + \rho^a)^2} d\rho \right)$$

The first integral is shown to be finite in the wave paper. The second integral is finite if

$$2a - \alpha + 1 > 1 \tag{2.8} \quad \boxed{\alpha > 2}$$

Second inequality has the same proof as in the wave paper however I do not know why

$$(n!)^2 \int_{(\mathbb{R}^d)^n} \prod_{k=1}^n \left(\frac{1}{1 + \left| \sum_{j=k}^n \xi_j \right|^2} \right)^2 \mu(d\xi_1) \cdots \mu(d\xi_n) \leq (n!)^2 \left[\sup_{\eta \in \mathbb{R}^d} \int_{(\mathbb{R}^d)^n} \prod_{k=1}^n \left(\frac{1}{1 + |\xi + \eta|^2} \right)^2 \mu(d\xi) \right]^n$$

Nick's Question 2.7 Lemma 3.7

$$\int_0^\infty e^{-t} \left\| \tilde{f}_n(\cdot, 0, t) \right\|_{\mathcal{H}^{\otimes n}}^n dt \leq \frac{2^{n[2(b+r)-\frac{b\alpha}{a}]} T_n}{(n!)^2} \leq 2^{n[2(b+r)-\frac{b\alpha}{a}]} (C'_\mu)^n$$

Proof: Details in my notebook

Nick's Question 2.8 Theorem 3.1 1.6 has a unique solution $\{u(t, x) : t > 0, x \in \mathbb{R}^d\}$ and

$$\sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \mathbb{E}(|u(t, x)|^p) < \infty \quad \forall p \geq 2 \text{ and } T > 0.$$

Proof: First using 2.5 (details in notebook) we can show that

$$\mathcal{F} \tilde{f}_n(\cdot, 0, ct)(\xi_1, \dots, \xi_n) = c^{n(b+r)} \mathcal{F} \tilde{f}_n(\cdot, 0, t)(c^{b/a} \xi_1, \dots, c^{b/a} \xi_n). \quad (2.9)$$

In order to prove the theorem we need to show that $\sum_{n \geq 0} \theta^n n! \left\| \tilde{f}_n(\cdot, 0, t) \right\|_{\mathcal{H}^{\otimes n}}^2 < \infty$. From the scaling property 2.6 we see that

$$\begin{aligned} \int_0^\infty e^{-t} \left\| \tilde{f}_n(\cdot, 0, t) \right\|_{\mathcal{H}^{\otimes n}}^2 dt &= \left\| \tilde{f}_n(\cdot, 0, 1) \right\|_{\mathcal{H}^{\otimes n}}^2 \int_0^\infty e^{-t} t^{2(b+r)-\frac{b\alpha}{a}} n dt \\ &= \left\| \tilde{f}_n(\cdot, 0, 1) \right\|_{\mathcal{H}^{\otimes n}}^2 \Gamma \left(\left[2(b+r) - \frac{b\alpha}{a} \right] n + 1 \right) \end{aligned}$$

as long as we have

$$2(b+r) - \frac{b\alpha}{a} > 0 \quad (2.10) \quad \boxed{\text{alpha3}}$$

By using Lemma 3.7 we can deduce that

$$\begin{aligned} \left\| \tilde{f}_n(\cdot, 0, 1) \right\|_{\mathcal{H}^{\otimes n}}^2 &= \frac{\int_0^\infty e^{-t} \left\| \tilde{f}_n(\cdot, 0, t) \right\|_{\mathcal{H}^{\otimes n}}^2 dt}{\Gamma \left(\left[2(b+r) - \frac{b\alpha}{a} \right] n + 1 \right)} \\ &\leq \frac{2^{n[2(b+r)-\frac{b\alpha}{a}]} T_n}{\Gamma \left(\left[2(b+r) - \frac{b\alpha}{a} \right] n + 1 \right)} \\ &\leq \frac{\left(2^{[2(b+r)-\frac{b\alpha}{a}]} C'_\mu \right)^n}{\Gamma \left(\left[2(b+r) - \frac{b\alpha}{a} \right] n + 1 \right)} \end{aligned}$$

By the same reasoning as in the proof in the wave paper (details in my notebook) we can say that there exists a constant depending on α , k_α , such that

$$\left\| \tilde{f}_n(\cdot, 0, 1) \right\|_{\mathcal{H}^{\otimes n}}^2 \leq k_\alpha \frac{\left(2^{[2(b+r)-\frac{b\alpha}{a}]} C'_\mu \right)^n}{(n!)^{2(b+r)-b\alpha/a}}$$

Therefore,

$$\sum_{n \geq 0} \theta^n n! \left\| \tilde{f}_n(\cdot, 0, t) \right\|_{\mathcal{H}^{\otimes n}}^2 \leq \sum_{n \geq 0} \theta^n n! k_\alpha \frac{\left(2^{[2(b+r)-\frac{b\alpha}{a}]} C'_\mu \right)^n}{(n!)^{2(b+r)-b\alpha/a}}$$

$$= \sum_{n \geq 0} \theta^n k_\alpha \frac{\left(2^{\lfloor 2(b+r) - \frac{b\alpha}{a} \rfloor} C'_\mu\right)^n}{(n!)^{2(b+r)-1-b\alpha/a}}$$

and the last sum is finite when

$$2(b+r) - 1 - b\alpha/a > 0 \tag{2.11}$$

alpha4