

Thesis Defense:
Invariant measures for the stochastic heat equation with
no drift term

Nicholas Eisenberg

March 29, 2022

The stochastic heat equation

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) - \frac{1}{2}\Delta_x u(t, x) = b(x, u(t, x))\dot{W}(t, x), & t \geq 0, x \in \mathbb{R}^d \\ u(0, \cdot) = \mu(\cdot) \end{cases}$$

- The Stochastic Heat Equation is interpreted as follows:

$$u(t, x) = \int_{\mathbb{R}^d} G(t, x - y)\mu(dy) + \int_0^t \int_{\mathbb{R}^d} G(t - s, x - y)b(u(s, y))W(ds, dy)$$

where $G(t, x)$ is the heat kernel

$$G(t, x) = (2\pi t)^{-d/2} \exp\left(-\frac{|x|^2}{2t}\right)$$

- A processes, $\{u(t, x) : t \geq 0, x \in \mathbb{R}^d\}$, that satisfies this relation is referred to as the *mild solution*.

The noise term $\dot{W}$

- $W = \{W(\phi) : \phi \in \mathcal{S}(\mathbb{R}^d)\}$ is a centered Gaussian process uniquely defined through the following covariance functional:

$$J(\phi, \psi) := \mathbb{E}[W(\phi)W(\psi)] = \int_0^\infty ds \int_{\mathbb{R}^d} \Gamma(dx)(\phi(s, \cdot) * \tilde{\psi}(s, \cdot))(x),$$

where $\tilde{\psi}(s, x) = \psi(s, -x)$. We have the following nomenclature.


- *Space-time white noise*: $\Gamma(dx) = \delta_0(dx)$.
- *Time white and spatially colored*: $\Gamma(dx) = f(x)dx$ where f is a non-negative and non-negative definite function referred to as the *Correlation function* and its Fourier transform,

$$\hat{f}(\xi) := \mathcal{F}f(\xi) = \int_{\mathbb{R}^d} \exp(-ix \cdot \xi) f(x)dx,$$

is referred to as the *spectral density*.

Proposition (DaPrato and Zabczyk [2, Propsition 4.19])

A space-time white noise, dB , is the distributional derivative of a Brownian sheet, $B(t, x)$.



White noise expansion DaPrato [2]

If $t \geq 0$ and $0 \leq x \leq \pi$ then,

$$B(t, x) = \sum_{n=0}^{\infty} \frac{\beta_n(t)}{n + \frac{1}{2}} \sqrt{\frac{2}{\pi}} \sin \left[\left(n + \frac{1}{2} \right) x \right]$$

Walsh [4]

$$\int_t^{t+\delta_t} \int_x^{x+\delta_x} dB(ds, dy) \sim N(0, \delta_t \delta_x)$$

Brownian Sheet Simulation (Walsh [4]):

$$B(t, x) \sim B(t - \delta_t, x) + B(t, x - \delta_x) - B(t - \delta_t, x - \delta_x) + N(0, \delta_t \delta_x)$$
$$B(t, 0) = B(0, x) = 0 \text{ for all } t, x > 0.$$

Existence and uniqueness with space-time white noise

- A necessary condition for global existence, *Dalang's condition* [1], is the following:

$$\int_0^t ds \int_{\mathbb{R}^d} dy G^2(s, y) < \infty, \quad \forall t > 0.$$

- Direct calculations yields that,


$$\int_0^t ds \int_{\mathbb{R}^d} dy G^2(s, y) = \frac{\Gamma(d/2)}{2(2\pi)^d} \int_0^t \frac{1}{s^{d/2}} ds,$$

which forces,


$$d/2 < 1 \quad \Longleftrightarrow \quad d < 2.$$

- Thus a global space-time white noise solution may only exist when the spatial dimension is 1.

$$u_t - \frac{1}{2}u_{xx} = \lambda u \dot{W} \text{ with } u_0(x) = \delta_0(x) \quad (\text{white noise})$$



$\lambda = 0$





$\lambda = 1$

- Simulations done by using a finite difference scheme [5].
- We partition (space) $[0, 3]$ into n_x pieces of size $\delta_x = 3/n_x$.
- We partition (time) $[0, .2]$ into n_t pieces of size $\delta_t = .2/n_t$.
- To control error, we assume $n_x^2 \delta_t < \frac{1}{2}$ which implies $n_t \gg n_x$.

$$u(t + \delta_t, x) = u(t, x) + n_x^2 \delta_t (u(t, x + \delta_x) + u(t, x - \delta_x) - 2u(t, x)) \\ + n_x \lambda u(t, x) \text{normal}(0, \delta_t \delta_x).$$

Intermittency of $u_t - \frac{1}{2}u_{xx} = \lambda u \dot{W}$ (white noise)



- Notice the enlargement of the peaks as λ increases. We have the following theorem.

Theorem (Chen and Dalang [1, Theorem 2.4])

The second moments of the stochastic heat equation driven by space-time white noise grow exponentially in time.

Will the second moments ever be bounded in time?

- Several slides below will require the second moments to be bounded in time:

$$\sup_{t \geq 0} \|u(t, x)\|_2 < \infty, \quad \text{where } \|X\|_p = [\mathbb{E}(|X|^p)]^{1/p}.$$

- This is possible, however this will require us to only consider $d \geq 3$, and therefore we must only consider a spatially colored noise.
- Recall from above, that the covariance of a spatially colored noise is given by the following:

$$\begin{aligned} \mathbb{E}[W(\phi)W(\psi)] &= \int_0^\infty ds \int_{\mathbb{R}^d} dx f(x)(\phi(s, \cdot) * \tilde{\psi}(s, \cdot))(x) \\ &= \int_0^\infty ds \int_{\mathbb{R}^{2d}} dx dy \phi(s, x) f(x - y) \psi(s, y) \\ &= (2\pi)^{-d} \int_0^\infty ds \int_{\mathbb{R}^d} d\xi \hat{f}(\xi) \mathcal{F}\phi(s, \xi) \overline{\mathcal{F}\psi(s, \xi)} \end{aligned}$$

$$u_t - \frac{1}{2} u_{xx} = b(u) \dot{W} \text{ with } u(0, \cdot) = \mu(\cdot) \quad (\text{colored noise})$$

Theorem (Chen and Huang [2, Theorem 1.2])

There exists a unique global mild solution under the following:

- $b : \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}$ such that there exists a $L_b, L_0 > 0$ so for all $u, v \in \mathbb{R}$ and $x \in \mathbb{R}^d$,

$$|b(x, u) - b(x, v)| \leq L_b |u - v| \quad \text{and} \quad |b(x, 0)| \leq L_0.$$

- The initial measure, $\mu(dx)$ (resp. $\varphi(x)dx$), satisfies for some $a > 0$,

$$\int_{\mathbb{R}^d} e^{-a|x|^2} |\mu|(dx) < \infty \quad \left(\text{resp. } \int_{\mathbb{R}^d} e^{-a|x|^2} |\varphi(x)| dx < \infty \right),$$

where $|\mu| = \mu^+ + \mu^-$ is the *Hahn decomposition*.

- The spectral density, $\hat{f}$, satisfies *Dalang's condition* for some $\beta > 0$:

$$\Upsilon(\beta) := \int_{\mathbb{R}^d} \frac{\hat{f}(d\xi)}{\beta + |\xi|^2} < \infty.$$

Bounded moments in time

Theorem (Chen and Kim [4, Theorem 1.3 and Lemma 2.5])

Suppose that we strengthen Dalang's condition and the Lipschitz assumption to the following:

$$\Upsilon(0) := (2\pi)^{-d} \int_{\mathbb{R}^d} \frac{\hat{f}(d\xi)}{|\xi|^2} < \infty \quad \text{and} \quad L_b^2 < \frac{1}{128\Upsilon(0)}.$$

Then the following holds:

$$\begin{aligned} \|u(t, x)\|_2 &\leq C_2 \left[1 + \left(G(t, \cdot) * |\mu| \right)(x) \right] \\ &= C_2 \left[1 + \int_{\mathbb{R}^d} G(t, x - y) |\mu|(dy) \right], \quad \text{for all } t \geq 0. \end{aligned}$$

- If $\mu = \delta_0$ then $\|u(t, x)\|_2 \leq C_2 [1 + (2\pi t)^{-d/2}]$.
- Later, it will suffice consider μ such that $\sup_{t \geq t_0} \|u(t, x)\|_2 < \infty$.

Transition functions and their semigroup

Definition (Markovian Transition Functions)

- Let H be a Hilbert space and $\mathcal{B}(H)$ be the borel sets of H .
- $\{P_t(x, \Gamma) : t \geq 0, x \in H, \Gamma \in \mathcal{B}(H)\}$ forms a family of *Markovian transition functions* if,

(1) $x \mapsto P_t(x, \Gamma)$ if measurable,

(2) $\Gamma \mapsto P_t(x, \Gamma)$ is a probability measure on H ,

(3) $P_{t+s}(x, \Gamma) = \int_H P_t(x, dy) P_s(y, \Gamma)$,

(4) $P_0(x, \Gamma) = \mathbf{1}_\Gamma(x)$.

Definition (Transition Semigroup)

The above transition functions form the following semi group on $\mathcal{B}(H)$, which denotes the bounded Borel measurable functions on H :

$$P_t \varphi(x) = \int_H \varphi(y) P_t(x, dy), \quad \varphi \in \mathcal{B}(H).$$

Invariant measures

Definition (Invariant measures)

- A probability measure, η , on H is said to be *invariant* for the transition functions P_t if for all $\Gamma \in \mathcal{B}(H)$ and $t > 0$,

$$\int_H P_t(x, \Gamma) \eta(dx) = \int_H P_0(x, \Gamma) \eta(dx) = \int_H \mathbf{1}_\Gamma(x) \eta(dx) = \eta(\Gamma),$$

which is equivalent to,

$$\int_H P_t \varphi(x) \eta(dx) = \int_H \varphi(x) \eta(dx), \quad \text{for all } \varphi \in \mathcal{B}(H).$$

- This implies that the η -average of $x \mapsto P_t(x, \Gamma)$ is $\eta(\Gamma)$ and it is independent of time:

$$\text{AVG}_\eta[P_t(\cdot, \Gamma)] = \frac{1}{\eta(H)} \int_H P_t(x, \Gamma) \eta(dx) = \int_H P_t(x, \Gamma) \eta(dx) = \eta(\Gamma).$$

Invariant measures for the stochastic heat equation

- First we define the law of the solution as follows:

$$\mathcal{L}(u(t, \cdot; \zeta))(\Gamma) = \mathbb{P}[\omega \in \Omega : u(t, \cdot; \zeta) \in \Gamma], \quad \text{for all } \zeta \in H \text{ and } \Gamma \in \mathcal{B}(H),$$

and for the mean time we are intentionally vague about H .

- Our intention is for

$$P_t(\zeta, \Gamma) := \mathcal{L}(u(t, \cdot; \zeta))(\Gamma)$$

to form a Markovian semigroup.

- Established theory by DaPrato and Zabczyk [2] requires the solution,

$$\zeta \mapsto u(t, \cdot; \zeta),$$

to be almost surely H -valued for $t > 0$ and $\zeta \in H$.

- So what is H ?

$H = L^2_\rho(\mathbb{R}^d)$: weighted square integrable functions

Definition (Admissible Weight)

A positive, bounded and continuous $L^1(\mathbb{R}^d)$ function, ρ , is an *admissible weight* if

$$\frac{\left(G(t, \cdot) * \rho(\cdot)\right)(x)}{\rho(x)} \leq C_\rho(T), \quad \text{for all } t \in [0, T] \text{ and } x \in \mathbb{R}^d.$$

Definition ($L^2_\rho(\mathbb{R}^d)$)

The space $H := L^2_\rho(\mathbb{R}^d)$ is the space of ρ -weighted square integrable functions:

$$\|\zeta\|_\rho^2 := \int_{\mathbb{R}^d} |\zeta(x)|^2 \rho(x) dx < \infty.$$

Moreover, H is a Hilbert space with inner product defined as

$$\langle \zeta, \psi \rangle_\rho := \int_{\mathbb{R}^d} \zeta(x) \psi(x) \rho(x) dx.$$

Weight examples

- The following two are examples of admissible weights:

$$\exp(-a|x|), \text{ for } a > 0 \quad \text{and} \quad \frac{1}{1+|x|^a}, \text{ for } a > d.$$

- One can show that the following is not admissible for any $a > 0$,

$$\exp(-a|x|^b), \text{ for all } b > 1.$$




Figure: $x \mapsto \frac{(G(t, \cdot) * \exp(-|\cdot|))(x)}{\exp(-|x|)}$




Figure: $x \mapsto \frac{(G(t, \cdot) * \exp(-|\cdot|^{1.25}))(x)}{\exp(-|x|^{1.25})}$

The solution is almost surely H -valued.

Theorem

Let ρ be an admissible weight and suppose that

$$\Upsilon(0) = \int_{\mathbb{R}^d} \frac{\widehat{f}(\xi)}{|\xi|^2} d\xi < \infty \quad \text{and} \quad L_b^2 < \frac{1}{128\Upsilon(0)}.$$

Now suppose our initial measure, μ , satisfies the following:

$$\mathcal{G}_\rho(t; |\mu|) := \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} G(t, x - y) |\mu|(dy) \right)^2 \rho(x) dx < \infty.$$

Then for all $t > 0$,

$$\mathbb{E} \left(\|u(t, \cdot; \mu)\|_\rho^2 \right) \leq C \mathcal{G}_\rho(t; 1 + |\mu|) < \infty.$$

In particular, for any $\zeta \in H$ and for all $T > 0$, there exists a C_T such that

$$\mathcal{G}_\rho(t; \zeta) \leq 2 \|\rho\|_{L^1(\mathbb{R}^d)} + 2C_T \|\zeta\|_\rho < \infty, \quad \text{for all } t \in [0, T].$$

Invariant measures for the stochastic heat equation

- So the solution acts as an operator, mapping H to H .
- Moreover, the Markov property of the solution [2, Th 9.21] implies that

$$P_t(\zeta, \Gamma) := \mathcal{L}(u(t, \cdot; \zeta))(\Gamma), \quad \text{for all } \zeta \in H \text{ and } \Gamma \in \mathcal{B}(H)$$

for a family of Markovian transition functions.

Definition

A measure η is *invariant* for the stochastic heat equation if η is invariant for the transition functions P_t . In other words,

$$\int_H P_t(\zeta, \Gamma) \eta(d\zeta) = \int_H \mathcal{L}(u(t, \cdot; \zeta))(\Gamma) \eta(d\zeta) = \eta(\Gamma), \quad \text{for all } \Gamma \in \mathcal{B}(H).$$

Weak convergence of measures

- Let $\mathcal{M}(H)$ and $C_b(H)$ denote the probability measures and bounded continuous functions on H .
- For $\varphi \in C_b(H)$ and $\nu \in \mathcal{M}(H)$,

$$\langle \varphi, \nu \rangle = \int_H \varphi(\zeta) \nu(d\zeta).$$

Definition

We say ν_n converges *weakly* to η in $\mathcal{M}(H)$ if for all $\varphi \in C_b(H)$,

$$\langle \varphi, \nu_n \rangle \rightarrow \langle \varphi, \nu \rangle, \quad \text{as } n \rightarrow \infty.$$

We will use the notation, $\nu_n \rightarrow \nu$, to denote the weak convergence of measures.

Some existence results

Theorem (DaPrato [2, Propositions 11.2 and 11.3])

If $\zeta_1, \zeta_2 \in H$ and $\eta_1, \eta_2 \in \mathcal{M}(H)$ are such that

$$\mathcal{L}\left(u(t, \cdot; \zeta_1)\right) \rightarrow \eta_1 \text{ as } t \rightarrow \infty \quad \text{or} \quad \frac{1}{t_n} \int_0^{t_n} \mathcal{L}\left(u(s, \cdot; \zeta_2)\right) ds \rightarrow \eta_2 \text{ as } n \rightarrow \infty,$$

for some $t_n \uparrow \infty$, then both η_1 and η_2 are invariant.

- Red implies Blue but not vice versa.
- For example, suppose $f(t) \rightarrow L$ as $t \rightarrow \infty$. Then for $\epsilon > 0$, there exists N so for all $n > N$ we have the following:

$$\lim_{n \rightarrow \infty} \frac{1}{n} \int_0^n f(t) dt \leq \lim_{n \rightarrow \infty} \frac{1}{n} \left(\int_0^N f(t) dt + \int_N^n L \pm \epsilon dt \right),$$

thus,

$$L - \epsilon \leq \lim_{n \rightarrow \infty} \frac{1}{n} \int_0^n f(t) dt \leq L + \epsilon.$$

Under what setup has an invariant measure been proven to exist?

- Research on invariant measures for $u_t - \frac{1}{2}\Delta u = f(u) + b(u)\dot{W}$ on the whole space with a globally Lipschitz b is not vast.
- Tessitore and Zabczyk [3] is the only other paper which studies $u_t - \frac{1}{2}\Delta u = b(u)\dot{W}$ on the whole space with a globally Lipschitz b .
- Tessitore and Zabczyk showed there exists an $\eta \in \mathcal{M}(H)$ such that,

$$\frac{1}{t_n} \int_1^{t_n} \mathcal{L}\left(u(s, \cdot; 1)\right) ds \rightarrow \eta, \quad \text{as } n \rightarrow \infty,$$

under the following assumption on the spectral density $\widehat{f}$:

$$\int_{\mathbb{R}^d} |\xi|^{2-d} \left[\left| \mathcal{F}^{-1}\left(\sqrt{\widehat{f}}\right) \right| * \left| \mathcal{F}^{-1}\left(\sqrt{\widehat{f}}\right) \right| \right](\xi) d\xi < \infty.$$

- If one could find a f such that $\mathcal{F}^{-1} \left(\sqrt{\widehat{f}} \right) \geq 0$ and $\widehat{f} \geq 0$ then

$$\begin{aligned}
 & \int_{\mathbb{R}^d} |\xi|^{2-d} \left[\left| \mathcal{F}^{-1} \left(\sqrt{\widehat{f}} \right) \right| * \left| \mathcal{F}^{-1} \left(\sqrt{\widehat{f}} \right) \right| \right] (\xi) \, d\xi \\
 &= \int_{\mathbb{R}^d} |\xi|^{2-d} \left[\mathcal{F}^{-1} \left(\sqrt{\widehat{f}} \right) * \mathcal{F}^{-1} \left(\sqrt{\widehat{f}} \right) \right] (\xi) \, d\xi \\
 &= K \int_{\mathbb{R}^d} |\xi|^{2-d} \left[\mathcal{F}^{-1} \left(\widehat{f} \right) \right] (\xi) \, d\xi \\
 &= K \int_{\mathbb{R}^d} |\xi|^{2-d} f(\xi) \, d\xi \\
 &= K \int_{\mathbb{R}^d} |\xi|^{-2} \widehat{f}(\xi) \, d\xi = \Upsilon(0),
 \end{aligned}$$

which is precisely Dalang's condition.

- $\mathcal{F}^{-1} \left(\sqrt{\widehat{f}} \right) \geq 0$ is restrictive.

- $\mathcal{F}^{-1}\left(\sqrt{\widehat{f}}\right) \geq 0$ requires $\sqrt{\widehat{f}}$ to be non-negative definite and this may not be true even if $\widehat{f}$ is non-negative and non-negative definite.

Example

Consider the pair

$$f(x) = (2\pi)^{-1} \sin(x^2)/x^2 \quad \text{and} \quad \widehat{f}(\xi) = \frac{1}{4} \max\{2 - |\xi|, 0\}.$$

Both f and $\widehat{f}$ is clearly non-negative and thus both are non-negative definite. However, direct calculation shows that

$$\mathcal{F}^{-1}\left(\sqrt{\widehat{f}}\right)(x) = \frac{\sqrt{\frac{\pi}{2}} \left(-\cos(2x) \mathcal{S}\left(\frac{2\sqrt{x}}{\sqrt{\pi}}\right) + \sin(2x) \mathcal{C}\left(\frac{2\sqrt{x}}{\sqrt{\pi}}\right) \right)}{x^{3/2}}$$

Where $\mathcal{S}$ and $\mathcal{C}$ are the sine and cosine Fresnel integrals,

$$\mathcal{C}(z) = \int_0^z \cos\left(\frac{\pi t^2}{2}\right) dt \quad \text{and} \quad \mathcal{S}(z) = \int_0^z \sin\left(\frac{\pi t^2}{2}\right) dt.$$

$\mathcal{F}^{-1}(\sqrt{\widehat{f}})(x)$ takes positive and negative values




Figure: One example that $\mathcal{F}^{-1}(\sqrt{\widehat{f}})(x)$ assumes both positive and negative values.

- So a simpler condition must be formulated in order to give a full example.

● Recall that

$$\mathbb{E} \left(\|u(t, \cdot; \mu)\|_{\rho} \right) \leq C \mathcal{G}_{\rho}(t; \mu) := C \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} G(t, x - y)(1 + |\mu|)(dy) \right)^2 \rho(x) dx$$

Theorem

Suppose that the initial measure, μ , the diffusion term, b , and the spectral density, $\hat{f}$ satisfy

$$\Upsilon_{\alpha}(0) := \int_{\mathbb{R}^d} \frac{\hat{f}(\xi) d\xi}{|\xi|^{2(1-\alpha)}} < \infty, \quad \limsup_{t>0} \mathcal{G}_{\rho}(t; \mu) < \infty, \quad \text{and} \quad L_b^2 < \frac{1}{2^7 \Upsilon(0)},$$

for some $\alpha \in (2^7 \Upsilon(0) L_b^2, 1)$ and also for $\alpha = 0$. Then there exists an invariant measure for the stochastic heat equation. More precisely, the following weak limit exists for each $t_0 > 0$:

$$\frac{1}{t_0} \int_{t_0}^{t_n + t_0} \mathcal{L} \left(u(s, \cdot; \mu) \right) ds \rightarrow \eta_{t_0},$$

where η_{t_0} is the invariant measure.

$\widehat{f}(\xi)$ and $\Upsilon_\alpha(0)$

- It is easy to verify that $\widehat{f}(\xi) = \max\{2 - |\xi|, 0\}$ is allowable using our condition above. Indeed, for any $\alpha \in [0, 1)$,

$$\begin{aligned}\Upsilon_\alpha(0) &= \int_{\mathbb{R}^d} \frac{\max\{2 - |\xi|, 0\}}{|\xi|^{2(1-\alpha)}} d\xi \\&= \int_0^\infty \frac{\max\{2 - r, 0\}}{r^{2(1-\alpha)}} r^{d-1} dr \\&= \int_0^2 \frac{2 - r}{r^{2(1-\alpha)}} r^{d-1} dr \\&= \int_0^2 2r^{d-1-2(1-\alpha)} - r^{d-2(1-\alpha)} dr \\&< \infty \text{ when } \begin{cases} \frac{d}{2} > 1 - \alpha & 0 < \alpha < 1 \\ \frac{d}{2} > 1 & \alpha = 0. \end{cases}\end{aligned}$$

- Thus, $\frac{d}{2} > 1$, being the more restrictive condition, is what we need to assume. In other words, the spatial dimension must be $d > 2$ which means $d \geq 3$.

A comment about $\Upsilon_\alpha(0) < \infty$

- For $\alpha \in (0, 1/2)$, $\Upsilon_{2\alpha}(0) < \infty$ implies Hölder continuity of the solution [2].
- This condition may be slightly "weakened" to the following:

$$\mathcal{H}_\alpha(t) := \int_0^t dr r^{-2\alpha} \int_{\mathbb{R}^d} \widehat{f}(d\xi) \exp(-r|\xi|^2) < \infty \quad \text{for all } t > 0 \text{ and some } \alpha \in \left(0, \frac{1}{2}\right)$$

Lemma

We have the following relations between $\Upsilon_\alpha(0)$, $\Upsilon(0)$ and $\mathcal{H}_\alpha(t)$:

- (1) *However, $\Upsilon_{2\alpha}(0) < \infty \implies \mathcal{H}_\alpha(t) < \infty$ for all $t > 0$.*
- (2) *$\lim_{t \rightarrow \infty} \mathcal{H}_\alpha(t) = \Gamma(1 - 2\alpha) \Upsilon_{2\alpha}(0)$.*
- (3) *When $\Upsilon(0) < \infty$ then $\Upsilon_{2\alpha}(0) < \infty \iff \mathcal{H}_\alpha(t) < \infty$ for all $t > 0$.*

Strategy

- The strategy is to prove tightness of the laws: when given $\epsilon \in (0, 1)$, to prove the existence of a compact set $\mathcal{K}$ such that

$$\mathcal{L}u(t, \cdot; \mu)(\mathcal{K}) \geq 1 - \epsilon \quad \text{for all } t \geq t_0 > 0.$$

- This implies that

$$A_n(\mathcal{K}) := \frac{1}{n} \int_{t_0}^{n+t_0} \mathcal{L}\left(u(s, \cdot; \mu)\right)(\mathcal{K}) ds \geq \frac{1}{n} \int_{t_0}^{n+t_0} 1 - \epsilon ds = 1 - \epsilon,$$

and so the sequence of measures, A_n is tight.

- Every tight sequence of measures has a weakly convergent subsequence that converges to an invariant measure, η . Hence, there exists $(t_n) \uparrow \infty$ such that,

$$A_{t_n} \rightarrow \eta$$

- Unfortunately this does not imply that

$$\mathcal{L}u(t, \cdot; \mu) \rightarrow \eta \quad \text{as } t \rightarrow \infty.$$

Permissible spectral density

- Recall that $\widehat{f}$ must satisfy for both $\alpha = 0$ and $\alpha \in (2^7 \Upsilon(0) L_b^2, 1)$ the following:

$$\Upsilon_\alpha(0) := \int_{\mathbb{R}^d} \frac{\widehat{f}(\xi) \, d\xi}{|\xi|^{2(1-\alpha)}} < \infty.$$

- Recall that $d \geq 3$ and so $\widehat{f}$ may have slow blow up at the origin but must have fast decay at $\pm\infty$ to counter act the thick tails of $|\xi|^{2(1-\alpha)}$.
- This excludes the Riesz kernel.
- However, the Bessel kernel has polynomial blowup at the origin and exponential decay at $\pm\infty$ and will be permissible.

Bessel kernel

- Let f_s denote the Bessel kernel. When $0 < s < d$ we have that

$$f_s(x) \sim \exp(-|x|/2) \text{ for } |x| > 2 \quad f_s(x) \sim \frac{1}{|x|^{d-s+2}} \text{ for } |x| < 2.$$

- The Fourier transform $\hat{f}$ is given by $\hat{f}(\xi) = (1 + |\xi|^2)^{-s/2}$.

Example

When considering the Bessel kernel, direct calculation yields that

$$\Upsilon_\alpha(0) = \frac{\Gamma\left(\frac{d}{2} - 1 + \alpha\right) \Gamma\left(\frac{s-d}{2} + 1 - \alpha\right)}{2^d \pi^{d/2} \Gamma(s/2)},$$

and when $\alpha = 0$, this simplifies to

$$\Upsilon(0) = \frac{\Gamma\left(\frac{2+s-d}{2}\right)}{2^{d-1} \pi^{d/2} (d-2) \Gamma(s/2)},$$

and we must assume that

$$0 < \alpha < 1 - \frac{(d-s)_+}{2} \quad \text{and} \quad s > d-2 > 0.$$

Quick verification for the Bessel kernel

- Recall that $\Upsilon_\alpha(0) = C \int_{\mathbb{R}^d} \frac{\widehat{f}(\xi)}{|\xi|^{2(1-\alpha)}} d\xi$ and $\widehat{f}(\xi) = (1 + |\xi|^2)^{-s/2}$.

$$\begin{aligned}\Upsilon_\alpha(0) &= C \int_{\mathbb{R}^d} \frac{1}{|\xi|^{2(1-\alpha)} (1 + |\xi|^2)^{s/2}} d\xi \\ &= C \int_0^\infty \frac{z^{d-1}}{z^{2(1-\alpha)} (1 + z^2)^{s/2}} dz,\end{aligned}$$

- Now by the change of variables $z = r^2/(1 + r^2)$, we can evaluate the above integral explicitly by transforming it to the Beta

$$\begin{aligned}C \int_0^\infty \frac{r^{d-1}}{r^{2(1-\alpha)} (1 + r^2)^{s/2}} dr &= C \int_0^1 z^{d/2+\alpha-2} (1 - z)^{(s-d)/2-\alpha} dz \\ &= C \frac{\Gamma(d/2 - 1 + \alpha) \Gamma((s-d)/2 + 1 - \alpha)}{2\Gamma(s/2)},\end{aligned}$$

Riesz type kernel

- The g_s denote the Riesz kernel for $0 < s < d$: $g_s(x) = |x|^{-s}$.
- g_s will not satisfy

$$\Upsilon(0) = C \int_{\mathbb{R}^d} \widehat{g}_s(\xi) |\xi|^{-2} d\xi = \int_{\mathbb{R}^d} |\xi|^{d-s} |\xi|^{-2} d\xi < \infty.$$

- Instead, we can use the following

$$\widehat{r}_s(\xi) := \widehat{r}_{s_1, s_2}(\xi) = \widehat{f}_{s_1}(\xi) + f_{s_2}(\xi),$$

where $s_1 > 0$, $0 < s_2 < d$ and f_{s_1} and f_{s_2} are Bessel kernels.

- $\widehat{r}_s$ has polynomial growth at the origin and polynomial decay at infinity:

$$\widehat{r}_s(\xi) \sim \begin{cases} |\xi|^{-(d-s_2)} & |\xi| \rightarrow 0 \\ |\xi|^{-s_1} & |\xi| \rightarrow \infty, \end{cases}$$

Riese type kernel

- Notice that

$$\begin{aligned}\Upsilon_\alpha(0) &= \int_{\mathbb{R}^d} \frac{\widehat{r}_s(\xi)}{|\xi|^{2(1-\alpha)}} d\xi \\ &= \int_{\mathbb{R}^d} \frac{\widehat{f}_{s_1}(\xi)}{|\xi|^{2(1-\alpha)}} d\xi + \int_{\mathbb{R}^d} \frac{f_{s_2}(\xi)}{|\xi|^{2(1-\alpha)}} d\xi \\ &= \int_{\mathbb{R}^d} \frac{\widehat{f}_{s_1}(\xi)}{|\xi|^{2(1-\alpha)}} d\xi + (2\pi)^{-d} \int_{\mathbb{R}^d} \frac{\widehat{f}_{s_2}(\xi)}{|\xi|^{d-2(1-\alpha)}} d\xi\end{aligned}$$

- We can easily calculate that

$$\Upsilon_\alpha(0) = \frac{\Gamma\left(\frac{d}{2} - 1 + \alpha\right) \Gamma\left(\frac{s_1 - d}{2} + 1 - \alpha\right)}{2^d \pi^{d/2} \Gamma(d/2) \Gamma(s_1/2)} + \frac{\Gamma(1 - \alpha) \Gamma\left(\alpha - 1 + \frac{s_2}{2}\right)}{2^{2d} \pi^{3d/2} \Gamma(d/2) \Gamma(s_2/2)} < \infty,$$

where the finiteness requires us to assume that

$$s_1 > d - 2(1 - \alpha) > 0 \quad \text{and} \quad d > s_2 > 2(1 - \alpha) > 0.$$

Allowable Initial Conditions:

(1) For any $\zeta \in L^\infty(\mathbb{R}^d)$ then for all $t \geq 0$

$$\mathbb{E} \left(\|u(t, \cdot; \mu)\|_\rho^2 \right) \leq \left(1 + \|\zeta\|_{L^\infty(\mathbb{R}^d)} \right)^2 \|\rho\|_{L^1(\mathbb{R}^d)} < \infty.$$

(2) For $\delta_0(\cdot)$ and for all $t > 0$,

$$\mathbb{E} \left(\|u(t, \cdot; \delta_0)\|_\rho^2 \right) \leq (2 + G(t, 0)^2) \|\rho\|_{L^1(\mathbb{R}^d)} < \infty.$$

(3) Consider $|\cdot|^{-\lambda}$ for any $\lambda \in (0, d)$. Then for all $t > 0$,

$$\mathbb{E} \left(\|u(t, \cdot; |\cdot|^{-\lambda})\|_\rho^2 \right) \leq 2(1 + Ct^{-\lambda}) \|\rho\|_{L^1(\mathbb{R}^d)} < \infty, \quad C = 2^{-\lambda} \left(\frac{\Gamma((d-\lambda)/2)}{\Gamma(d/2)} \right)^2,$$

Non-permissible $L_\rho^2(\mathbb{R}^d)$ -valued initial conditions:

(1) $\zeta(x) = |x|^\alpha$. When $\rho(x) = \exp(-|x|)$, then we can show that

$$C_1(1 + t^\alpha) \leq \mathcal{G}_\rho(t; |\zeta|) \leq C_2(1 + t^\alpha).$$

Comments about the initial conditions

- Previous theory was only able to consider $\zeta \in H = L^2_\rho(\mathbb{R}^d)$ initial conditions as candidates for the following limit:

$$\lim_{n \rightarrow \infty} \frac{1}{t_n} \int_0^{t_n} \mathcal{L}(u(s, \cdot; \zeta)) ds.$$

- Now $\mu \notin H$ is now allowed
- The heat operator smooths
- $u(t, \cdot; \mu) \in H$ for $t > 0$
- For $t > t_0 > 0$,

$$\mathcal{L}u(t, \cdot; \mu) = \mathcal{L}u(t - t_0, \cdot; u(t_0, \cdot; \mu))$$




Figure: $u_t - u_{xx} = 0$ with a bumpy initial condition

References



L. Chen and R. Dalang (2015)

Moments and growth indices for the nonlinear stochastic heat equation with rough initial conditions

The Annals of Probability 43, 6, 3006–3051



L. Chen and J. Huang

Comparison principle for stochastic heat equation on $\mathbb{R}^d$

Ann. Probab., 47.2 (2019) 989–1035.



L. Chen, Y. Hu, D. Nualart (2019)

Nonlinear stochastic time-fractional slow and fast diffusion equations on $\mathbb{R}^d$.

Stochastic Processes and their Applications, 129 5073–5112.



L. Chen and K. Kim (2019)

Nonlinear stochastic heat equation driven by a spatially colored noise: moments and intermittency.

Acta Mathematica Scientia 39B(3), 645–668



A. Davie and J. Gaines (2000)

Convergence of numerical schemes for the solution of parabolic stochastic partial differential equations

Mathematics of Computation 70, 223, 121—134

References



R. Dalang (1999)

Extending the martingale measure stochastic integral with applications to spatially homogeneous s.p.d.e.'s

Electron. J. Probab.



G. DaPrato and J. Zabczyk (2014)

Stochastic Equations in Infinite Dimensions. 2nd Edition.

Cambridge University Press



G. Tessitore and J. Zabczyk (1996)

Invariant measures for the stochastic heat equation

Probability and Mathematical Statistics 18, 2, 271–287



J. Walsh (1986)

Introduction to Stochastic Partial Differential Equations