

Invariant measures for the stochastic heat equation on $\mathbb{R}^d$ without drift term

Le Chen
Auburn University
le.chen@auburn.edu

Nicholas Eisenberg
Auburn University
nze0019@auburn.edu

April 28, 2022

Abstract

This paper deals with the long term behavior of the solution to the nonlinear stochastic heat equation $\partial u / \partial t - \frac{1}{2} \Delta u = b(u) \dot{W}$ for a Lipschitz continuous diffusion coefficient b which is not necessarily bounded such as $b(u) = u$. Using the theory of the stochastic integral laid out by John Walsh, we provide conditions which will guarantee the existence of an invariant measure for a broad range of initial conditions which includes L^2_ρ as well as the Dirac delta distribution δ_0 .

MSC 2010 subject classification: 60H15, 60H07, 60F05.

Keywords: Stochastic heat equation, parabolic Anderson model, invariant measure, Dirac delta initial condition.

1 Introduction

We consider the following *stochastic heat equation* (SHE):

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) - \frac{1}{2} \Delta u(t, x) = b(x, u(t, x)) \dot{W}(t, x) & x \in \mathbb{R}^d, t > 0, \\ u(0, \cdot) = \mu(\cdot), \end{cases} \quad (1.1) \quad \boxed{\text{E:SHE}}$$

The noise, $\dot{W}(t, s)$, is a centered Gaussian noise that is white in time and homogeneously colored in space defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with natural filtration $(\mathcal{F}_t)$:

$$\mathcal{F}_t = \sigma(W_s(A) : 0 \leq s \leq t, A \in \mathcal{B}_b(\mathbb{R}^d)) \vee \mathcal{N},$$

where $\mathcal{B}_b(\mathbb{R}^d)$ denotes the bounded Borel sets on $\mathbb{R}^d$ and $\mathcal{N}$ are the $\mathbb{P}$ -null sets of Ω . Its covariance structure, J , is defined on $C_c^\infty(\mathbb{R}^{d+1}) \times C_c^\infty(\mathbb{R}^{d+1})$ as follows:

$$J(\psi, \phi) := \mathbb{E} [\dot{W}(\psi) \dot{W}(\phi)] = \int_0^\infty ds \int_{\mathbb{R}^d} \Gamma(dx) (\psi(s, \cdot) * \tilde{\phi}(s, \cdot))(x), \quad (1.2) \quad \boxed{\text{E:Cor}}$$

where $\tilde{\phi}(x) := \phi(-x)$, “ $*$ ” refers to the convolution in spatial variable, and Γ is a nonnegative and nonnegative definite tempered measure on $\mathbb{R}^d$ that is commonly referred to as the *correlation measure*. The Fourier transform of Γ (in the generalized sense) is also a nonnegative and nonnegative definite tempered measure, which is called *spectral measure* and is denoted by $\hat{f}(d\xi)$.

Moreover, in the case where Γ has density f , namely, $\Gamma(dx) = f(x)dx$, then $\widehat{f}(d\xi) = \widehat{f}(\xi)d\xi$. The initial condition μ is a deterministic, locally finite, regular, signed Borel measure that satisfies some integrability condition at the infinity (see (3.5) below). Let $\mu = \mu_+ - \mu_-$ be its *Hahn decomposition* and $|\mu| = \mu_+ + \mu_-$.

Any jointly measurable $L^2(\Omega)$ -continuous $\mathcal{F}_t$ -adapted processes, $u = (u(t, x) : t \geq 0, x \in \mathbb{R}^d)$, that solves almost surely the following stochastic integral equation is referred to as the *mild solution* to (1.1):

$$u(t, x) = \int_{\mathbb{R}^d} G(t, x - y) \mu(dy) + \int_0^t \int_{\mathbb{R}^d} b(y, u(s, y)) G(t - s, x - y) W(ds, dy), \quad (1.3) \quad \boxed{\text{E:gmsol}}$$

where $G(t, x) = (2\pi t)^{-d/2} \exp\left(-\frac{|x|^2}{2t}\right)$ is the heat kernel and the stochastic integral is the *Walsh integral* (see [Wal86; Dal+09]).

Regarding the function $b(t, x)$, we assume that for some constants $L_b > 0$ and $L_0 \geq 0$, it holds that

$$|b(x, u) - b(x, v)| < L_b |u - v| \quad \text{and} \quad |b(x, 0)| \leq L_0 \quad \text{for all } u, v \in \mathbb{R}, x \in \mathbb{R}^d. \quad (1.4) \quad \boxed{\text{E:lipcon}}$$

In particular, our assumption allows the linear case $b(x, u) = \lambda u$, which is usually referred to the *parabolic Anderson model* [CM94].

The aim of this paper is to investigate the conditions required to guarantee the existence of invariant measures for the solution to (1.1), which is a crucial step towards the study of the ergodicity of the system. We will see that for any $t_0 > 0$ that the solution of (1.1), $u(t_0, \cdot)$, belongs to the Hilbert space $\mathcal{E}_{t_0} = (E_{t_0}, \|\cdot\|_{E_{t_0}})$; see Remark 1.1 below. Then by an invariant measure for (1.1), we mean a measure, η , on $\mathcal{B}(E_{t_0})$ such that

$$\eta(A) = \int_E \mathcal{L}(u(t, \cdot, \varphi))(A) \eta(d\zeta), \quad \text{for all } t \geq t_0, \quad (1.5) \quad \boxed{\text{D:InvMeas?}}$$

where $A \in \mathcal{B}(L^2_\rho(\mathbb{R}^d))$ and $\mathcal{L}(u(t, \cdot, \varphi))(A)$ is the law of (1.1) starting from $\varphi \in E_{t_0}$;

$$\mathcal{L}(u(t, \cdot, \varphi))(A) = \mathbb{P}[\omega \in \Omega : u(t, \cdot, \varphi)(\omega) \in A], \quad t \geq t_0. \quad (1.6) \quad \boxed{\text{D:law?}}$$

Due the *Krylov-Bogoliubov theorem* (see [DZ96, Theorem 3.1.1]), we see that that existence of the measure can depend on the initial condition and under suitable conditions, for example the assumptions of Theorem 1.4 below, the following measure, η , is invariant for (1.1):

$$\eta(A) = \lim_{n \rightarrow \infty} \frac{1}{T_n} \int_{t_0}^{T_n + t_0} \mathcal{L}(u(t, \cdot, \varphi))(A) dt, \quad \text{for any } t_0 > 0. \quad (1.7) \quad \boxed{\text{E:InvMeasForm}}$$

In the above, $\{T_n\}$ is an appropriately chosen sequence of positive numbers increasing to infinity.

We note that proving uniqueness of the invariant measure is a much harder question. An open question is if φ_1 and φ_2 are two such initial conditions that are suitable for (1.7) and if η_1 and η_2 are the two invariant measures that arise from φ_1 and φ_2 respectively then does $\eta_1 = \eta_2$? Another question one can ask in the absence of uniqueness is what initial data are applicable to (1.7) and can we give explicit examples? We address the second question in Examples 2.1 and 2.2 and also in the discussion below.

Existence in unbounded domains has been studied in few papers such as [AM03; MSY20; MSY16; TZ98]. Since the spatial domain is unbounded, one needs to introduce a weight function ρ and the corresponding weighted space $L^2_\rho(\mathbb{R}^d)$ as usual; see Section 3 below for more details.

Following the standard procedure as laid out by Da Prato and Zabczyk [DGZ92], the main effort is to show that:

$$\sup_{t>0} \mathbb{E} \left(\|u(t, \cdot)\|_{L^2_\rho(\mathbb{R}^d)}^2 \right) < \infty. \quad (1.8) \quad \boxed{\text{E:BddMnt}}$$

However, this solution is usually *intermittent*, namely, the probability moments to (1.1) have a certain exponential growth in t . In [AM03; MSY20; MSY16], a drift term $f(x, u)$ is included in (1.1) in a crucial way to help to cancel the otherwise exponential growing moments. The absence of such a drift term in this paper makes the problem more challenging. In order to have moments bounded in time, as required in (1.8), one has to first identify the exact conditions, under which the moments to (1.1) do not possess exponential growth. This question has been answered in [CK19, Theorem 1.3 and Lemma 2.5], where necessary and sufficient conditions are given. More precisely, for bounded second moments one has to have the spatial dimension $d \geq 3$, and in addition, the spectral measure, $\hat{f}$, and Lipschitz constant, L_b , (the Lyapunov exponent of b) in (1.4) must satisfy the following:

$$\Upsilon(0) := \int_{\mathbb{R}^d} \frac{\hat{f}(d\xi)}{|\xi|^2} < \infty \quad \text{and} \quad 64L_b^2 < \frac{1}{4\Upsilon(0)}; \quad (1.9) \quad \boxed{\text{E:U0finite}}$$

see the Assumption (3.8) below.

In the setting of equation (1.1), Tessitore and Zabczyk [TZ98] prove existence of an invariant measure for (1.1) in $L^2_\rho(\mathbb{R}^d)$ under the assumption that there exists a $\varphi \in L^2_\rho(\mathbb{R}^d) \cap L^2_{\hat{\rho}}(\mathbb{R}^d)$ where $\rho(\hat{\rho})^{-1} \in L^1(\mathbb{R}^d)$ and the solution starting from φ is bounded in probability in $L^2_{\hat{\rho}}(\mathbb{R}^d)$ and also that the spectral density $\hat{f}$ satisfies

$$\hat{f} \in L^p(\mathbb{R}^d) \quad \text{where} \quad \frac{d-2}{d} < \frac{1}{p}; \quad (1.10) \quad \boxed{\text{E:CorTZ?}}$$

see Hypothesis 2.1 (*ibid.*). However as was illustrated in [TZ98, Theorem 3.3], in order to apply this theorem, the following additional assumptions were needed: $d \geq 3$ and the Lipschitz constant of b must satisfy $L(b)^{-2} > 1$, where

$$\tilde{\Gamma} := \left| \sqrt{\mathcal{F}(f)} \right| * \left| \sqrt{\mathcal{F}(f)} \right| \quad \text{and} \quad 1 := \frac{\Gamma(d/2 - 1)}{4\pi^{d/2}} \int_{\mathbb{R}^d} \tilde{\Gamma}(d\zeta) |\zeta|^{2-d}, \quad (1.11) \quad \boxed{\text{E:TZspec}}$$

where Γ is the Gamma function. With these assumptions, they were able to prove that (1.1) starting from the constant 1 satisfies (1.8) and thus is bounded in probability, verifying the existence of an invariant measure. Moreover, the measure takes the form of (1.7) above with $\varphi = 1$.

The additional assumptions given by Tessitore and Zabczyk parallel our assumptions (1.9) and (??). However, it is not at all trivial to check which spectral densities, $\hat{f}(d\xi)$, satisfy (1.11) and no such examples were given in [TZ98]. For example, it may not be obvious from (1.11) that one can not take the spectral density of the noise to be the Riesz kernel, $\hat{f}(\xi) = |\xi|^{d-\alpha}$, for any $0 < \alpha < d$ but this can be easily seen from our condition (1.9). The condition that $d \geq 3$ forces us to only consider a colored noise as in (1.2) and so the loss of the ability to apply the Riesz kernel is unfortunate. However, as we will see in Example 2.1 below, we are able to apply the Bessel kernel, $B_s(x)$, with $s < d$, which has exponential decaying tails and power growth at the origin. So in absence of the Riesz kernel, we still may apply our results to a kernel, namely the Bessel kernel, that has origin blow up.

Moreover by applying a restart argument (see Remark 1.1 and Example 2.1 below), our method allows us to verify (1.8), and in return (1.7), for more general initial conditions $\mu(\cdot)$

which include:

Dirac delta measure $\delta_0(dx)$ and $\varphi(x)dx$ with $\varphi \in C^\infty(\mathbb{R}^d)$.

Thus, we can expand our choice of initial data to include singular distributions, which are outside the space of $L_\rho^2(\mathbb{R}^d)$. We will expand on this in the following remark.

(R:Restart) Remark 1.1 (Restarted SHE). Suppose that $u(t_0, \cdot, \mu)$ is the solution to (1.1) starting from an initial condition, μ , that satisfies the assumptions of Theorem 1.3. Then (1.16) and Theorem 3.5 imply that $u(t_0, \cdot, \mu)$ belongs to the Hilbert space, $\mathcal{E}_{t_0} := (E_{t_0}, \|\cdot\|_{E_{t_0}})$, where E_{t_0} is the space of random fields, $\{\varphi_{t_0}(x)\}_{x \in \mathbb{R}^d}$, that are $\mathcal{F}_{t_0} \times \mathcal{B}(\mathbb{R}^d)$ measurable with the following finite norm:

$$\|\varphi_{t_0}\|_{E_{t_0}}^2 := \mathbb{E} \left(\|\varphi_{t_0}(\cdot)\|_{L_\rho^2(\mathbb{R}^d)}^2 \right) < \infty.$$

Then, for any $t_0 > 0$, consider the following restarted equation:

$$\begin{cases} \frac{\partial v}{\partial t}(t, x) - \frac{1}{2} \Delta v(t, x) = b(x, v(t, x)) \dot{W}_{t_0}(t, x) & x \in \mathbb{R}^d, t > 0, \\ v(0, \cdot) = u(t_0, \cdot), \end{cases} \quad (1.12) \quad \boxed{\text{E:restart}}$$

where $\dot{W}_{t_0}(t, x) := \dot{W}(t + t_0, x)$ denotes the time shifted noise, ie,

$$\int_0^t \int_{\mathbb{R}^d} W_{t_0}(ds, dy) = \int_{t_0}^{t+t_0} \int_{\mathbb{R}^d} W(ds, dy).$$

Moreover, the Markov property of the solution to (1.1) implies that for any $t \geq 0$, we have $u(t + t_0, x) \stackrel{\mathcal{L}}{=} v(t, x)$ where $\mathcal{L}$ denotes equality in law. Since we are interested in studying the long time behavior of (1.1), we obtain the same information by choosing to study the restarted equation.

This is a very import observation. Recall that Tessitore et.al. [TZ98, Theorem 3.3] proved that there exists an invariant measure, η , on $\mathcal{B}(L_\rho^2(\mathbb{R}^d))$ of the form (1.7) for the constant initial data $\varphi = 1$. Because of the restart property of the equation, we may rewrite (1.7) as the following:

$$\begin{aligned} \eta(A) &= \lim_{n \rightarrow \infty} \int_{t_0}^{T_n+t_0} \mathcal{L}(u(t, \cdot, 1))(A) dt \\ &= \lim_{n \rightarrow \infty} \int_{t_0}^{T_n+t_0} \mathcal{L} \left[v \left(t - \frac{t_0}{2}, \cdot, u \left(\frac{t_0}{2}, \cdot, 1 \right) \right) \right] (A) dt \\ &= \lim_{n \rightarrow \infty} \int_{t'_0}^{T_n+t'_0} \mathcal{L} \left[v \left(t, \cdot, u \left(\frac{t_0}{2}, \cdot, 1 \right) \right) \right] (A) dt, \quad t'_0 := t_0/2, \end{aligned}$$

where we recall that $A \in \mathcal{B}(L_\rho^2(\mathbb{R}^d))$ and v denotes the solution to (1.12) starting from $u(t_0/2, \cdot, 1)$ with the time-shifted noise $\dot{W}_{t_0/2}$. Naturally the invariant measure is usually defined on the Hilbert space of functions where the initial condition lives; in the case of [TZ98], this is $L_\rho^2(\mathbb{R}^d)$. The above shows that we can extend the measure η to $\mathcal{B}(E_{t_0})$ since we can associate the equation (1.1) starting from $\mu = 1$ with the above restarted process v that starts from $u(t_0, \cdot, 1) \in E_{t_0}$. This is no contradiction since clearly $\mathcal{B}(L_\rho^2(\mathbb{R}^d)) \subset \mathcal{B}(E_{t_0})$.

This observation will allow us to show existence of an invariant measure in $\mathcal{B}(E_{t_0})$ for our equation (1.1) starting from an initial condition outside of $L_\rho^2(\mathbb{R}^d)$. For example, we will see

in Example 2.1 below that we may start our equation at the Dirac delta measure δ_0 , which is clearly not in $L^2_\rho(\mathbb{R}^d)$. However, because of the smoothing properties of the heat operator, after any amount of time $t_0 > 0$, the solution $u(t_0, \cdot, \delta_0) \in E_{t_0}$. Thus, whether we originally start our equation (1.1) at the constant 1, as in [TZ98], or at a distribution that satisfies the assumptions of Theorem 1.4 main below, like the Dirac delta distribution δ_0 , then either way we end up in the same function space. If one were able to prove uniqueness of the invariant measure, then the invariant measure which is derived from the Dirac delta initial condition and the invariant measure which is derived from the constant 1 initial condition would be the same.

For the remainder of the paper, μ will always denote the initial condition to (1.1) and $u(t, \cdot) = u(t, \cdot, \mu)$ will always denote the solution of (1.1) starting from μ . In addition, $v_{t_0}(t, \cdot) = v(t, \cdot, u(t_0, \cdot, \mu))$ will always denote the solution to the restarted equation (1.12) starting from $u(t_0, \cdot, \mu)$ and driven by the time shifted noise $\dot{W}_{t_0}$. When there is no confusion, we may write v instead of v_{t_0} .

Notation **Notation 1.2.** Denote $\|\cdot\|_\rho := \|\cdot\|_{L^2_\rho(\mathbb{R}^d)}$ and $\|X\|_p := (\mathbb{E}(|X|^p))^{1/p}$. In addition, we define

$$J_u(t, x, \mu) := (G(t, \cdot) * \mu)(x) = \int_{\mathbb{R}^d} G(t, x - y) \mu(dy), \quad \text{and} \quad (1.13) \text{ ?E:Ju?}$$

$$J_u^*(t, x) := J_0(t, x, \mu^*)(t, x) \quad \text{with} \quad \mu^* := 1 + |\mu|. \quad (1.14) \text{ [E:J*u]}$$

We first present a preliminary result which is used to prove Theorem 1.4 below.

Theorem 1.3. Let $u(t, x)$ be the solution to (1.1) starting from μ . Assume that

(i) $\rho : \mathbb{R}^d \rightarrow \mathbb{R}_+$ is an nonnegative $L^1(\mathbb{R}^d)$ function;

(ii) the initial measure μ satisfies

$$\mathcal{G}_\rho^u(t) := \int_{\mathbb{R}^d} (J_u^*(t, x))^2 \rho(x) dx < \infty \quad \text{for all } t > 0; \quad (1.15) \text{ [E:G]}$$

(iii) the correlation measure Γ and the Lipschitz constant L_b of b satisfy conditions in (1.9).

Then for all $t > 0$, $u(t, \cdot) \in L^2_\rho(\mathbb{R}^d)$ a.s. Moreover, for some constant $C > 0$ which does not depend on t , it holds that

$$\mathbb{E} \left(\|u(t, \cdot)\|_\rho^2 \right) \leq C \mathcal{G}_\rho^u(t) < \infty. \quad (1.16) \text{ [E:uInLrho]}$$

Proof. Under condition (iii), we can apply Fubini's Theorem and the moment bound (3.9) below to see that for some constant $C > 0$ independent of t , which may vary from line to line, it holds that

$$\begin{aligned} \mathbb{E} \left(\|u(t, \cdot)\|_\rho^2 \right) &= \mathbb{E} \left(\int_{\mathbb{R}^d} |u(t, x)|^2 \rho(x) dx \right) \\ &\leq C \int_{\mathbb{R}^d} \left(1 + (G(t, \cdot) * |\mu|)(x) \right)^2 \rho(x) dx \\ &= C \int_{\mathbb{R}^d} \left((G(t, \cdot) * (1 + |\mu|))(x) \right)^2 \rho(x) dx \\ &= C \mathcal{G}_\rho(t) < \infty. \end{aligned}$$

This proves Theorem 1.3. □

Here is the main theorem of the paper:

Theorem 1.4. *Let $u(t, x)$ be the solution to (1.1) starting from μ . Assume that all conditions in Theorem 1.3 are satisfied. Moreover, we assume that*

- (i) *if ρ is an admissible weight (see Definition 3.1 below) and there exists another admissible weight $\hat{\rho}$ such that*

$$\int_{\mathbb{R}^d} \frac{\rho(x)}{\hat{\rho}(x)} dx < \infty; \quad (1.17) \quad \boxed{\text{E:rho rho}}$$

- (ii) *the weight function $\hat{\rho}$ and the initial condition satisfy the following condition:*

$$\limsup_{t>0} \mathcal{G}_{\hat{\rho}}^u(t + t_0) < \infty; \quad (1.18) \quad \boxed{\text{E:InitData}}$$

- (iii) *for some $\alpha \in (0, 1/2)$, the spectral measure $\hat{f}$ satisfies that*

$$\mathcal{H}_\alpha(t) := \int_0^t dr r^{-2\alpha} \int_{\mathbb{R}^d} \hat{f}(d\xi) \exp(-r|\xi|^2) < \infty \quad \text{for all } t > 0; \quad (1.19) \quad \boxed{\text{E:H}}$$

- (iv) *for the constant α in (ii), there exists $q > 1/\alpha$ (i.e., $q^{-1} < \alpha < 2^{-1}$) such that*

$$\int_0^t \left[\mathcal{G}_{\hat{\rho}}^u(s + t_0) \mathcal{H}_\alpha(s) \right]^{q/2} ds < \infty \quad \text{for all } t > 0; \quad (1.20) \quad \boxed{\text{E:GH-int}}$$

- (v) *the second condition in (1.9) is strengthened accordingly with q given in (iv) to*

$$qL_b^2 < \frac{1}{128\Upsilon(0)}.$$

Then we have that

- (a) *for any $t_0 > 0$, the sequence of laws of $\{\mathcal{L}u(t, \cdot)\}_{t \geq t_0}$ is tight, i.e., for any $\epsilon \in (0, 1)$, there exists a compact set $\mathcal{K} \subset L_\rho^2(\mathbb{R}^d)$ such that*

$$\mathcal{L}u(t, \cdot)(\mathcal{K}) := \mathbb{P}(u(t, \cdot) \in \mathcal{K}) \geq 1 - \epsilon, \quad \text{for all } t \geq t_0 > 0; \quad (1.21) \quad \boxed{\text{E:cpt}}$$

- (b) *there exists an invariant measure for the laws $\{\mathcal{L}u(t, \cdot)\}_{t > t_0}$.*

We will postpone the proof of Theorem 1.4 (a) to Section 4. The proof of part (b) based on (a) is routine:

Proof of Theorem 1.4 (b). Fix an arbitrary $T_0 > 0$ and denote

$$U(T) := \frac{1}{T} \int_{T_0}^{T+T_0} \mathcal{L}(u(t, \cdot)) dt, \quad T > 0.$$

We claim that the family of laws $U(T, \cdot)$ for $T > 0$ is tight in $L_\rho^2(\mathbb{R}^d)$. Indeed, for any $\epsilon \in (0, 1)$, by part (a), there exists a compact set $\mathcal{K} \in L_\rho^2(\mathbb{R}^d)$ such that (1.21) holds. This implies that

$$U(T)(\mathcal{K}) = \frac{1}{T} \int_{T_0}^{T+T_0} \mathcal{L}(u(t, \cdot))(\mathcal{K}) dt \geq \frac{1}{T} \int_{T_0}^{T+T_0} (1 - \epsilon) dt = 1 - \epsilon, \quad \text{for all } T > 0.$$

Let $\{T_n\}_{n \in \mathbb{N}}$ be any deterministic sequence such that $T_n \uparrow \infty$. Since $\{U(T_n)\}_{n \geq 1}$ is a tight sequence of measures, there exists a subsequence $\{U(T_{n_m})\}_{m \geq 1}$ that converges weakly to a measure, say, η , on $L_\rho^2(\mathbb{R}^d)$. Then one can apply for the *Krylov-Bogoliubov theorem* (see, e.g., [DZ96, Theorem 3.1.1]) to conclude that the measure η is the invariant measure for $\mathcal{L}(u(t, \cdot))$, $t \geq T_0$. Finally, we can choose T_0 to be arbitrarily close to zero. This completes the proof of part (b) of Theorem 1.4. $\square$

2 Examples

Example 2.1 (Bessel kernel and δ_0 initial data). We end this section with an example of an applicable initial data and spectral density. Note that because of (1.20), the initial data and spectral density must exist in tandem. Let's consider the initial data $\mu(\cdot) = \delta_0(\cdot)$ and let the correlation function of W be $f(x) = B_s(x)$, where B_s denotes the Bessel kernel with parameter s . It is shown in [Gra14, Section 1.2.2] that there exists $c(s, d)$ and $C(s, d)$ such that for $0 < s < d$

$$B_s(x) \leq C(s, d) \exp(-|x|/2) \quad \text{for } |x| \geq 2,$$

and for $H(x) = |x|^{s-d} + 1 + O(|x|^{s-d+2})$,

$$\frac{1}{c(s, d)} \leq \frac{B_s(x)}{H_s(x)} \leq c(s, d) \quad \text{for } |x| \leq 2.$$

Hence $B_s(x)$ has blow up at the origin and exponential decay at infinity. Moreover,

$$\mathcal{F}G_s(\xi) = \frac{1}{(1 + |\xi|^2)^{s/2}}. \quad (2.1) \quad \boxed{\text{E:FB}}$$

It is easy to see that δ_0 satisfies (3.5). Moreover, it is easy to see that if $d - 2 < s < d$ then $\Upsilon(0) < \infty$ and so the conditions of Theorem (3.5) and Theorem 3.6 are satisfied and a unique mild solution, u , exists that starts from δ_0 and is driven by a Gaussian noise with correlation function $B_s(x)$. We denote $v(t, x)$ to be the restarted solution as in (1.12). We now show that v satisfies the conditions in Theorem 1.4. In other words, it remains to prove that $v(0, \cdot) = u(t_0, \cdot)$ and $B_s(x)$ satisfy the conditions (1.18), (1.19) and (1.20).

Let $q > 2$ be arbitrary. Starting with (1.18),

$$\begin{aligned} \mathcal{G}_{\hat{\rho}, q}(s) &= \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} G(s, x - y) \mu_q^*(dy) \right)^2 \rho(x) dx \\ &= \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} G(s, x - y) (1 + \|v(0, y)\|_q)(dy) \right)^2 \hat{\rho}(x) dx \\ &= \int_{\mathbb{R}^d} \left(1 + \int_{\mathbb{R}^d} G(s, x - y) \|v(0, y)\|_q dy \right)^2 \hat{\rho}(x) dx \\ &\leq 2 \|\hat{\rho}\|_{L^1(\mathbb{R}^d)} + 2 \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} G(s, x - y) \|u(t_0, y)\|_q dy \right)^2 \hat{\rho}(x) dx, \end{aligned}$$

where in the last line we used Recall that $u(t_0, x) = v(0, x)$. In the following, C is a generic constant independent of s that may change from line to line. By applying the moment bound (3.9) we see that

$$\begin{aligned} \mathcal{G}_{\hat{\rho}, q}(s) &\leq 2 \|\hat{\rho}\|_{L^1(\mathbb{R}^d)} + 2C \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} G(s, x - y) [1 + (G(t_0, \cdot) * \delta_0)(y)] dy \right)^2 \hat{\rho}(x) dx \\ &= 2 \|\hat{\rho}\|_{L^1(\mathbb{R}^d)} + C \int_{\mathbb{R}^d} \left(1 + \int_{\mathbb{R}^d} G(s, x - y) (G(t_0, \cdot) * \delta_0)(y) dy \right)^2 \hat{\rho}(x) dx \\ &= 2 \|\hat{\rho}\|_{L^1(\mathbb{R}^d)} + C \int_{\mathbb{R}^d} (1 + (G(t_0 + s, \cdot) * \delta_0)(x))^2 \hat{\rho}(x) dx \\ &\leq 2 \|\hat{\rho}\|_{L^1(\mathbb{R}^d)} + 2C \|\hat{\rho}\|_{L^1(\mathbb{R}^d)} + 2C \int_{\mathbb{R}^d} (G(t_0 + s, \cdot) * \delta_0)(x)^2 \hat{\rho}(x) dx \\ &\leq C \|\hat{\rho}\|_{L^1(\mathbb{R}^d)} + C \int_{\mathbb{R}^d} (G(t_0 + s, x))^2 \hat{\rho}(x) dx \end{aligned}$$

$$\begin{aligned}
&= C \|\widehat{\rho}\|_{L^1(\mathbb{R}^d)} + C \frac{(2\sqrt{\pi})^{-d}}{(t_0 + s)^{d/2}} \int_{\mathbb{R}^d} G\left(\frac{t_0 + s}{2}, x\right) \widehat{\rho}(x) dx \\
&=: \mathbf{G}(s),
\end{aligned}$$

and this verifies (1.18) because $t_0 > 0$ is fixed. Moreover, this implies that for any $T > 0$, $\mathcal{G}_{\widehat{\rho},q}(s)$ is bounded on $s \in [0, T]$. This will be needed below.

Now let's verify (1.19) for the Bessel kernel, B_s . By recalling (2.1), we have

$$\begin{aligned}
\mathcal{H}_\alpha(s) &= \int_0^s dr r^{-2\alpha} \int_{\mathbb{R}^d} d\xi \frac{\exp(-r|\xi|^2)}{(1 + |\xi|^2)^{s/2}} \\
&= \sigma(\mathbb{S}^{d-1}) \int_0^t dr r^{-2\alpha} \int_0^\infty dz \frac{\exp(-rz^2)}{(1 + z^2)^{s/2}} z^{d-1}.
\end{aligned}$$

Let's examine the $\int_0^\infty dz$ integral and apply the change of variables $y = z^2$. This yields,

$$\int_0^\infty dz \frac{\exp(-rz^2)}{(1 + z^2)^{s/2}} z^{d-1} = \frac{1}{2} \int_0^\infty dy e^{-ry} (1 + y)^{s/2} y^{\frac{d}{2}-1} dy = \frac{\Gamma(d/2)}{2} U\left(\frac{d}{2}, \frac{d+2-s}{2}, r\right),$$

where $U(\alpha, \beta, t)$ is the confluent hypergeometric U function (see [Olv+10, Chapter 13]) and the last equality can be found in [Olv+10, p. 13.4.4]. Thus, to show the finiteness of $\mathcal{H}_\alpha(s)$ for each $s > 0$, it suffices to show that the following integral is finite:

$$\int_0^s r^{-2\alpha} U\left(\frac{d}{2}, \frac{d+2-s}{2}, r\right) dr. \quad (2.2) \quad \boxed{\text{E:BesselHG}}$$

The behavior of $U(\alpha, \beta, t)$ around the origin is given in [Olv+10, Section 13.2 (iii)] and is given in cases on β , which in our case is $(d+2-s)/2$. Recall above that we are taking $d-2 < s < d$. For this reason, the only case in [Olv+10, Section 13.2 (iii)] that is applicable is $1 < (d+2-s)/2 < 2$ and in this case

$$r^{-2\alpha} U\left(\frac{d}{2}, \frac{d+2-s}{2}, r\right) = O\left(r^{1-2\alpha-(d+2-s)/2} + r^{-2\alpha} + r^{2-2\alpha-(d+2-s)/2}\right). \quad (2.3) \quad \{?\}$$

So in order (2.2) to be finite we need

$$\min\{1 - 2\alpha - (d+2-s)/2, -2\alpha, 2 - 2\alpha - (d+2-s)/2\} \geq -1.$$

Recall that we are assuming $\alpha \in (q^{-1}, 2^{-1})$ so $-2\alpha > -1$. Thus it suffices to set

$$1 - 2\alpha - (d+2-s)/2, -2\alpha \geq -1 \quad \Longleftrightarrow \quad 2 - 4\alpha - d + s > 0.$$

This is indeed possible as $2 - 4\alpha > 0$ and $d - 2 < s < d$, thus one can take s sufficiently close to d so that $(2 - 4\alpha) - (d - s) > 0$. This completes the verification for (1.19).

Lastly we need to verify (1.20). But essentially all the work for this has been completed. The verification of (1.19) implies that $\mathcal{H}_\alpha(s)$ is continuous on $s \in [0, T]$ for any $T > 0$ and we already saw above that $\mathcal{G}_{\widehat{\rho},q}(s)$ was bounded on $[0, T]$. These two facts imply (1.20), thus allowing us to apply Theorem (1.4) and verify the existence of an invariant measure to (1.1) with initial data, $\mu(\cdot) = u(t_0, \cdot)$ and noise $\dot{W}_{t_0}$ with correlation function given to be the Bessel kernel B_s . Recall that the invariant measures derived from specific initial conditions take the form of (1.7). Since for all $t > 0$ we have $u(t + t_0, x) \stackrel{\mathcal{L}}{=} v(t, x)$, what we have actually done was verify the existence of an invariant measure, η , for (1.1) starting from δ_0 with correlation function $B_s(x)$. Indeed,

$$\eta(A) = \lim_{n \rightarrow \infty} \int_{t_0}^{T_n + t_0} \mathcal{L}(v(t, \cdot, u(t_0, \cdot)))(A) dt$$

$$\begin{aligned}
&= \lim_{n \rightarrow \infty} \int_{t_0}^{T_n+t_0} \mathcal{L}(u(t+t_0, \cdot, \delta_0))(A) \, dt \\
&= \lim_{n \rightarrow \infty} \int_{t'_0}^{T_n+t'_0} \mathcal{L}(u(t, \cdot, \delta_0))(A) \, dt, \quad t'_0 = 2t_0.
\end{aligned}$$

Hence we may start (1.1) at δ_0 .

(Ex:More) **Example 2.2** (More examples). We would like to point out that the following initial data are also applicable when considering the correlation function to be the Bessel kernel $B_s(x)$ with $s < d$:

$$\text{Lebesgue measure } dx \quad \text{and} \quad \varphi(x)dx \quad \text{with } \varphi \in C^\infty(\mathbb{R}^d).$$

The interested reader may go through the same steps above to see that (1.18), (1.19) and (1.20) are all finite.

Example 2.3 (Bounded initial conditions). When the initial condition is bounded, namely, $\mu(dx) = \phi(x)dx$ with $\phi \in L^\infty(\mathbb{R}^d)$, our conditions in our Theorems 1.3 and 1.4 will be greatly simplified. (Give details here...)

3 Preliminaries

(S:Prelim) Let ρ be some nonnegative function on $\mathbb{R}^d$. We denote by $L_\rho^2(\mathbb{R}^d)$ the weighted Hilbert space with the inner product $\langle f, g \rangle_\rho := \int_{\mathbb{R}^d} f(x)g(x)\rho(x) \, dx$ and set $\|f\|_\rho^2 := \int_{\mathbb{R}^d} f(x)^2\rho(x) \, dx$. Following Tessitore and Zabczyk [TZ98], we make the following definition:

(D:Rho) **Definition 3.1.** $\rho : \mathbb{R}^d \mapsto \mathbb{R}$ is called an *admissible weight* if ρ is a strictly positive, bounded and continuous function in $L^1(\mathbb{R}^d)$ such that for $T > 0$ there exists a constant $C_\rho(T)$ such that

$$(G(t, \cdot) * \rho(\cdot))(x) \leq C_\rho(T)\rho(x) \quad \text{for all } t \in [0, T] \text{ and } x \in \mathbb{R}^d, \quad (3.1) \quad \boxed{\text{E:aw}}$$

or equivalently,

$$\sup_{x \in \mathbb{R}^d, t \in [0, T]} \frac{(G(t, \cdot) * \rho(\cdot))(x)}{\rho(x)} \leq C_\rho(T). \quad (3.2) \quad \text{?E:awbd1?}$$

Here are several examples of admissible weights (see Example in Section 2 of [TZ98])

$$\begin{aligned}
\rho(x) &= \exp(-a|x|), & a > 0 \quad \text{and} \\
\rho(x) &= (1 + |x|^a)^{-1}, & a > d.
\end{aligned} \quad (3.3) \quad \boxed{\text{E:admRho}}$$

(R:admRho) **Remark 3.2.** The smaller the weight function $\rho(\cdot)$ (not necessarily to be admissible here) is, the larger the space $L_\rho^2(\mathbb{R}^d)$ is. For example, one may choose ρ to be either a nonnegative function with compact support or the heat kernel itself $G(1, \cdot)$. In both cases, ρ is smaller than those in (3.3) (up to a constant). However, one can easily check that the admissible condition (3.1) excludes these two cases. It is much less obvious that the following weight function is not admissible:

$$\rho(x) = \exp(-a|x|^b), \quad x \in \mathbb{R}^d, \quad \text{with } a > 0 \text{ and } b \in (1, 2) \text{ fixed.}$$

(P:weight)? **Proposition 3.3** (Proposition 2.1 of [TZ98]). *For any admissible weight ρ , the operators on $L_\rho^2(\mathbb{R}^d)$ defined by $\varphi \mapsto (G(t, \cdot) * \varphi(\cdot))(x)$ can be extended to a C_0 -semigroup on L_ρ^2 . Moreover, if $\widehat{\rho}$ is another admissible weight such that*

$$\int_{\mathbb{R}^d} \frac{\rho(x)}{\widehat{\rho}(x)} dx < \infty,$$

then for any $t < 0$, the operators defined above are compact from $L_\rho^2(\mathbb{R}^d)$ to $L_{\widehat{\rho}}^2(\mathbb{R}^d)$.

Find out if the following lemma has been used, if not we may remove the Lemma.

$\langle \text{L:RhoDom} \rangle$ **Lemma 3.4.** *For any admissible weight function ρ and $a > 0$, there exists a constant $K_{\rho,a} > 0$ such that*

$$\exp(-a|x|^2) \leq K_{\rho,a} \rho(x), \quad \text{for all } x \in \mathbb{R}^d. \quad (3.4) \quad \langle \text{E:RhoDom} \rangle$$

Proof. Let ρ be an arbitrary admissible weight function and $a > 0$. By (3.1),

$$\begin{aligned} \frac{\exp(-a|x|^2)}{\rho(x)} &= \frac{\exp(-a|x|^2)}{\rho(x)} \frac{(G(1, \cdot) * \rho(\cdot))(x)}{(G(1, \cdot) * \rho(\cdot))(x)} \\ &\leq \left(\sup_{x \in \mathbb{R}^d, t \in [0,1]} \frac{(G(t, \cdot) * \rho(\cdot))(x)}{\rho(x)} \right) \frac{\exp(-a|x|^2)}{(G(1, \cdot) * \rho(\cdot))(x)} \\ &\leq C_\rho(1) \frac{\exp(-a|x|^2)}{(G(1, \cdot) * \rho(\cdot))(x)}. \end{aligned}$$

Noticing that for some constant $\Theta > 0$, $\rho(x) \geq \Theta \prod_{\ell=1}^d \mathbb{I}_{[-1,1]}(x_\ell)$, a simple exercise shows that

$$C_{\rho,a} := \sup_{x \in \mathbb{R}^d} \frac{\exp(-a|x|^2)}{(G(1, \cdot) * \rho(\cdot))(x)} \leq \frac{1}{\Theta} \left(\sup_{x \in \mathbb{R}} \frac{\exp(-ax^2)}{(G(1, \cdot) * \mathbb{I}_{[-1,1]}(\cdot))(x)} \right)^d < \infty.$$

This proves the lemma with $K_{\rho,a} = C_\rho(1)C_{\rho,a}$. $\square$

$\langle \text{T:Solve} \rangle$ **Theorem 3.5** (Theorem 1.2 [CH19]). *Suppose that*

(i) *the initial measure μ satisfies the following integrability condition:*

$$\int_{\mathbb{R}^d} \exp(-a|x|^2) \|\mu(dx)\|_2 < \infty \quad \text{for all } a > 0, \quad (3.5) \quad \langle \text{A:icon} \rangle$$

(ii) *the spectral measure $\hat{f}$ satisfies Dalang's condition, namely,*

$$\Upsilon(\beta) := (2\pi)^{-d} \int_{\mathbb{R}^d} \frac{\hat{f}(d\xi)}{\beta + |\xi|^2} < \infty \quad \text{for some and therefore any } \beta > 0. \quad (3.6) \quad \langle \text{A:dalcon} \rangle$$

then (1.1) has a unique random field solution starting from μ . Moreover, the solution is $L^2(\Omega)$ continuous and is adapted to the filtration induced by W .

$\langle \text{T:ExUn} \rangle$ **Theorem 3.6** (Theorem 1.7 [CH19]). *Under the assumptions of Theorem 3.5, for any $t > 0$ and $x \in \mathbb{R}^d$, the solution to (1.1), $u(t, x)$, given by (1.3) is in $L^p(\Omega)$ and for $p \geq 2$*

$$\|u(t, x)\|_p \leq [\bar{\varsigma} + \sqrt{2}(G(t, \cdot) * |\mu|)(x)] H(t; \gamma_p)^{1/2}, \quad (3.7) \quad \langle \text{E:mb} \rangle$$

where $\bar{\varsigma} = L_0/L_b$, $\gamma_p = 32pL_b^2$ (see (1.4) for L_0 and L_b) and the function $H(t; \gamma_p)$ is nondecreasing in t (see [CH19] for the expression of H). Moreover, if

$$\Upsilon(0) < \infty \quad \text{and} \quad \gamma_p < \frac{1}{4\Upsilon(0)}, \quad (3.8) \quad \langle \text{A:dalcon0} \rangle$$

then this function $H(t; \gamma_p)$ is bounded in t . Hence, by choosing $C = (\sqrt{2} \vee \bar{\varsigma}) \sup_{t \geq 0} H(t; \gamma_p)^{1/2} < \infty$, we have that

$$\|u(t, x)\|_p \leq C \left(1 + (G(t, \cdot) * |\mu|)(x) \right). \quad (3.9) \quad \langle \text{E:mb0} \rangle$$

Corollary 3.7. *Under the assumption (3.8) above, the solution $v = v_{t_0}$ to the restarted equation (1.12) satisfies the following moment bound for any $q \geq 2$:*

$$\|v(t, x)\|_q \leq C \left(1 + (G(t + t_0, \cdot) * |\mu|)(x) \right), \quad \forall t \geq 0, \quad (3.10) \quad \boxed{\text{E:MomBddRestart}}$$

where $C > 0$ does not depend on t .

Proof. The proof is a simple consequence of Theorem 3.6. Indeed, for each $t \geq 0$

$$\|v(t, x)\|_q = \|u(t + t_0, x)\|_q \leq C \left(1 + (G(t + t_0, \cdot) * |\mu|)(x) \right).$$

□

The following factorization property of the heat kernel will be used in this paper

$$G(t, x)G(s, y) = G\left(\frac{ts}{t+s}, \frac{sx+ty}{t+s}\right) G(t+s, x-y), \quad (3.11) \quad \boxed{\text{E:gsim}}$$

which can be easily verified.

4 Proof of part (a) of Theorem 1.4

$\langle \text{S:mainproof} \rangle$ Now, for $\alpha \in (0, 1)$, $t > 0$ and $x \in \mathbb{R}^d$, define formally

$$\begin{aligned} (F_\alpha f)(t, x) &:= \int_0^t \int_{\mathbb{R}^d} (t-s)^{\alpha-1} G(t-s, x-y) f(s, y) \, ds dy \quad \text{and} \\ (Y_\alpha f)(t, x) &:= \int_0^t \int_{\mathbb{R}^d} (t-s)^{-\alpha} G(t-s, x-y) f(s, y) W(ds, dy). \end{aligned} \quad (4.1) \quad \boxed{\text{?E:Ops?}}$$

For F_α , the Step 1 of the proof of [TZ98, Theorem 3.1] showed the following proposition:

$\langle \text{P:F-Comp} \rangle$ **Proposition 4.1.** *Let ρ and $\hat{\rho}$ be given in condition (i) of Theorem 1.4 (see (1.17)). For any $q > 2$, $t_0 > 0$ and $\alpha \in (q^{-1}, 2^{-1})$, the operator F_α as an operator from $L^q((0, t_0); L_\rho^2)$ to $L_\rho^2(\mathbb{R}^d)$ is compact.*

As for Y_α , we have the following result:

$\langle \text{L:yineq} \rangle$ **Lemma 4.2.** *For any $\alpha \in (0, 1/2)$, denote (with slight abuse of notation)*

$$Y_{v_{t_0}}(s, t) = Y_v(s, y) := [Y_\alpha b(\circ, v(\cdot, \circ))](s, y) = \int_0^s \int_{\mathbb{R}^d} (s-r)^{-\alpha} G(s-r, y-z) b(z, v(r, z)) W_{t_0}(dr, dz). \quad (4.2) \quad \boxed{\text{E:Y}}$$

Under the assumptions of Theorem 1.3 and (iii) of Theorem 1.4, then for all $q \geq 2$, $s > 0$ and $y \in \mathbb{R}^d$,

$$\|Y_v(s, y)\|_q^2 \leq C_q [J_u^*(s + t_0, y)]^2 \mathcal{H}_\alpha(s). \quad (4.3) \quad \boxed{\text{E:NormY}}$$

Moreover, if

(iv') for some $q \geq 2$ and $\alpha \in (0, 1/2)$, the weight function ρ (not necessarily admissible)

$$\int_0^t [\mathcal{G}_\rho^u(s + t_0) \mathcal{H}_\alpha(s)]^{q/2} ds < \infty, \quad \text{for all } t > 0, \quad (4.4) \quad \boxed{\text{E:GH-int'}}$$

then there exists a constant $\Theta = \Theta(q, L_b, L_0, \alpha)$ such that

$$\mathbb{E} \left(\int_0^t \|Y_v(s, \cdot)\|_\rho^q ds \right) \leq \Theta \int_0^t [\mathcal{G}_\rho^u(s+t_0) \mathcal{H}_\alpha(s)]^{q/2} ds < \infty. \quad (4.5) \quad \boxed{\text{E:yineq}}$$

Proof. In the proof, we use C to denote a generic constant that may change its value at each appearance. First, an application of Minkowski's inequality shows that

$$\mathbb{E} \left(\|Y_v(s, \cdot)\|_\rho^q \right) = \left\| \int_{\mathbb{R}^d} Y_v(s, y)^2 \rho(y) dy \right\|_{q/2}^{q/2} \leq \left(\int_{\mathbb{R}^d} \|Y_v(s, y)\|_q^2 \rho(y) dy \right)^{q/2}. \quad (4.6) \quad \boxed{\text{E:EYqr}}$$

Now we look to find an appropriate upper bound for $\|Y(s, y)\|_q^2$. By the Burkholder-Davis-Gundy inequality, as well as Minkowski's integral inequality to see that

$$\begin{aligned} \|Y_v(s, y)\|_q^2 &\leq C_q \int_0^s dr (s-r)^{-2\alpha} \iint_{\mathbb{R}^{2d}} dz_1 dz_2 G(s-r, y-z_1) \|b(z_1, v(r, z_1))\|_q f(z_1-z_2) \\ &\quad \times G(s-r, y-z_2) \|b(z_2, v(r, z_2))\|_q. \end{aligned}$$

Note that for the Lipschitz condition in (1.4), we have that

$$|b(x, v)| \leq |b(x, v) - b(x, 0)| + |b(x, 0)| \leq L_b |v| + L_0 \leq C(1 + |v|), \quad C := L_b \vee L_0.$$

We apply this and the moment bound (3.10) to $\|b(z_i, v(r, z_i))\|_q$ above to see that

$$\|b(z_i, v(r, z_i))\|_q \leq C \left(1 + \|v(r, z_i)\|_q \right) \leq C \left(1 + (G(r+t_0, \cdot) * |\mu|)(z_i) \right) = C J_u^*(r+t_0, z_i), \quad i = 1, 2, \quad (4.7) \quad \boxed{\text{E:NormB}}$$

where J_u^* is defined in (1.14). Therefore,

$$\begin{aligned} \|Y_v(s, y)\|_q^2 &\leq C \int_0^s dr (s-r)^{-2\alpha} \iint_{\mathbb{R}^{2d}} dz_1 dz_2 f(z_1-z_2) \prod_{i=1}^2 \left(G(s-r, y-z_i) J_u^*(r+t_0, z_i) \right) \\ &= C \int_0^s dr (s-r)^{-2\alpha} \iint_{\mathbb{R}^{2d}} \mu^*(d\sigma_1) \mu^*(d\sigma_2) \iint_{\mathbb{R}^{2d}} dz_1 dz_2 \\ &\quad \times f(z_1-z_2) \prod_{i=1}^2 \left(G(s-r, y-z_i) G(r+t_0, z_i-\sigma_i) \right) \\ &= C \int_0^s dr (s-r)^{-2\alpha} \iint_{\mathbb{R}^{2d}} \mu^*(d\sigma_1) \mu^*(d\sigma_2) G(s+t_0, y-\sigma_1) G(s+t_0, y-\sigma_2) \\ &\quad \times \iint_{\mathbb{R}^{2d}} dz_1 dz_2 f(z_1-z_2) \prod_{i=1}^2 G \left(\frac{(r+t_0)(s-r)}{s+t_0}, z_i - \sigma_i \frac{r+t_0}{s+t_0} - \frac{s-r}{s+t_0} y \right) \\ &\leq C(2\pi)^{-2d} \int_0^s dr (s-r)^{-2\alpha} \iint_{\mathbb{R}^{2d}} \mu^*(d\sigma_1) \mu^*(d\sigma_2) G(s+t_0, y-\sigma_1) G(s+t_0, y-\sigma_2) \\ &\quad \times \int_{\mathbb{R}^d} \widehat{f}(d\xi) \exp \left(-\frac{(r+t_0)(s-r)}{s+t_0} |\xi|^2 \right), \end{aligned}$$

where we have applied (3.11) and Plancherel's theorem. Therefore,

$$\|Y_v(s, y)\|_q^2 \leq C(2\pi)^{-2d} [J_u^*(s+t_0, y)]^2 \int_0^s dr (s-r)^{-2\alpha} \int_{\mathbb{R}^d} \widehat{f}(d\xi) \exp \left(-\frac{(r+t_0)(s-r)}{s+t_0} |\xi|^2 \right).$$

Notice the function

$$t_0 \mapsto \frac{r+t_0}{s+t_0} \quad \text{for } t_0 > 0,$$

is increasing in t_0 whenever $s > r > 0$. Thus we have that

$$\|Y_v(s, y)\|_q^2 \leq C(2\pi)^{-2d} [J_u^*(s+t_0, y)]^2 \int_0^s dr (s-r)^{-2\alpha} \int_{\mathbb{R}^d} \widehat{f}(d\xi) \exp\left(-\frac{r(s-r)}{s} |\xi|^2\right). \quad (4.8) \quad \boxed{\text{E_NormY}}$$

Furthermore, by symmetry of $r(s-r)/s$ and the fact that $r(s-r)/s \geq r/2$ for all $r \in [0, s/2]$, we see that the above double integral is bounded by

$$\begin{aligned} &\leq 2 \int_0^{s/2} dr r^{-2\alpha} \int_{\mathbb{R}^d} \widehat{f}(d\xi) \exp\left(-\frac{r}{2} |\xi|^2\right) \\ &= 2^{2(1-\alpha)} \int_0^{s/4} dr r^{-2\alpha} \int_{\mathbb{R}^d} \widehat{f}(d\xi) \exp(-r |\xi|^2) \\ &\leq 2^{2(1-\alpha)} \int_0^s dr r^{-2\alpha} \int_{\mathbb{R}^d} \widehat{f}(d\xi) \exp(-r |\xi|^2) \\ &= 2^{2(1-\alpha)} \mathcal{H}_\alpha(s). \end{aligned}$$

Plugging the above bound back to (4.8) proves (4.3).

It remains to prove (4.5). By the definition of $\mathcal{G}_\rho^u(t)$ in (1.15) and by (4.3), we see that

$$\int_{\mathbb{R}^d} \|Y_v(s, y)\|_q^2 \rho(y) dy \leq C \mathcal{G}_\rho^u(s+t_0) \mathcal{H}_\alpha(s).$$

Plugging the above expression to the far right-hand of (4.6) proves (4.5). Finally, the finiteness of the upper bound in (4.5) is guaranteed by condition (4.4). This completes the proof of Lemma 4.2. $\square$

(L:Factorize)

Lemma 4.3 (Factorization lemma). *Let $u(t, x, \mu)$ be the solution to (1.1) and Y_v be as in (4.2). Under the conditions of Theorem 1.3 and (iii) of Theorem 1.4, for all $\alpha \in (0, 1/2)$, the following factorization holds*

$$\frac{\sin(\alpha\pi)}{\pi} \int_0^t (t-s)^{\alpha-1} [G(t-s, \cdot) * Y_v(s, \cdot)](x) ds = \int_0^t \int_{\mathbb{R}^d} G(t-r, x-z) b(z, v(r, z)) W_{t_0}(dr, dz).$$

As a consequence,

$$v(t, x) = [G(t, \cdot) * u(t_0, \cdot, \mu)](x) + \frac{\sin(\alpha\pi)}{\pi} [F_\alpha Y_v](t, x), \quad \text{for all } t > 0 \text{ and } x \in \mathbb{R}^d. \quad (4.9) \quad \text{?E:1fac?}$$

Proof. The lemma is straightforward provided that one can switch the orders of stochastic and ordinary integrals:

$$\begin{aligned} &\int_0^t (t-s)^{\alpha-1} [G(t-s, \cdot) * Y_v(s, \cdot)](x) ds \\ &= \int_0^t ds (t-s)^{\alpha-1} \int_{\mathbb{R}^d} dy G(t-s, x-y) \\ &\quad \times \int_0^s \int_{\mathbb{R}^d} (s-r)^{-\alpha} G(s-r, y-z) b(z, v(r, z)) W_{t_0}(dr, dz) \\ &= \int_0^t ds (t-s)^{\alpha-1} \int_0^s \int_{\mathbb{R}^d} (s-r)^{-\alpha} G(t-r, x-z) b(z, v(r, z)) W_{t_0}(dr, dz) \end{aligned} \quad (4.10) \quad \boxed{\text{E:Fubini}_1}$$

$$\begin{aligned}
&= \int_0^t \int_{\mathbb{R}^d} W(dr, dz) G(t-r, x-z) b(v(r, z)) \int_r^t ds (s-r)^{-\alpha} (t-s)^{\alpha-1} \\
&= \frac{\pi}{\sin(\alpha\pi)} \int_0^t \int_{\mathbb{R}^d} G(t-r, x-z) b(v(r, z)) W_{t_0}(dr, dz),
\end{aligned} \tag{4.11} \quad \boxed{\text{E:Fubini_2}}$$

where the last step is the Beta integral which requires that $\alpha \in (0, 1)$. It remains to justify the two applications of the stochastic Fubini's theorem (see Theorem 5.30 of Chapter one in [Dal+09], or also [Wal86] or Theorem 4.18 of [DZ92]) in (4.10) and (4.11) in the following two steps.

Step 1. In this step, we justify the change of orders in (4.10). Note that t, x and s are fixed. It suffices to prove the following condition:

$$\begin{aligned}
I_1 &:= \int_{\mathbb{R}^d} dy G(t-s, x-y) \int_0^s dr (s-r)^{-2\alpha} \iint_{\mathbb{R}^{2d}} dz_1 dz_2 \\
&\quad \times f(z_1 - z_2) \left(\prod_{i=1}^2 G(s-r, y-z_i) \right) \mathbb{E} \left(\prod_{i=1}^2 b(z_i, v(r, z_i)) \right) \\
&= \int_{\mathbb{R}^d} dy G(t-s, x-y) \|Y_v(s, y)\|_2^2 \\
&< +\infty.
\end{aligned}$$

But this follows immediately from (4.3). Indeed

$$\begin{aligned}
\int_{\mathbb{R}^d} dy G(t-s, x-y) \|Y_v(s, y)\|_2^2 &\leq C \int_{\mathbb{R}^d} dy G(t-s, x-y) (J_u^*(s+t_0, y))^2 \mathcal{H}_\alpha(s) \\
&= C \mathcal{H}_\alpha(s) \int_{\mathbb{R}^d} dy G(t-s, x-y) (J_u^*(s+t_0, y))^2.
\end{aligned}$$

Then Condition (1.15) implies that $I_1 < \infty$ as $\rho(y)$ cannot decay as fast as $y \mapsto G(t-s, x-y)$ (See Remark 3.2).

Step 2. Similarly, as for (4.11), we need to show that

$$\begin{aligned}
I_2 &:= \int_0^t ds (t-s)^{\alpha-1} \int_0^s dr (s-r)^{-2\alpha} \iint_{\mathbb{R}^{2d}} dz_1 dz_2 \\
&\quad \times f(z_1 - z_2) \left(\prod_{i=1}^2 G(t-r, x-z_i) \right) \mathbb{E} \left(\prod_{i=1}^2 b(z_i, v(r, z_i)) \right) < \infty.
\end{aligned}$$

By the Cauchy Schwartz inequality, (4.7) and because $\alpha \in (0, 1/2)$,

$$\begin{aligned}
I_2 &\leq C \int_0^t ds (t-s)^{\alpha-1} \int_0^s dr (s-r)^{-2\alpha} \iint_{\mathbb{R}^{2d}} dz_1 dz_2 f(z_1 - z_2) \prod_{i=1}^2 \left(G(t-r, x-z_i) J_u^*(r+t_0, z_i) \right) \\
&= C' \int_0^t dr (t-r)^{-\alpha} \iint_{\mathbb{R}^{2d}} dz_1 dz_2 f(z_1 - z_2) \prod_{i=1}^2 \left(G(t-r, x-z_i) J_u^*(r+t_0, z_i) \right).
\end{aligned}$$

Now by the same arguments as those leading to (4.3) (with 2α there replaced by α), we see that

$$I_2 \leq C (J_u^*(t, x))^2 \int_0^t dr r^{-\alpha} \int_{\mathbb{R}^d} \widehat{f}(d\xi) \exp\left(-\frac{r(t-r)}{t} |\xi|^2\right),$$

which is finite by (1.19) where we replace α with $\alpha/2$ and repeat the same steps right after Equation (4.8). This completes the proof of Lemma 4.3. $\square$

We are now ready to prove part (a) of Theorem 1.4.

Proof of Theorem 1.4 (a). Recall that Lemma 4.3 implies that for any $t_0 > 0$,

$$v(t_0, x) = \left(G(t_0, \cdot) * u(t_0, \cdot, \mu) \right)(x) + \frac{\sin(\alpha\pi)}{\pi} [F_\alpha Y_v](t_0, x).$$

Now define the set $\mathcal{K}(\Lambda)$ as

$$\mathcal{K}(\Lambda) = \{ (G(t_0, \cdot) * y(\cdot))(x) + (F_\alpha h)(x), \quad \text{where } \|y\|_{\hat{\rho}} \leq \Lambda \text{ and } \|h\|_{L^q(0,1,L^2_{\hat{\rho}})} \leq \Lambda \}.$$

Then by Proposition 4.1, $\mathcal{K}(\Lambda)$ is relatively compact in $L^2_{\hat{\rho}}$ for $\Lambda > 0$. Moreover, by Chebyshev's inequality, (1.16) and (4.5) we have

$$\begin{aligned} \mathbb{P}\{v(t_0, \cdot) \notin \mathcal{K}(\Lambda)\} &\leq \mathbb{P}\left[\left(\int_0^{t_0} \|Y_v(s, \cdot)\|_{\hat{\rho}}^q ds\right)^{1/q} > \frac{\pi\Lambda}{\sin(\alpha\pi)}\right] + \mathbb{P}\left[\|u(t_0, \cdot, \mu)\|_{\hat{\rho}} > \Lambda\right] \\ &\leq \frac{\sin^q(\alpha\pi)}{\pi^q \Lambda^q} \mathbb{E} \int_0^{t_0} \|Y_v(s, \cdot)\|_{\hat{\rho}}^q ds + \frac{1}{\Lambda^2} \mathbb{E} \left(\|u(t_0, \cdot, \mu)\|_{\hat{\rho}}^2 \right) \\ &\leq \frac{\sin^q(\alpha\pi)}{\pi^q \Lambda^q} \Theta \int_0^{t_0} (\mathcal{G}_{\hat{\rho}}^u(s+t_0) \mathcal{H}_\alpha(s))^{q/2} ds + \frac{1}{\Lambda^2} \mathbb{E} \left(\|u(t_0, \cdot, \mu)\|_{\hat{\rho}}^2 \right) < \infty. \end{aligned} \tag{4.12} \text{E:Restartbdd}$$

With all this said, (4.12) implies that

$$\mathbb{P}\{v(t_0, \cdot) \in \mathcal{K}(\Lambda)\} \geq 1 - \frac{\sin^q(\alpha\pi)}{\pi^q \Lambda^q} \left(\Theta \int_0^{t_0} (\mathcal{G}_{\hat{\rho}}^u(s+t_0) \mathcal{H}_\alpha(s))^{q/2} ds \right) - \frac{1}{\Lambda^2} \mathbb{E} \left(\|u(t_0, \cdot, \mu)\|_{\hat{\rho}}^2 \right),$$

and we can choose $\Lambda > 0$ big enough such that

$$\frac{\sin^q(\alpha\pi)}{\pi^q \Lambda^q} \left(\Theta \int_0^{t_0} (\mathcal{G}_{\hat{\rho}}^u(s+t_0) \mathcal{H}_\alpha(s))^{q/2} ds \right) + \frac{1}{\Lambda^2} \mathbb{E} \left(\|u(t_0, \cdot, \mu)\|_{\hat{\rho}}^2 \right) < \epsilon,$$

thus we get that

$$\mathbb{P}\{v(t_0, \cdot) \in \mathcal{K}(\Lambda)\} \geq 1 - \epsilon.$$

We can now deduce from this and the semigroup property of the solution to the heat equation that for any $t > t_0$,

$$\mathbb{P}\{v(t, \cdot) \in \mathcal{K}(\Lambda)\} \geq (1 - \epsilon) \mathbb{P}\{\|v(t - t_0, \cdot)\|_{\hat{\rho}} < \Lambda\}. \tag{4.13} \text{E:vtight}$$

We now apply Chebyshev's inequality and (1.18) to see that for all $t > t_0$ that there exists a $C > 0$ such that

$$\mathbb{P}\{\|v(t - t_0, \cdot)\|_{\hat{\rho}} \geq \Lambda\} \leq \frac{\mathbb{E} \left(\|v(t - t_0, \cdot)\|_{\hat{\rho}}^2 \right)}{\Lambda^2} = \frac{\mathbb{E} \left(\|u(t, \cdot, \mu)\|_{\hat{\rho}}^2 \right)}{\Lambda^2} \leq C \frac{\mathcal{G}_{\hat{\rho}}^u(t)}{\Lambda^2} \leq \frac{C}{\Lambda^2}.$$

Thus, by enlarging Λ if necessary,

$$\mathbb{P}\{\|v(t - t_0, \cdot)\|_{\hat{\rho}} \geq \Lambda\} = \mathbb{P}\{\|u(t, \cdot, \mu)\|_{\hat{\rho}} \geq \Lambda\} \leq \epsilon, \quad \forall t \geq t_0.$$

Combining this with (4.13) yields

$$\mathbb{P}\{u(t, \cdot, \mu) \in \mathcal{K}(\Lambda)\} \geq (1 - \epsilon), \quad t \geq t'_0,$$

where $t'_0 = 2t_0$. Since $t_0 > 0$ is arbitrary, then this completes the proof of (1.21). $\square$

References

- [AM03] Sigurd Assing and Ralf Manthey. “Invariant measures for stochastic heat equations with unbounded coefficients”. In: *Stochastic Process. Appl.* 103.2 (2003), pp. 237–256 (cit. on pp. 2, 3).
- [CM94] René A. Carmona and S. A. Molchanov. “Parabolic Anderson problem and intermittency”. In: *Mem. Amer. Math. Soc.* 108.518 (1994), pp. viii+125 (cit. on p. 2).
- [CD15] Le Chen and Robert Dalang. “Moments and Growth Indices for the Nonlinear Stochastic Heat Equation with Rough Initial Conditions”. In: *Ann. Probab.* 43 (2015), pp. 3006–3051.
- [CH19] Le Chen and Jingyu Huang. “Comparison principle for stochastic heat equation on $\mathbb{R}^d$ ”. In: *Ann. Probab.* 47.2 (2019), pp. 989–1035 (cit. on p. 10).
- [CK19] Le Chen and Kunwoo Kim. “Nonlinear stochastic heat equation driven by spatially colored noise: moments and intermittency”. In: *Acta Math. Sci. Ser. B (Engl. Ed.)* 39.3 (2019), pp. 645–668 (cit. on p. 3).
- [DZ96] G. Da Prato and J. Zabczyk. *Ergodicity for infinite-dimensional systems*. Vol. 229. Cambridge University Press, Cambridge, 1996, pp. xii+339. ISBN: 0-521-57900-7 (cit. on pp. 2, 6).
- [DGZ92] Giuseppe Da Prato, D. Gatarek, and Jerzy Zabczyk. “Invariant measures for semi-linear stochastic equations”. In: *Stochastic Anal. Appl.* 10.4 (1992), pp. 387–408 (cit. on p. 3).
- [DZ92] Giuseppe Da Prato and Jerzy Zabczyk. *Stochastic equations in infinite dimensions*. Vol. 44. Cambridge University Press, Cambridge, 1992, pp. xviii+454. ISBN: 0-521-38529-6 (cit. on p. 14).
- [Dal+09] Robert Dalang et al. *A minicourse on stochastic partial differential equations*. Vol. 1962. Held at the University of Utah, Salt Lake City, UT, May 8–19, 2006, Edited by Khoshnevisan and Firas Rassoul-Agha. Springer-Verlag, Berlin, 2009, pp. xii+216. ISBN: 978-3-540-85993-2 (cit. on pp. 2, 14).
- [Gra14] Loukas Grafakos. *Modern Fourier analysis*. Third. Vol. 250. Springer, New York, 2014, pp. xvi+624. ISBN: 978-1-4939-1229-2; 978-1-4939-1230-8 (cit. on p. 7).
- [MSY16] Oleksandr Misiats, Oleksandr Stanzhytskyi, and Nung Kwan Yip. “Existence and uniqueness of invariant measures for stochastic reaction-diffusion equations in unbounded domains”. In: *J. Theoret. Probab.* 29.3 (2016), pp. 996–1026 (cit. on pp. 2, 3).
- [MSY20] Oleksandr Misiats, Oleksandr Stanzhytskyi, and Nung Kwan Yip. “Invariant measures for stochastic reaction-diffusion equations with weakly dissipative nonlinearities”. In: *Stochastics* 92.8 (2020), pp. 1197–1222 (cit. on pp. 2, 3).
- [Olv+10] Frank W. J. Olver et al. *NIST handbook of mathematical functions*. With 1 CD-ROM (Windows, Macintosh and UNIX). U.S. Department of Commerce, National Institute of Standards and Technology, Washington, DC; Cambridge University Press, Cambridge, 2010, pp. xvi+951. ISBN: 978-0-521-14063-8 (cit. on p. 8).
- [TZ98] Gianmario Tessitore and Jerzy Zabczyk. “Invariant measures for stochastic heat equations”. In: *Probab. Math. Statist.* 18.2, Acta Univ. Wratislav. No. 2111 (1998), pp. 271–287 (cit. on pp. 2–5, 9, 11).

- [Wal86] John B. Walsh. “An introduction to stochastic partial differential equations”. In: *École d’été de probabilités de Saint-Flour, XIV—1984*. Vol. 1180. 1986, pp. 265–439 (cit. on pp. 2, 14).