

Research Statement

1 Overview

I am currently a PhD student studying at Auburn University under the supervision of [Dr. Le Chen](#) and I will be defending my thesis in August 2022. Overall, my research deals with a space-time fractional stochastic partial differential equation driven by some type of Gaussian noise which varies depending on the problem in question. I will refer this equation as the *Interpolated Stochastic Heat and Wave Equation* (ISHWE),

$$\begin{cases} \left(\partial_t^\beta + \frac{\nu}{2}(-\Delta)^{\alpha/2} \right) u(t, x) = I_t^\gamma \left[\sigma(u(t, x)) \dot{W}(t, x) \right] & x \in \mathbb{R}^d, t > 0 \\ u(0, \cdot) = u_0 & \beta \in (0, 1] \\ u(0, \cdot) = u_0, \quad \partial_t u(0, \cdot) = v_0 & \beta \in (1, 2). \end{cases} \quad (1.1) \quad \boxed{\text{E:ISHWE}}$$

The ISHWE interpolates the well known *Stochastic Wave and Stochastic Heat Equations*, SWE and SHE respectively, in the sense that when we let $\alpha = \beta = 2$ and $\gamma = 0$ then we recover the SWE and when $\alpha = 2$ and $\beta = 1$ and $\gamma = 0$ then we recover the SHE. I have worked on problems that have dealt with proving the existence and uniqueness of a solution, deriving the exact blow-up time for a local solution, proving the existence of invariant measures of the solution, deriving exact moment asymptotics, and proving regularity of the solution.

2 Past Thesis Projects

2.1 Invariant Measures of the SHE

My first project dealt with proving the existence of an invariant measure to the Stochastic Heat Equation,

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) - \frac{1}{2} \Delta_x u(t, x) = \sigma(u(t, x)) \dot{W}(t, x) & x \in \mathbb{R}^d, t > 0 \\ u(0, x) = \mu(x), \end{cases} \quad (2.1) \quad \boxed{\text{E:SHE}}$$

where σ is globally Lipschitz satisfying

$$L_\sigma := \sup_{x \in \mathbb{R}^d} \frac{|\sigma(x)|}{|x|} < \infty$$

and the measure μ satisfies some integrability conditions given below in Theorem [2.1](#). The noise $\dot{W}$ is a Gaussian noise that is white in time and colored in space with spatial correlation function f . In formally speaking, this means

$$\mathbb{E} \left(\dot{W}(t, x) \dot{W}(t, z) \right) = \delta_0(t - s) f(x - z)$$

where δ_0 is the Dirac delta distribution. See [\[13, pg. 51\]](#) for more details.

The question of the existence of invariant measures in unbounded domains with dimension $d \geq 3$ has been handled in few papers such as [\[21, 1, 18, 19\]](#). One thing to note is that the just mentioned papers use the theory of the stochastic integral laid out by Da Prato and Jabczyk in [\[14\]](#) while in my work we used the theory developed by Walsh [\[22\]](#). The former develops a theory of integration with respect to Hilbert-space-valued processes while the later emphasizes

integration with respect to worthy martingale measures. The choice of either theory is one of mathematical convince as both have been proven to be equivalent [12].

To make a quick mention about the requirement, $d \geq 3$. Chen and Kim [8, Theorem 1.3] proved that a phase transition to the solution of (2.1) happens if and only if $d \geq 3$. In other words, in dimensions $d = 1$ and $d = 2$, the solution to (2.1) will exhibit exponential growth in time no matter how we control the coefficient $\rho(x)$. In order to prove the existence of an invariant measure, we will need to require that the second moments to the solution of (2.1) be bounded in t and this forces us to require $d \geq 3$.

Da Prato and Jabczyk have [16] established the following procedure

- Show the existence of a Markovian solution to the SPDE in question in a certain function space where the transition group satisfies the Feller Property
- Show that the solution starting from a particular initial condition is bounded in probability

Da Prato and Zabczyk [15] have also shown that the first bullet point is true for a wider class of SPDEs which includes (2.1). Zabczyk and Tessitore [21] follow this procedure to show the existence of an invariant measure for the laws of the solution solutions to (2.1) in two separate cases:

1. The initial measure is a function, $\varphi \in L_\rho^2 \cap L_{\hat{\rho}}^2$ where both ρ and $\hat{\rho}$ satisfy the condition that $\rho \cdot (\hat{\rho})^{-1} \in L^1(\mathbb{R}^d)$ and the solution is bounded in probability.
2. The initial measure is a constant function while assuming certain integrability properties of the spectral density.

By now using the theory of Walsh and a very important bound on $\mathbb{E}|u(t, x)|^2$ given by Chen [9], we are able to mimic the proof in [21], but in the language of Walsh, and show the existence of an invariant measure for a more broad group of initial conditions, which includes the Dirac delta distribution δ_0 .

InvMeasDelta

Theorem 2.1. Assume $d \geq 3$ and consider any pair of admissible weights ρ and $\hat{\rho}$ such that

$$\int_{\mathbb{R}^d} \frac{\rho(x)}{\hat{\rho}(x)} dx < \infty.$$

Also, suppose that the initial measure μ satisfies the following integrability condition:

$$\limsup_{t \geq 0} \int_{\mathbb{R}^d} dx \hat{\rho}(x) \left[\int_{\mathbb{R}^d} |\mu|(dy) G(t, x - y) \right]^2 < \infty, \quad (2.2)$$

A:InvAssumpt

and that the spectral measure $\hat{f}$ satisfies both

$$\Upsilon(0) := (2\pi)^{-d} \int_{\mathbb{R}^d} \frac{\hat{f}(d\xi)}{|\xi|^2} < \infty, \quad \text{and} \quad \int_0^s dr (s - r)^{-2\alpha} \int_{\mathbb{R}^d} \hat{f}(d\xi) \exp(-(s - r)|\xi|^2) < \infty. \quad (2.3)$$

A:InvAssumpt

Then the solution, $u(t, x)$, of (2.1) starting from μ satisfies

1. $u(t, \cdot) \in L_\rho^2(\mathbb{R}^d) \cap L_{\hat{\rho}}^2(\mathbb{R}^d)$
2. $u_{t_0}(t, x) := u(t + t_0, x)$ is bounded in probability in $L_\rho^2(\mathbb{R}^d)$

In return, there exists an invariant measure for the laws of the solution, $\{\mathcal{L}(u(t, \cdot))\}_{t \geq t_0}$, in $L_\rho^2(\mathbb{R}^d)$.

2.2 Solvability and Exact Moment Asymptotics for the ISHWE [7]

In this project, we studied the the ISHWE with constant one initial condition:

$$\begin{cases} (\partial_t^b + \frac{\nu}{2}(-\Delta)^{a/2}) u(t, x) = I_t^r [\sigma(u(t, x)) \dot{W}(t, x)] & x \in \mathbb{R}^d, t > 0 \\ u(0, \cdot) = 1 & \beta \in (0, 1] \\ u(0, \cdot) = 1, \quad \partial_t u(0, \cdot) = 0 & \beta \in (1, 2) \end{cases} \quad (2.4) \quad \text{E:ISHWEone}$$

where $a \in (0, 2]$, $b \in (0, 2)$, $r \geq 0$, $\nu > 0$ and $\theta > 0$. The fractional operators ∂_t^b , $(-\Delta)^{a/2}$ and I_t^r respectively denote the Caputo derivative, the fractional Laplacian and the Riemann Liouville fractional integral. The noise $\dot{W}$ is a time-independent multiplicative Gaussian noise, in other words, $\mathbb{E}[W(\phi)W(\psi)] = \int_{\mathbb{R}^d} \mathcal{F}\phi(\xi)\mathcal{F}\psi(\xi)\mu(d\xi)$, where μ is the spectral measure with density given as $\mu(d\xi) = \varphi(\xi)d\xi$, and $\varphi(\xi) = \prod_{i=1}^k C(\alpha_i, d_i)|\xi_{(i)}|^{d_i-\alpha_i}$. We now define the mild solution to (2.4).

Definition 2.2 (Mild, local and global solutions). (1) For $T \in (0, \infty]$, a random field $u = \{u(t, x) : t \in (0, T), x \in \mathbb{R}^d\}$ is called a *mild solution* to the equation (2.4) if for all $x \in \mathbb{R}^d$ and s, t fixed with $0 < s \leq t < T$, $y \rightarrow G(t-s, x-y)u(s, y)$ is Skorohod integrable and the following stochastic integral equation holds almost surely

$$u(t, x) = 1 + \sqrt{\theta} \int_0^t \left(\int_{\mathbb{R}^d} G(t-s, x-y)u(s, y)W(\delta y) \right) ds, \quad (2.5) \quad \text{E:IntEq}$$

where G is the Green's function of the deterministic version of (2.4), ie when $\sigma(u(t, x)) \dot{W}(t, x)$ is replaced by a nice deterministic function.

(2) Let $u(t, x)$ be a mild solution to (2.4) (or (2.5)) and fix $p \geq 1$. We call $u(t, x)$ a *global $L^p(\Omega)$ -solution*, or simply an *$L^p(\Omega)$ -solution* if $\|u(t, x)\|_p < \infty$ for all $t > 0$ and $x \in \mathbb{R}^d$.

(3) If there exist $0 < T_1 \leq T_2 < \infty$ such that $\|u(t, x)\|_p$ is finite for all $t \in (0, T_1)$ and $x \in \mathbb{R}^d$, but $\|u(t, x)\|_p$ diverges to infinity whenever $t > T_2$, the mild solution $u(t, x)$ in this case is called a *local $L^p(\Omega)$ -solution*.

Because of the constant one initial condition, we are able to apply a Piccard iteration scheme to suggest that $u(t, x)$ and $\mathbb{E}(|u(t, x)|^2)$ should be given as the following:

$$u(t, x) = 1 + \sum_{k \geq 1} \theta^{k/2} I_k(f_k(\cdot, x, t)), \quad \text{and} \quad \mathbb{E}(|u(t, x)|^2) = \sum_{k \geq 0} \theta^k k! \left\| \tilde{f}(\cdot, x, t) \right\|_{\mathcal{H}^{\otimes k}}^2, \quad (2.6) \quad \text{E:ISHWEoneS}$$

where $I_k : \mathcal{H}^{\otimes k}((\mathbb{R}^d)^k) \rightarrow \mathcal{H}_k$ is the k^{th} order Skorohod integral and $\mathcal{H}_k$ is the k^{th} Wiener chaos space and the kernels $f_k(\cdot, x, t)$ are obtained through the iteration to be

$$f_n(x_1, \dots, x_k; x, t) = \int_0^t \int_0^{t_n} \cdots \int_0^{t_2} G(t-t_n, x-x_n) \cdots G(t_2-t_1, x_2-x_1) dt_1 \cdots dt_n.$$

This is made legitimate in the next Theorem.

Theorem 2.3 ([7, Theorem 1.6]). *We assume some mild assumptions on the parameters of (2.4) which guarantees the non-negativity of $G(t, x)$. This assumption does not exclude the SHE and SWE.*

(1) (2.4) has a unique (global) solution $u(t, x)$, given in (2.6), in $L^p(\Omega)$ for all $p \geq 2$, $t > 0$, and $x \in \mathbb{R}^d$ provided that

$$0 < \alpha < \min \left(\frac{a}{b} [2(b+r) - 1], 2a, d \right). \quad (2.7) \quad \text{E:alpha}$$

(2) Otherwise, if

$$r \in [0, 1/2) \quad \text{and} \quad 0 < \alpha = \frac{a}{b} [2(b+r) - 1] \leq d, \quad (2.8) \quad \text{E:critical}$$

then (2.4) has a local solution in the sense that

(2-i) For any $p \geq 2$, (2.4) has a unique solution $u(t, x)$ in $L^p(\Omega)$ for all $p \geq 2$ and $x \in \mathbb{R}^d$, but only for $t \in (0, T_p)$ where

$$T_p := \frac{\nu^{\alpha/a}}{2\theta(p-1)\mathcal{M}_a^{(2a-\alpha)/a}}. \quad (2.9) \quad \boxed{\text{E:Tp}}$$

(2-ii) For any $t > T_2$, the series defining $\mathbb{E}(|u(t, x)|^2)$ in (2.6) diverges, that is, the $L^2(\Omega)$ -solution $u(t, x)$ to (2.4) does not exist whenever $t > T_2$.

The constant $\mathcal{M}_a$ is defined in [7, Equation (3.11)] and is shown to be finite in Theorem 3.5 (*ibid.*). The second main result of this project deals with finding an exact formula the asymptotic behavior of the moments of the solution.

Theorem 2.4 ([7, Corollary 1.8]). *Under the assumptions of Theorem 2.3 and if we assume (2.7), then*

1. For all $p \geq 2$ fixed, it holds that

$$\begin{aligned} \lim_{t \rightarrow \infty} t^{-\beta} \log \mathbb{E}(|u(t, x)|^p) = & p(p-1)^{\frac{1}{2(b+r) - \frac{b\alpha}{a} - 1}} \left(\frac{1}{2}\right) \left(\frac{2a}{2a(b+r) - b\alpha}\right)^{\beta} \\ & \times \left(\theta \nu^{-\alpha/a} \mathcal{M}_a^{\frac{2a-\alpha}{a}}\right)^{\frac{a}{2a(b+r) - b\alpha - a}} \left(2(b+r) - \frac{b\alpha}{a} - 1\right); \end{aligned} \quad (2.10) \quad \boxed{\text{E:momentasy}}$$

2. For all $t > 0$ fixed, it holds that

$$\begin{aligned} \lim_{p \rightarrow \infty} p^{-\beta} \log \mathbb{E}(|u(t, x)|^p) = & t^{\beta} \left(\frac{1}{2}\right) \left(\frac{2a}{2a(b+r) - b\alpha}\right)^{\beta} \\ & \times \left(\theta \nu^{-\alpha/a} \mathcal{M}_a^{\frac{2a-\alpha}{a}}\right)^{\frac{a}{2a(b+r) - b\alpha - a}} \left(2(b+r) - \frac{b\alpha}{a} - 1\right). \end{aligned} \quad (2.11) \quad \boxed{\text{E:momentasy}}$$

It is important to know that under the case of the SWE, ie $a = b = 2$ and $r = 0$ then our results recover the work of Balan *et al* [5] and under the case of the SHE, ie $a = 2$, $b = 1$ and $r = 0$ our results recover the work of Chen [6, Theorem 1.1 and Theorem 1.2]] when we set $\alpha_0 = 0$.

3 Current Thesis Project

3.1 Solvability of the ISHWE With Superlinear Diffusion Term

My current project deals with proving the existence and uniqueness a of global solution to (1.1) with a super-linear diffusion term σ that satisfies the following growth condition: as $|x| \rightarrow \infty$,

$$|\sigma(x)| \leq \sigma_1 + \sigma_2 |x| [\ln(|x|)]^{\delta}, \quad (3.1) \quad \boxed{\text{C:SigSupLin}}$$

where $\sigma_1 = c(\sigma) = \sigma(0)$ and $\delta > 0$. The work is an extension of Millet and Sanz-Solé [20]. It is often the case that blow of the solution to SPDE's will occur when considering a super-linear diffusion coefficient. For example, Asogwa *et al* [2] study (1.1) where $\beta \in (0, 1]$, $\alpha \in (0, 2)$, and $\gamma = 1 - \beta$. They prove in Theorem 1.2 that blow up will occur in finite time when the diffusion coefficient has super linear growth and is bounded below by a non-zero constant. The techniques in their paper can also be easily adapted (1.1) for $\beta \in (0, 2)$, $\alpha \in (0, 2)$ and $\gamma > 0$.

In this project, we require the initial conditions, $u_0 = 1$ and $v_0 = 0$, and the noise $\dot{W}$ may either be space-time white or white in time and colored in space. When the noise is colored in space, we take the correlation function to be the Riesz kernel. As of now, I have more or less

completed the case of space-time white noise and the same techniques should also apply to the colored noise case as well.

I will now outline the steps which prove the existence of a global solution to (1.1) driven by space-time white noise. We first consider the case where the diffusion coefficient σ is globally Lipschitz. In this case, existence and uniqueness is proven in [10, Theorem 3.1]. The next step was to prove the Proposition 3.1 stated below. We note that this Hölder regularity proven in this result recovers [17, Theorem 1.9].

P:Holder

Proposition 3.1. *Suppose that $u_0 = 1$ and $v_0 = 0$ and $\Theta := 2\alpha + \frac{\alpha}{\beta} \min\{2\gamma - 1, 0\}$. Then u has a version, still denoted by u , that is η_1 -Hölder continuous in time and η_2 -Hölder continuous in space with $0 < \eta_1 < q/2$ and $0 < \eta_2 < \theta/2$ where $0 < \theta < (\Theta - 1) \wedge 2$ and*

1. $0 < q \leq \rho$ under when $0 < \beta \leq 1$ and $\gamma \leq \lceil \beta \rceil - \beta$,
2. $0 < q < \rho$ under $0 < \alpha < 2$ and $0 < \beta < 2$.

Moreover, if we consider

$$a > (8L^2(\sigma)K^2)^{1/\rho} \quad \text{and} \quad p \in \left[2, \frac{a^\rho}{4L^2(\sigma)K^2}\right],$$

where K is a specific constant, whose definition at the moment is unimportant, then we have that for any $R > 0$,

$$\mathbb{E} \left[\sup_{t \in [0, T], |x| \leq R} |u(t, x)|^p \right] \leq 2^{p-1} + C(p, \theta, T, R) \left[\mathcal{M}_1^p + \mathcal{M}_2^p e^{apT} \left(2 + \frac{c(\sigma)}{L(\sigma)} \right)^p \right].$$

After proving this proposition, we can consider proving the existence of a global solution to (1.1) with a super-linear diffusion coefficient satisfying (3.1). For simplicity, suppose $d = 1$. What we do is truncate σ in following sense:

$$\sigma_N(x) = \sigma(x) \mathbf{1}_{\{|x| \leq N\}} + \sigma(N) \mathbf{1}_{\{x > N\}} + \sigma(-N) \mathbf{1}_{\{x \leq -N\}}.$$

This can also be done analogously in higher dimensions. For each N , σ_N is globally Lipschitz and so for each N we have the solution $u_N(t, x)$ where u_N is the solution to (1.1) with σ replaced by σ_N . If we define τ_N to be the following stopping time,

$$\tau_N := \inf \left\{ t > 0 : \sup_{|x| \leq R} |u_N(t, x)| \geq N \right\} \wedge T, \quad (3.2)$$

then through a bit of analysis we deduce that $\tau_N \uparrow \infty$ as $N \rightarrow \infty$ and that on $\{t < \tau_N\}$, $u_N(t, x) = u_{N+k}(t, x)$ for any $k \in \mathbb{N}$. Lastly, we define on each $\{t < \tau_N\}$, $(u(t, x) : (t, x) \in [0, T] \times \mathbb{R})$ by $u(t, x) = u_N(t, x)$. A little more analysis shows that u is a global solution to (1.1). Thus we have our next theorem.

Theorem 3.2. *Consider the initial conditions $u_0 = 1$ and $v_0 = 0$ and suppose that σ satisfies (3.1). Then for any $M > 0$ and $T > 0$, there exists a random field solution to (1.1) denoted as $\{u(t, x) : (t, x) \in [0, T] \times B_M(0)\}$. This solution satisfies*

$$\sup_{t \in [0, T], |x| \leq M} |u(t, x)| < \infty, \text{ a.s.} \quad (3.3)$$

One last thing to note here is that the solution u constructed in the above theorem is not defined in $L^2(\Omega)$. In order to define u , we use an extension of the Walsh integral which can be found in [11]

4 Future Projects

In this section I will introduce four possible future projects.

4.1 Sharp Moment Bounds for the Solution to the ISHWE Driven by a Spatially Colored Gaussian Noise

This project will try to generalize the following lemma to the ISHWE.

Lemma 4.1 ([9, Lemma 2.2]). *Suppose $\alpha = 2$, $\beta = 1$, $\gamma = 0$ and $\sigma(x) = \lambda x$ in (1.1). Suppose that the measure μ is such that $\int_{\mathbb{R}^d} \exp(-a|x|^2) |\mu|(dx) < \infty$. Denote $J_0(t, x)$ as the solution to the deterministic homogeneous SHE. Let $g : [0, \infty) \times \mathbb{R}^d \rightarrow [0, \infty)$ satisfy*

$$\int_0^t ds \iint_{\mathbb{R}^{2d}} dy_1 dy_2 G(t-s, x-y_1) G(t-s, x-y_2) f(y_1-y_2) g(s, y_1) g(s, y_2) < \infty.$$

In addition suppose,

$$g^2(t, x) \leq J_0^2(t, x) + \lambda^2 \int_0^t ds \iint_{\mathbb{R}^{2d}} dy_1 dy_2 G(t-s, x-y_1) G(t-s, x-y_2) f(y_1-y_2) g(s, y_1) g(s, y_2).$$

Then,

$$g(t, x) \leq (|\mu| * G(t, \cdot))(x) H(t; 2\lambda^2)^{1/2}.$$

There are two issues that arise. First is the loss of the following identity for the fundamental solution of the SHE:

$$G(s, x) G(t-s, y) = G\left(\frac{s(t-s)}{t}, \frac{sy - (t-s)x}{t}\right) G(t, x+y).$$

Second is the spatially colored noise. Because of the noise, we have that

$$\mathbb{E}(|u(t, x)|^2) = J_0^2(t, x) + \lambda^2 \int_0^t ds \iint_{\mathbb{R}^{2d}} dy_1 dy_2 G(t-s, x-y_1) G(t-s, x-y_2) f(y_1-y_2) \mathbb{E}[u(s, y_1) u(s, y_2)].$$

In the space-time white noise case, we have that $f(x-y) = \delta_0(x-y)$, which simplifies the expression above.

4.2 Solvability of the ISHWE With Superlinear Diffusion Term Driven by a Spatially Colored Gaussian Noise

This project will be a natural extension to my current project, which was explained in Section 3.1 above. The goal will be to prove the following results.

Theorem 4.2. *For $p \geq 2$, $T > 0$ and $R > 0$, there exists a series of constants K_1 , K_2 and K_3 such that*

$$\frac{\|u(t, x) - u(s, z)\|}{|x - z|^{v_1} + |t - s|^{v_2}} \leq K_1 (K_2 + K_2 \exp(Ta) \mathcal{N}_{a,p}(u))$$

where v_1 and v_2 are to be determined and $\mathcal{N}_{a,p} = \sup_{t \geq 0, x \in \mathbb{R}^d} \exp(-at) \|u(t, x)\|_p$.

Theorem 4.3. *Consider the initial conditions $u_0 = 1$ and $v_0 = 0$ and suppose that σ satisfies (3.1). Then for any $M > 0$ and $T > 0$, there exists a random field solution to (1.1) driven by a spatially colored noise denoted as $\{u(t, x) : (t, x) \in [0, T] \times B_R(0)\}$. This solution satisfies*

$$\sup_{t \in [0, T], |x| \leq R} |u(t, x)| < \infty, a.s. \quad (4.1)$$

To overcome the introduction of the spatially correlation, we have the following lemma which will be useful.

L:intYfYbdd

Lemma 4.4. *Suppose that the spatial correlation function is given to be the Riesz Kernel, $f(x) = x^{-\lambda}$, and that $\lambda < 2\alpha$. Then there exists a constant $C(\beta, \gamma) =: C > 0$ such that for any $x, y \in \mathbb{R}^d$ we have that*

$$\int_{\mathbb{R}^{2d}} Y(t, x - z)Y(t, y - w)f(w - z) \, dzdw \leq CK_f \left(\frac{\nu}{2}\right)^{-\lambda/\alpha} t^{2(\beta+\gamma-1)-\beta\lambda/\alpha}. \quad (4.2) \quad \text{E:YYfBdd}$$

Using this lemma we will be able to prove an analogous result as [20, Proposition 4.4] which was key to complete the project in the space-time white noise case.

4.3 ISHWE Driven by a Lévy Noise

This project will aim to generalize the recent results due to Balan [3] to the ISHWE. The goal will be to show existence of the solution of (1.1) driven by a Lévy white noise, $L(A)$, defined as follows:

$$L(A) = b|A| + \int_{A \times \{|z| \leq 1\}} z \tilde{J}(dt, dx, dz) + \int_{A \times \{|z| > 1\}} z J(dt, dx, dz)$$

where $A \in \mathcal{B}_0(\mathbb{R}_+ \times \mathbb{R}^d)$ is the class of finite Lebesgue-measured sets, $|\cdot|$ denotes the Lebesgue measure and $b \in \mathbb{R}$. J is a Poisson random measure on $\mathbb{R}_+ \times \mathbb{R}^d \times \mathbb{R}$ and $\tilde{J}$ is the compensated version of J .

Under the assumption that there exists $0 < q \leq p$ such that

$$\gamma_1 := \int_{|z| \leq 1} |z|^p \nu(dz) < \infty, \quad \text{and} \quad \gamma_2 := \int_{|z| > 1} |z|^q \nu(dz) < \infty,$$

Balan [3] proved the following result for the SWE.

Theorem 4.5. *Assume that γ_1 and γ_2 are finite and if $d = 2$ then $p < 2$. Let $h(x) = 1 + |x|^\eta$ for some $d/q < \eta < (2 - d(p - 1))/(p - q)$. Then the truncated SWE driven by a truncated Lévy measure,*

$$L_N(A) := b|A| + \int_{A \times \{|z| \leq 1\}} z \tilde{J}(dt, dx, dz) + \int_{A \times \{1 < |z| \leq Nh(x)\}} z J(dt, dx, dz),$$

has a unique solution denoted as $\{u_N(t, x) : t \geq 0, x \in \mathbb{R}^d\}$. Moreover,

$$\sup_{0 \leq t \leq T} \sup_{|x| \leq R} \mathbb{E}|u_N(t, x)|^p < \infty,$$

for any $T > 0$ and $R > 0$. In $d = 1$, the above supremum holds for $p \leq 2$.

4.4 Spatial Integral of the Solution of the ISHWE Driven by a Time-Independent Noise

This project will aim to generalize the recent results due to Balan and Yuan [4] to the ISHWE. In this project we will consider (1.1) driven by a time-independent Gaussian noise as was done in my previous work with Dr. Le Chen in [7]. We will need to consider the constant 1 initial condition with 0 initial velocity: $u_0 = 1$ and $v_0 = 0$. We also take $\sigma(u) = u$. As was proven in [7, Theorem 1.6 and 3.3], under these conditions, there exists a unique solution that can be expressed through the following Wiener expansion:

$$u(t, x) = 1 + \sum_{k \geq 1} I_k(f_k(\cdot, x, t)),$$

where $B_R(0)$ is the ball centered at the origin. We will want to investigate the quantity

$$F_R(t) := \int_{B_R(0)} u(t, x) - 1 \, dx.$$

Balan and Yuan [4] proved the following theorem, which will be the main focus of this future project.

Theorem 4.6 ([4, Theorem 1.3]). *If $d = 1$, then we take the noise to be white. If $d > 1$, then the noise will be colored with correlation function, $\gamma \in L^1(\mathbb{R}^d)$ which satisfies Dalang's condition. We have the following:*

1. for any $t > 0$ and $s > 0$,

$$\mathbb{E}[F_R(t)F_R(s)] \sim K(t, s)R^d \text{ as } R \rightarrow \infty, \text{ where } K(t, x) := \omega_d \int_{\mathbb{R}^d} \rho_{t,s}(z) dz < \infty,$$

and in particular, $\sigma_R^2(t) \sim K(t, t)R^d$ as $R \rightarrow \infty$ where $\sigma_R^2 = \mathbb{E}(|F_R(t)|^2)$ and $\rho_{t,s} := \mathbb{E}[(u(t, x) - 1)(u(s, y) - 1)]$,

2. for any $t > 0$,

$$d_{TV} \left(\frac{F_R(t)}{\sigma_R(t)}, Z \right) \leq C_t R^{-d/2},$$

where d_{TV} is the total variation distance and Z is a standard normal random variable,

3. there exists a continuous modification of the process $\{R^{-d/2}F_R(t)\}_{t \geq 0}$ which converges in distribution in $C[0, \infty)$ as $R \rightarrow \infty$, to a zero-mean Gaussian processes with covariance $K(t, s)$.

References

- AM03 [1] Assing, Sigurd and Ralf Manthey. Invariant measures for stochastic heat equations with unbounded coefficients. *Stochastic Process. Appl.* 103 (2003), no. 2, 237–256.
- AMN19 [2] Asogwa, M., Mijena, J., and Nane, E. (2019) Blow-Up Results for Space-Time Fractional Stochastic Partial Differential Equations. *Potential Analysis* (2020) 53: 357 – 386.
- Balan22 [3] Balan, R. Stochastic wave equation with Lévy white noise *arXiv:2111.14242*
- BY22 [4] Balan, R. and Yuan W. Spatial integral of the solution to hyperbolic Anderson model with time-independent noise *arXiv:2201.02319*
- BCC21 [5] Balan, R. M., Chen, L., and Chen, X. (2021). Exact asymptotics of the stochastic wave equation with time-independent noise. *Ann. Inst. Henri Poincaré Probab. Stat.*, to appear.
- Chen19AIHP [6] Chen X. (2019). Parabolic Anderson model with rough or critical Gaussian noise. *Ann. Inst. Henri Poincaré Probab. Stat.* **55**, 941–976.
- CE21 [7] Chen, L. and Eisenberg, N. (2021). Interpolating the Stochastic Heat and Wave Equations with Time-independent Noise: Solvability and Exact Asymptotics. *arXiv:2108.11473*. Accepted pending revision to *Stochastics and Partial Differential Equations: Analysis and Computation*.
- CK15SHE [8] Chen, Le and Kunwoo Kim. Nonlinear stochastic heat equation driven by spatially colored noise: moments and intermittency. *Acta Math. Sci. Ser. B*, 39B (2019), No. 3, 645–668.

- [9] Chen, Le and Jingyu Huang. Comparison principle for stochastic heat equation on $\mathbb{R}^d$. *Ann. Probab.* 47 (2019), no. 2, 989–1035.
- [10] Chen, L., Hu, Y., and Nualart, D. (2019). Nonlinear stochastic time-fractional slow and fast diffusion equations on $\mathbb{R}^d$. *Stochastic Process. Appl.* **129**, 5073–5112
- [11] Dalang, R. and Sanz-Solè, M. Stochastic Partial Differential Equations. Book in preparation.
- [12] Dalang, Robert C. and Lluís Quer-Sardanyons. Stochastic integrals for spde’s: a comparison. *Expo. Math.* 29 (2011), no. 1, 67–109.
- [13] Dalang, Robert, Davar Khoshnevisan, Carl Mueller, David Nualart, and Yimin Xiao. *A minicourse on stochastic partial differential equations*. Held at the University of Utah, Salt Lake City, UT, May 819, 2006. Edited by Khoshnevisan and Firas Rassoul-Agha. Lecture Notes in Mathematics, 1962. Springer-Verlag, Berlin, 2009. xii+216 pp.
- [14] Da Prato, Giuseppe and Jerzy Zabczyk. *Stochastic equations in infinite dimensions*. Encyclopedia of Mathematics and its Applications, 44. Cambridge University Press, Cambridge, 1992. xviii+454 pp.
- [15] Da Prato, Giuseppe and Jerzy Zabczyk. *Invariant measures for semilinear stochastic equations*. Stochastic Anal. Appl. 10, 387-408.
- [16] Da Prato, Giuseppe and Jerzy Zabczyk. *Ergodicity for infinite-dimensional systems*. London Mathematical Society Lecture Note Series, 229. Cambridge University Press, Cambridge, 1996.
- [17] Foondun, M and Nane E. (2016) Asymptotic properties of some space-time fractional stochastic equations. *Mathematische Zeitschrift* 2015, **287**, 493 – 519
- [18] Misiats, Oleksandr, Oleksandr Stanzhytskyi, and Nung-Kwan Yip. Ito’s Formula in infinite dimension and its application to the existence of invariant measures.
- [19] Misiats, Oleksandr, Oleksandr Stanzhytskyi, and Nung-Kwan Yip. Existence and uniqueness of invariant measures for stochastic reaction-diffusion equations in unbounded domains. *J. Theoret. Probab.* 29 (2016), no. 3, 996–1026.
- [20] Sanz-Sole, M. and Millet, A. (2021) Global solutions to stochastic wave equations with superlinear coefficients. *Stochastic Processes and their Applications*, Elsevier, 2021, 139, pp.175-211.
- [21] Tessitore, Gianmario and Jerzy Zabczyk. Invariant measures for stochastic heat equations. *Probab. Math. Statist.* 18 (1998), no. 2, Acta Univ. Wratislav. No. 2111, 271–287.
- [22] Walsh, John B. *An Introduction to Stochastic Partial Differential Equations*. In: École d’été de probabilités de Saint-Flour, XIV—1984, 265–439. Lecture Notes in Math. 1180, Springer, Berlin, 1986.