

1 Introduction

We study the interpolated stochastic heat and wave equation (ISHWE) for spatial dimension on $\mathbb{R}^d$,

$$\begin{cases} \left(\partial_t^\beta + \frac{\nu}{2}(-\Delta)^{\alpha/2} \right) u(t, x) = I_t^\gamma \left[\sigma(u(t, x)) \dot{W}(t, x) \right] & x \in \mathbb{R}^d, t > 0 \\ u(0, \cdot) = u_0 & \beta \in (0, 1] \\ u(0, \cdot) = u_0, \quad \partial_t u(0, \cdot) = v_0 & \beta \in (1, 2). \end{cases} \quad (1.1) \quad \boxed{\text{E:SPDE}}$$

2 Preliminaries

For any $t \in [0, T]$ and $x \in \mathbb{R}^d$, we write (assuming $\beta \in (1, 2)$)

$$u(t, x) = J_0(t, x) + I(t, x)$$

where

$$\begin{aligned} J_0(t, x) &= \begin{cases} [Z(t, \cdot) * u_0](x) & \beta \in (0, 1] \\ [Z(t, \cdot) * v_0](x) + \frac{\partial}{\partial t}[Z(t, \cdot) * u_0](x) & \beta \in (1, 2) \end{cases} \\ I(t, x) &= \int_0^t \int_{\mathbb{R}^d} Y(t-s, x-y) \sigma(u(s, y)) W(ds, dy). \end{aligned}$$

The function Y is defined through the Fox-H function and its Fourier transform can be shown to be given as

$$\mathcal{F}Y(t, \cdot)(\xi) = t^{\beta+\gamma-1} E_{\beta, \beta+\gamma}(-2^{-1} \nu t^\beta |\xi|^\alpha)$$

The following is for space-time white noise. For colored noise we need $\rho(\lambda) > 0$ and $2\alpha > d$.

3 Space-time White Noise

In this section we study (1.1) for driven by space-time white noise. To establish the existence and uniqueness of random field solutions in the to (1.1), we will need to assume Dalang's condition:

$$\int_0^t ds \int_{\mathbb{R}^d} dy |Y(s, y)|^2 < \infty, \quad \text{for all } t > 0, \quad (3.1) \quad \boxed{\text{C:Dalang}}$$

which is equivalent to

$$d < 2\alpha + \frac{\alpha}{\beta} \min\{2\gamma - 1, 0\} =: \Theta, \quad (3.2) \quad \boxed{\text{E:DalangEqu}}$$

which is further equivalent to

$$\rho(d) > 0 \quad \text{and} \quad d < 2\alpha, \quad (3.3) \quad \boxed{\text{E:DalangEqu}}$$

where

$$\rho(x) := 2\beta + 2\gamma - 1 - \beta x / \alpha. \quad (3.4) \quad \boxed{\text{D:theta}}$$

3.1 Globally Lipschitz Coefficient

In this section, we assume that the coefficient σ is globally Lipschitz and therefore satisfies the following inequality:

$$|\sigma(x)| \leq c(\sigma) + L(\sigma)|x|,$$

where $c(\sigma) = \sigma(0)$ and $L(\sigma)$ is the Lipschitz constant of σ . We assume that $L(\sigma) > 0$. We now state an existence and uniqueness result proven in [4].

Theorem 3.1. Under (3.1), the SPDE (1.1) has a unique (in the sense of versions) random field solution $\{u(t, x) : (t, x) \in (0, \infty) \times \mathbb{R}^d\}$ if the initial data are such that

$$\hat{C}_t := \sup_{t \in [0, T], x \in \mathbb{R}^d} |J_0(t, x)| < \infty. \quad (3.5)$$

Moreover, the following statements are true:

1. $(t, x) \mapsto u(t, x)$ is $L^p(\Omega)$ -continuous for all $p \geq 2$,
2. For all even integers $p \geq 2$, all $t \geq 0$ and $x, y \in \mathbb{R}^d$,

$$\|u(t, x)\|_p^2 \leq 2J_0^2(t, x) + \left[\varsigma^2 + 2\hat{C}_t^2\right] \exp(C\lambda^{2/(1-\sigma)}t). \quad (3.6)$$

Lemma 3.2. [4, Theorem 4.1 and Lemma 5.5] Assume that $d < 2\alpha$, $\beta \in (0, 2)$, and $\gamma \geq 0$. Then

$$\int_{\mathbb{R}^d} Y(t, x) dx = t^{\beta+\gamma-1}$$

and

$$\int_{\mathbb{R}^d} Y^2(t, x) dx = C_{\#} t^{2(\beta+\gamma-1)-d\beta/\alpha} = C_{\#} t^{\rho-1},$$

for all $t > 0$, where $\rho = \rho(d)$ and

$$C_{\#} := \frac{2}{\Gamma(d/2)(4\pi)^{d/2}} \int_0^\infty u^{d-1} E_{\beta, \beta+\gamma}^2(-u^\alpha) du.$$

Now further assume that $\beta \in (1, 2)$, then

$$\int_{\mathbb{R}} Z(t, x) dx = t^{\lceil \beta \rceil - 1} = t.$$

Remark 3.3. When $\alpha = \beta = 2$, $\gamma = 0$ and $d = 1$ then Lemma 3.2 gives us that $\int_{\mathbb{R}} G^2(t, x) dx = t/2$ which coincides with the $L^2(\mathbb{R})$ norm of the wave kernel.

In the proof of the next proposition, we will use the following two facts:

$$\sup_{t \geq 0} t^k e^{-at} = k^k (ea)^{-k} \text{ for } k \in \mathbb{N} \text{ and } \sup_{t \geq 0} \int_0^t s e^{-as} ds = a^{-2} \text{ for } a > 0. \quad (3.7)$$

E:SUPidenti

Proposition 3.4. [11, Proposition 3.2] Let u_0 and v_0 be Borel functions satisfying $\|u_0\|_\infty + \|v_0\|_\infty < \infty$. Suppose that $d < 2\alpha$ and $\rho = \rho(d) > 0$, where $\rho(x)$ is defined in (3.4). Then there exists a universal constant $K := K(\alpha, \beta, \gamma, d) > 0$ such that for any

$$a > (8L^2(\sigma)K^2)^{1/\rho}.$$

and

$$p \in \left[2, \frac{a^\rho}{4L^2(\sigma)K^2}\right].$$

we have that

$$N_{a,p}(u) \leq 2\mathcal{T}_0 + \frac{c(\sigma)}{L(\sigma)} \quad (3.8)$$

E:Nbdd

where

$$\mathcal{T}_0 = \begin{cases} (\lceil \beta \rceil - 1)^{\lceil \beta \rceil - 1} (ea)^{1 - \lceil \beta \rceil} \|u_0\|_\infty & \beta \in (0, 1] \\ (\lceil \beta \rceil - 1)^{\lceil \beta \rceil - 1} (ea)^{1 - \lceil \beta \rceil} \|v_0\|_\infty + \|u_0\|_\infty & \beta \in (1, 2). \end{cases}$$

Moreover for any $T > 0$,

$$\sup_{x \in \mathbb{R}^d} \mathbb{E}(|u(t, x)|^p) \leq \exp(apt) \left[2\mathcal{T}_0 + \frac{c(\sigma)}{L(\sigma)}\right]^p, \quad t \in [0, T]. \quad (3.9)$$

E:Pnormtbdd

Proof. First we fix an $a > 0$ and $p \in [2, \infty)$. Using Lemma 3.2 we see that,

$$\begin{aligned} |J(t, x)| &= \begin{cases} |(Z(t, \cdot) * u_0)(x)| & \beta \in (0, 1] \\ |(Z(t, \cdot) * v_0)(x) + \frac{\partial}{\partial t}(Z(t, \cdot) * u_0)(x)| & \beta \in (1, 2) \end{cases} \\ &\leq \begin{cases} t^{\lceil \beta \rceil - 1} \|u_0\|_\infty & \beta \in (0, 1] \\ t^{\lceil \beta \rceil - 1} \|v_0\|_\infty + \|u_0\|_\infty & \beta \in (1, 2) \end{cases} \end{aligned}$$

Now using (3.7), we see that

$$\begin{aligned} N_{a,p}(J) &\leq \begin{cases} \sup_{t \geq 0} \sup_{x \in \mathbb{R}} t^{\lceil \beta \rceil - 1} \|u_0\|_\infty e^{-at} & \beta \in (0, 1] \\ \sup_{t \geq 0} \sup_{x \in \mathbb{R}} (t^{\lceil \beta \rceil - 1} \|v_0\|_\infty + \|u_0\|_\infty) e^{-at} & \beta \in (1, 2) \end{cases} \\ &= \begin{cases} (\lceil \beta \rceil - 1)^{\lceil \beta \rceil - 1} (ea)^{1 - \lceil \beta \rceil} \|u_0\|_\infty & \beta \in (0, 1] \\ (\lceil \beta \rceil - 1)^{\lceil \beta \rceil - 1} (ea)^{1 - \lceil \beta \rceil} \|v_0\|_\infty + \|u_0\|_\infty & \beta \in (1, 2). \end{cases} \end{aligned} \quad (3.10) \quad \boxed{\text{E: Jbdd}}$$

Now we find an upper bound for $N_{a,p}(I)$. By simultaneously applying the Burkholder-Davies-Gundy inequality and triangle inequality, we get that

$$\begin{aligned} \|I(t, x)\|_p^2 &\leq 4p \left\| \int_0^t ds \int_{\mathbb{R}^d} dy Y^2(t-s, x-y) \sigma^2(u(s, y)) \right\|_{p/2} \\ &\leq 4p \int_0^t ds \int_{\mathbb{R}^d} dy Y^2(t-s, x-y) \|\sigma^2(u(s, y))\|_{p/2}. \end{aligned}$$

Now by applying Lemma 3.2 and the Lipschitz property of σ we get that

$$\|I(t, x)\|_p^2 \leq 8pc^2(\sigma)C_\# \int_0^t (t-s)^{\rho-1} ds + 8pL^2(\sigma)N_{a,p}^2(u) \int_0^t ds \int_{\mathbb{R}^d} dy e^{2as} Y^2(t-s, x-y).$$

Now under Dalang's condition (3.3), the first integral above can be calculated as

$$\int_0^t (t-s)^{\rho-1} ds = \frac{t^\rho}{\rho}.$$

By applying Lemma 3.2, we can calculate the second integral as

$$\int_0^t ds \int_{\mathbb{R}^d} dy e^{2as} Y^2(t-s, x-y) = C_\# \int_0^t ds e^{2as} (t-s)^{\rho-1}.$$

Putting this together gives us that

$$\|I(t, x)\|_p^2 \leq 8pc^2(\sigma)C_\# \frac{t^\rho}{\rho} + 8pL^2(\sigma)N_{a,p}^2(u)C_\# \int_0^t ds e^{2as} (t-s)^{\rho-1}.$$

Now by multiplying both sides by e^{-2at} , we see that

$$\begin{aligned} \|I(t, x)\|_p^2 e^{-2at} &\leq \frac{8pc^2(\sigma)C_\#}{\rho} t^\rho e^{-2at} \\ &\quad + 8pL^2(\sigma)N_{a,p}^2(u)C_\# \int_0^t ds e^{-2a(t-s)} (t-s)^{\rho-1}. \end{aligned}$$

We now do a change of variables and again recall the definition of the Gamma function to see that

$$\|I(t, x)\|_p^2 e^{-2at} \leq \frac{8pc^2(\sigma)C_\#}{\rho} t^\rho e^{-2at} + 8pL^2(\sigma)N_{a,p}^2(u)C_\# \int_0^\infty du e^{-2au} (u)^{\rho-1}$$

$$= \frac{8pc^2(\sigma)C_{\#}}{\rho} t^{\rho} e^{-2at} + 8pL^2(\sigma)N_{a,p}^2(u) \frac{C_{\#}\Gamma(\rho)}{(2a)^{\rho}}$$

where the last line is true if Dalang's condition [4] holds. If we now apply (3.7) and also recall that $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$, then we see that

$$N_{a,p}(I) \leq c(\sigma) \sqrt{\frac{8pC_{\#}}{\rho}} \left(\frac{\rho}{2ea}\right)^{\rho/2} + L(\sigma)N_{a,p}(u) \sqrt{\frac{8pC_{\#}\Gamma(\rho)}{(2a)^{\rho}}}. \quad (3.11) \quad \boxed{\text{E:SoleMil3.}}$$

To simplify things, we now let

$$K := \max \left\{ \sqrt{\frac{8C_{\#}}{\rho}} \left(\frac{\rho}{2e}\right)^{\rho/2}, \sqrt{\frac{8C_{\#}\Gamma(\rho)}{2^{\rho}}} \right\}$$

then,

$$N_{a,p}(I) \leq \frac{K\sqrt{p}}{a^{\rho/2}} (c(\sigma) + L(\sigma)N_{a,p}(u)). \quad (3.12) \quad \boxed{\text{E:Ibdd}}$$

Now by combining (3.10) and (3.12), we see that

$$N_{a,p}(u) \leq \begin{cases} ([\beta] - 1)^{[\beta]-1} (ea)^{1-[\beta]} \|u_0\|_{\infty} & \beta \in (0, 1] \\ ([\beta] - 1)^{[\beta]-1} (ea)^{1-[\beta]} \|v_0\|_{\infty} + \|u_0\|_{\infty} & \beta \in (1, 2). \end{cases} \quad (3.13)$$

$$+ K \frac{\sqrt{p}c(\sigma)}{a^{\rho/2}} + K \frac{\sqrt{p}L(\sigma)}{a^{\rho/2}} N_{a,p}(u). \quad (3.14) \quad \boxed{\text{E:Nbddgen}}$$

It is important to note that if we now restrict ourselves to the case of $\alpha = \beta = 2$, $d = 1$ and $\gamma = 0$, then (3.11) and (3.14) recover [11, (3.9) and (3.10)] respectively. Now choose $a > 0$ such that

$$a > (8L^2(\sigma)K^2)^{1/\rho}.$$

This implies that the following interval is nonempty:

$$\left[2, \frac{a^{\rho}}{4L^2(\sigma)K^2} \right].$$

In addition, for any p in this interval, we have that

$$\frac{\sqrt{p}L(\sigma)K}{a^{\rho/2}} \leq \frac{1}{2},$$

and we also have

$$\frac{\sqrt{p}}{a^{\rho/2}} \leq \frac{1}{2KL(\sigma)}.$$

This along with (3.14) implies that

$$N_{a,p}(u) \leq 2 \begin{cases} ([\beta] - 1)^{[\beta]-1} (ea)^{1-[\beta]} \|u_0\|_{\infty} & \beta \in (0, 1] \\ ([\beta] - 1)^{[\beta]-1} (ea)^{1-[\beta]} \|v_0\|_{\infty} + \|u_0\|_{\infty} & \beta \in (1, 2) \end{cases} + \frac{c(\sigma)}{L(\sigma)}$$

and this is precisely (3.8). Note that (3.9) follows immediately from (3.8). $\square$

We will need to recall the following result proved in [4] in order to prove Proposition 3.7 below.

P:YInc

Proposition 3.5. [2, Proposition 2.1 and 2.2] Suppose that $\alpha \in (0, 2]$, $\beta \in (0, 2)$, $\gamma > 0$, and (3.2) holds. Consider $\rho = \rho(d)$ which is defined above in (3.4). Then $Y(t, x) = Y_{\alpha, \beta, \gamma, d}(t, x)$ satisfies the following:

P:YIncCase1

1. Suppose that $0 < \theta < (\Theta - d) \wedge 2$ where Θ is given in (3.2), $0 < t, r \leq T$ for some $T > 0$, $\beta \leq 1$ and $\gamma \leq \lceil \beta \rceil - \beta$. Then there exists some $C := C(\alpha, \beta, \gamma, \nu, d, \theta, T)$ such that

$$\int_0^T ds \int_{\mathbb{R}^d} dy |Y(t-s, x-y) - Y(r-s, z-y)|^2 \leq C \left(|t-r|^\rho + |x-z|^\theta \right).$$

P:YIncCase2

2. If we only assume that $\alpha \in (0, 2]$, $\beta \in (0, 2)$, $\gamma > 0$, $0 < t, r \leq T$ for some $T > 0$ and (3.2) holds then for $0 < \theta < (\Theta - d) \wedge 2$ there exists some $C := C(\alpha, \beta, \gamma, \nu, d, \theta, T)$ such that

$$\int_0^T ds \int_{\mathbb{R}^d} dy |Y(t-s, x-y) - Y(r-s, z-y)|^2 \leq C \left(|t-r|^q + |x-z|^\theta \right),$$

where

$$0 < q < \rho.$$

Remark 3.6. In Proposition 3.5 above, we may often ignore the dependence of C on $\alpha, \beta, \gamma, \nu, d$ and just write $C(\theta, T)$.

P:NormInc

Proposition 3.7. Suppose that $u_0 = 1$ and that $v_0 = 0$ so that $u(t, x) = 1 + I(t, x)$. Let $a > 0$ and suppose that $0 < \theta < (\Theta - d) \wedge 2$ and $\rho = \rho(d) > 0$. Then for any $x, z \in \mathbb{R}$, $p \geq 2$ and for any $T > 0$ with $t \in [0, T]$,

$$\frac{\|u(t, x) - u(r, z)\|_p}{(|t-r|^q + |x-z|^\theta)^{1/2}} \leq C(p, \theta, T) [\mathcal{M}_1 + \mathcal{M}_2 e^{aT} N_{a,p}(u)], \quad (3.15)$$

E:NormInc_N

where

$$\mathcal{M}_1 = \sqrt{p}c(\sigma) \quad \text{and} \quad \mathcal{M}_2 = \sqrt{p}L(\sigma),$$

and

1. $0 < q \leq \rho$ under Case 1 of Proposition 3.5,
2. $0 < q < \rho$ under Case 2 of Proposition 3.5.

Moreover, for any

$$a > (8L^2(\sigma)K^2)^{1/\rho}.$$

and

$$p \in \left[2, \frac{a^\rho}{4L^2(\sigma)K^2} \right]$$

we have that

$$\frac{\|u(t, x) - u(r, z)\|_p}{(|t-r|^q + |x-z|^\theta)^{1/2}} \leq C(p, \theta, T) \left[\mathcal{M}_1 + \mathcal{M}_2 e^{aT} \left(2 + \frac{c(\sigma)}{L(\sigma)} \right) \right]. \quad (3.16)$$

E:NormInc_c

Proof. Note that because of the initial conditions, we see that $\|u(t, x) - u(r, z)\|_p = \|I(t, x) - I(r, z)\|_p$. We now apply Burkholder-Davies-Gundy inequality along with the triangle inequality, as was done in the proof of Proposition 3.4, to see that

$$\|I(t, x) - I(r, z)\|_p^2 \leq 4p \left\| \int_0^T ds \int_{\mathbb{R}^d} dy |Y(t-s, x-y) - Y(r-s, z-y)|^2 \sigma^2(u(s, y)) \right\|_{p/2}$$

$$\leq 4p \int_0^T ds \int_{\mathbb{R}^d} dy |Y(t-s, x-y) - Y(r-s, z-y)|^2 \|\sigma^2(u(s, y))\|_{p/2}.$$

Now by applying the Lipchitz condition on σ gives us that

$$\begin{aligned} \|I(t, x) - I(r, z)\|_p^2 &\leq 8p \int_0^T ds \int_{\mathbb{R}^d} dy |Y(t-s, x-y) - Y(r-s, z-y)|^2 \left(c^2(\sigma) + L^2(\sigma) \|u(s, y)\|_p^2 \right) \\ &\leq 8p \left[c^2(\sigma) \int_0^T ds \int_{\mathbb{R}^d} dy |Y(t-s, x-y) - Y(r-s, z-y)|^2 \right. \\ &\quad \left. + L^2(\sigma) \int_0^T ds \int_{\mathbb{R}^d} dy |Y(t-s, x-y) - Y(r-s, z-y)|^2 \exp(2as) N_{a,p}^2(u) \right] \\ &\leq 8p \left[c^2(\sigma) \int_0^T ds \int_{\mathbb{R}^d} dy |Y(t-s, x-y) - Y(r-s, z-y)|^2 \right. \\ &\quad \left. + L^2(\sigma) \exp(2aT) N_{a,p}^2(u) \int_0^T ds \int_{\mathbb{R}^d} dy |Y(t-s, x-y) - Y(r-s, z-y)|^2 \right]. \end{aligned}$$

We now apply Proposition 3.5 to see that

$$\begin{aligned} \|I(t, x) - I(r, z)\|_p^2 &\leq 8p \left[c^2(\sigma) C(\theta, T) \left(|t-r|^q + |x-z|^\theta \right) \right. \\ &\quad \left. + L^2(\sigma) \exp(2aT) N_{a,p}^2(u) C(\theta, T) \left(|t-r|^q + |x-z|^\theta \right) \right], \end{aligned}$$

where $C(\theta, T)$ is as in Proposition 3.5. Now by taking the square root of both sides and applying the identity $\sqrt{x+y} \leq \sqrt{x} + \sqrt{y}$ yields

$$\|I(t, x) - I(r, z)\|_p \leq 2\sqrt{2C(\theta, T)p} \left[c(\sigma) + L(\sigma) \exp(aT) N_{a,p}(u) \right] \left(|t-r|^q + |x-z|^\theta \right)^{1/2}.$$

This proves (3.15). It is clear that (3.16) follows directly from (3.15) and Proposition 3.4. $\square$

The holder continuity in the next proposition needs to be checked. Also I need to change the $[-R, R]$ to $|x| \leq R$ in the following proof so we can consider higher dimension than $d = 1$.

Check the following Holder continuity with [2].

P:Holder

Proposition 3.8. Suppose that u_0 and v_0 are as on Proposition 3.7 above. Then u has a version, still denoted by u , that is η_1 -Hölder continuous in time and η_2 -Hölder continuous in space with $0 < \eta_1 < q/2$ and $0 < \eta_2 < \theta/2$ where $0 < \theta < (\Theta - d) \wedge 2$ and

1. $0 < q \leq \rho$ under Case 1 of Proposition 3.5,
2. $0 < q < \rho$ under Case 2 of Proposition 3.5.

Moreover, if we consider

$$a > (8L^2(\sigma)K^2)^{1/\rho} \quad \text{and} \quad p \in \left[2, \frac{a^\rho}{4L^2(\sigma)K^2} \right],$$

where K is the constant from Proposition 3.4, then we have that

$$\mathbb{E} \left[\sup_{t \in [0, T], |x| \leq R} |u(t, x)|^p \right] \leq 2^{p-1} + C(p, \theta, T, R) \left[\mathcal{M}_1^p + \mathcal{M}_2^p e^{apT} \left(2 + \frac{c(\sigma)}{L(\sigma)} \right)^p \right].$$

Proof. Proposition 3.7 implies that

$$\|u(t, x) - u(r, z)\|_p \leq C(p, \theta, T) [\mathcal{M}_1 + \mathcal{M}_2 e^{aT} N_{a,p}(u)] \left(|t - r|^q + |x - z|^\theta \right)^{1/2}. \quad (3.17)$$

By Kolmogorov's continuity theorem, there exists a version of u , which we denote by u , that is η_1 -Hölder continuous in time and η_2 -Hölder continuous in space with $0 < \eta_1 < q/2$ and $0 < \eta_2 < \theta/2$.

Next, note that by triangle inequality,

$$\begin{aligned} |u(t, x)|^p &\leq 2^{p-1} (\|u_0\|_\infty^p + |u(t, x) - u_0(x)|^p) \\ &\leq 2^{p-1} \left(\|u_0\|_\infty^p + \sup_{t \in [0, T], x \in [-R, R]} (|u(t, x) - u_0(x)|^p) \right) \\ &= 2^{p-1} \left(1 + \sup_{t \in [0, T], x \in [-R, R]} (|u(t, x) - 1|^p) \right) \end{aligned}$$

Due to the continuity of u on the compact set $[0, T] \times [-R, R]$, there exists a point $(t_0, x_0) \in [0, T] \times [-R, R]$ such that

$$\sup_{t \in [0, T], x \in [-R, R]} (|u(t, x) - 1|^p) = |u(t_0, x_0) - 1|^p.$$

Thus, for any $(t, x) \in [0, T] \times [-R, R]$,

$$|u(t, x)|^p \leq 2^{p-1} (1 + |u(t_0, x_0) - 1|^p)$$

which in return implies that

$$\sup_{t \in [0, T], x \in [-R, R]} |u(t, x)|^p \leq 2^{p-1} (1 + |u(t_0, x_0) - 1|^p).$$

Now by taking the expectation of both sides and applying Proposition 3.7 gives us

$$\begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T], x \in [-R, R]} |u(t, x)|^p \right] &\leq 2^{p-1} + 2^{p-1} C(p, \theta, T)^p \left((2T)^{q/2} + (2R)^{\theta/2} \right)^p [\mathcal{M}_1 + \mathcal{M}_2 e^{aT} N_{a,p}(u)]^p \\ &= 2^{p-1} + C(p, \theta, T, R) [\mathcal{M}_1^p + \mathcal{M}_2^p e^{apT} N_{a,p}(u)^p]. \end{aligned}$$

Now if we consider

$$a > (8L^2(\sigma)K^2)^{1/\rho} \quad \text{and} \quad p \in \left[2, \frac{a^\rho}{4L^2(\sigma)K^2} \right],$$

then by Proposition 3.4, we have that

$$\mathbb{E} \left[\sup_{t \in [0, T], x \in [-R, R]} |u(t, x)|^p \right] \leq 2^{p-1} + C(p, \theta, T, R) \left[\mathcal{M}_1^p + \mathcal{M}_2^p e^{apT} \left(2 + \frac{c(\sigma)}{L(\sigma)} \right)^p \right].$$

□

Remark 3.9. The Hölder continuity in time proven above in Proposition 3.8 recovers the result proven by Foondun and Nane in [5, Theorem 1.9] if we restrict ourself to the case where $\beta \in (0, 1)$ and $\gamma = 1 - \beta$.

Nick's Question 3.1
Then

Suppose we consider bounded initial conditions u_0 and v_0 .

$$\begin{aligned} J_0(t, x) &= \begin{cases} [Z(t, \cdot) * u_0](x) & \beta \in (0, 1) \\ [Z(t, \cdot) * v_0](x) + \frac{\partial}{\partial t}[Z(t, \cdot) * u_0](x) & \beta \in (1, 2) \end{cases} \\ &=: \begin{cases} [Z(t, \cdot) * u_0](x) & \beta \in (0, 1) \\ [Z(t, \cdot) * v_0](x) + [Z^*(t, \cdot) * u_0](x) & \beta \in (1, 2) \end{cases}. \end{aligned}$$

We notice that

$$J_0(t, x) - J_0(s, z) = \begin{cases} [Z(t, \cdot) * u_0](x) - [Z(s, \cdot) * u_0](z) & \beta \in (0, 1) \\ [Z(t, \cdot) * v_0](x) - [Z(s, \cdot) * v_0](z) \\ \quad + [Z^*(t, \cdot) * u_0](z) - [Z^*(s, \cdot) * u_0](x) & \beta \in (1, 2) \end{cases}.$$

Lets just examine the $\beta \in (1, 2)$ case since that case deals with both Z and Z^* . We have that

$$|J_0(t, x) - J_0(s, z)| \leq \int_{\mathbb{R}} |Z(t, x - y) - Z(s, z - y)| |v_0(y)| dy + \int_{\mathbb{R}} |Z^*(t, x - y) - Z^*(s, z - y)| |u_0(y)| dy.$$

For simplicity lets look at the time and space increments separately. For time we have that

$$|J_0(t, x) - J_0(s, x)| \leq \int_{\mathbb{R}} |Z(t, x - y) - Z(s, x - y)| |v_0(y)| dy + \int_{\mathbb{R}} |Z^*(t, x - y) - Z^*(s, x - y)| |u_0(y)| dy.$$

Foondun and Nane [5, Proof of Theorem 1.13] stop right and claim that the above integrals are smooth functions in t . Then it seems that they conclude that they are η -Hölder continuous in t with $\eta = \frac{(1-\beta\gamma/\alpha)}{2}$ where γ is the Reisz Kernel exponent. In other words they say that

$$|J_0(t, x) - J_0(s, x)| \leq C|t - s|^\eta.$$

Are smooth functions also η -Hölder continuous for any $\eta > 0$? If we can argue as in [5], then we can relax our initial conditions to be bounded measurable functions u_0 and v_0 . This is also an improvement from Sanz-Sole and Millet [11] since they assume the initial conditions are continuous.

Nick's Question 3.2 Continue of Question 3.1

Suppose that $\beta \in (0, 1]$.

Time Increment: Suppose $0 \leq s < t$. Then,

$$\begin{aligned} |J(t, x) - J(s, x)| &\leq \|u_0\|_\infty \int_{\mathbb{R}^d} |Z(t, x - y) - Z(s, x - y)| dy \\ &= \|u_0\|_\infty |t - s| \int_{\mathbb{R}^d} \left| \frac{\partial}{\partial t} Z(\xi(s, t), x - y) \right| dy \end{aligned} \quad (3.18)$$

E:DZnorm

where $s \leq \xi(s, t) \leq t$. Now we will calculate $\frac{\partial}{\partial t} Z(\xi(s, t), x - y)$.

$$\begin{aligned} \frac{\partial}{\partial t} Z(\xi(s, t), x) &= \frac{\partial}{\partial t} \left(\pi^{-d/2} t^{[\beta]-1} |x|^{-d} H_{2,3}^{2,1} \left(\frac{|x|^\alpha}{2^{\alpha-1} \nu t^\beta} \middle| \begin{matrix} (1, 1) & ([\beta], \beta) \\ (d/2, \alpha/2) & (1, 1) & (1, \alpha/2) \end{matrix} \right) \right) \\ &= \frac{\partial}{\partial t} \left(\pi^{-d/2} |x|^{-d} H_{2,3}^{2,1} \left(\frac{|x|^\alpha}{2^{\alpha-1} \nu t^\beta} \middle| \begin{matrix} (1, 1) & (1, \beta) \\ (d/2, \alpha/2) & (1, 1) & (1, \alpha/2) \end{matrix} \right) \right) \end{aligned}$$

$$= \pi^{-d/2} |x|^{-d} \frac{\partial}{\partial t} \left(H_{2,3}^{2,1} \left(\frac{1}{(ct)^\beta} \begin{vmatrix} (1,1) & (1,\beta) \\ (d/2, \alpha/2) & (1,1) & (1, \alpha/2) \end{vmatrix} \right) \right)$$

where $c = (2^{\alpha-1} \nu / |x|^{-\alpha})^{1/\beta}$. We now apply [8, (2.2.6)] to see that,

$$\begin{aligned} \pi^{-d/2} |x|^{-d} \frac{\partial}{\partial t} H_{2,3}^{2,1} \left(\frac{1}{(ct)^\beta} \begin{vmatrix} (1,1) & (1,\beta) \\ (d/2, \alpha/2) & (1,1) & (1, \alpha/2) \end{vmatrix} \right) \\ = \frac{\pi^{-d/2} |x|^{-d}}{t} H_{3,4}^{3,1} \left(\frac{1}{(ct)^\beta} \begin{vmatrix} (1,1) & (1,\beta) & (0,\beta) \\ (1,\beta) & (d/2, \alpha/2) & (1,1) & (1, \alpha/2) \end{vmatrix} \right) \\ = \frac{\pi^{-d/2} |x|^{-d}}{t} H_{3,4}^{3,1} \left(\frac{1}{(ct)^\beta} \begin{vmatrix} (1,1) & (0,\beta) & (1,\beta) \\ (1,\beta) & (d/2, \alpha/2) & (1,1) & (1, \alpha/2) \end{vmatrix} \right) \\ = \frac{\pi^{-d/2} |x|^{-d}}{t} H_{2,3}^{2,1} \left(\frac{1}{(ct)^\beta} \begin{vmatrix} (1,1) & (0,\beta) \\ (d/2, \alpha/2) & (1,1) & (1, \alpha/2) \end{vmatrix} \right), \end{aligned}$$

where in the last line we used [8, (2.1.2)]. Replacing c above gives us that

$$\begin{aligned} \frac{\partial}{\partial t} Z(\xi(s, t), x) &= \frac{\pi^{-d/2} |x|^{-d}}{t} H_{2,3}^{2,1} \left(\frac{1}{(ct)^\beta} \begin{vmatrix} (1,1) & (0,\beta) \\ (d/2, \alpha/2) & (1,1) & (1, \alpha/2) \end{vmatrix} \right) \\ &= \frac{\pi^{-d/2} |x|^{-d}}{t} H_{2,3}^{2,1} \left(\frac{|x|^\alpha}{2^{\alpha-1} \nu t^\beta} \begin{vmatrix} (1,1) & (0,\beta) \\ (d/2, \alpha/2) & (1,1) & (1, \alpha/2) \end{vmatrix} \right). \end{aligned}$$

We need to calculate (3.18). In order to do this, we can calculate $\mathcal{F} \left(\frac{\partial}{\partial t} Z(\xi(s, t), \cdot) \right) (0)$. However if we examine the proof of [3, Lemma 4.2], then it appears that

$$\mathcal{F} \left(\frac{\partial}{\partial t} Z(\xi(s, t), \cdot) \right) (\xi) = \frac{1}{t} H_{1,2}^{1,1} \left(2^{-1} \nu t^\beta |\xi|^\alpha \begin{vmatrix} (0,1) \\ (0,1) & (1,\beta) \end{vmatrix} \right).$$

This can no longer be written as a Mittag-Leffler function since by [3, (A.19)] we would have

$$\frac{1}{t} H_{1,2}^{1,1} \left(2^{-1} \nu t^\beta |\xi|^\alpha \begin{vmatrix} (0,1) \\ (0,1) & (1,\beta) \end{vmatrix} \right) = \frac{1}{t} E_{\beta,0} \left(-2^{-1} \nu t^\beta |\xi|^\alpha \right),$$

and $E_{\alpha,\beta}$ is not defined for $\beta = 0$. So I am unsure of how to calculate (3.18) in this case. I need to show that

$$\mathcal{F} \left(\frac{\partial}{\partial t} Z(\xi(s, t), \cdot) \right) (0) = \frac{1}{t} H_{1,2}^{1,1} \left(0 \begin{vmatrix} (0,1) \\ (0,1) & (1,\beta) \end{vmatrix} \right) < \infty.$$

3.2 Existence of a Global Solution with Superlinear Diffusion Coefficient

We now impose the following condition on σ : As $|x|, |z| \rightarrow \infty$,

$$|\sigma(x) - \sigma(z)| \leq \sigma_2 |x - z| [\ln_+(|x - z|)]^\delta, \quad (3.19)$$

where $\sigma_2, \delta > 0$ and $\ln_+(z) = \ln(z \vee e)$ for $z > 0$. Further, this implies that as $|x| \rightarrow \infty$,

$$|\sigma(x)| \leq \sigma_1 + \sigma_2 |x| [\ln(|x|)]^\delta, \quad (3.20)$$

where $\sigma_1 = c(\sigma) = \sigma(0)$.

Remark 3.10. Because of the absence of the drift coefficient in (1.1), we may drop [11, Assumption C1].

Theorem 3.11. *Consider the initial conditions $u_0 = 1$ and $v_0 = 0$ and suppose that σ satisfies (3.19). Then for any $M > 0$, there exists a random field solution to (1.1) in $[-M, M]$ denoted as $(u(t, x) : (t, x) \in [0, T] \times [-M, M])$. This solution is unique and satisfies*

$$\sup_{t \in [0, T], x \in [-R, R]} |u(t, x)| < \infty, a.s. \quad (3.21)$$

Proof. We first solve a *truncated* version of (1.1). By this, we mean (1.1) with σ replaced by σ_N where

$$\sigma_N(x) = \sigma(x) \mathbf{1}_{\{|x| \leq N\}} + \sigma(N) \mathbf{1}_{\{x > N\}} + \sigma(-N) \mathbf{1}_{\{x \leq -N\}}.$$

Note that since σ satisfies (3.19), then σ_N is Lipschitz continuous with Lipschitz constant $L(\sigma_N) := \sigma_2 \ln(2N)^\delta$. In other words,

$$|\sigma_N(x)| \leq \sigma_1 + \sigma_2 \ln(2N)^\delta |x|.$$

Hence we can apply Theorem 3.1 and see that there exists a unique solution of the truncated version of (1.1), which we denote by $u_N := \{u_N(t, x) : (t, x) \in [0, T] \times \mathbb{R}\}$. In addition, Proposition 3.8 implies that the solution has a version, still denoted by u_N , which is η_1 -Hölder continuous in time and η_2 -Hölder continuous in space with $0 < \eta_1 < q/2$ and $0 < \eta_2 < \theta/2$.

We will now apply Proposition 3.8 to find an upper bound on the p -norm of u_N . For this, consider

$$a > (8L^2(\sigma_N)K^2)^{1/\rho} \quad \text{and} \quad p \in \left[2, \frac{a^\rho}{4L^2(\sigma_N)K^2}\right].$$

Then Proposition 3.8 implies that

$$\mathbb{E} \left(\sup_{t \in [0, T], x \in [-R, R]} |u_N(t, x)|^p \right) \leq 2^{p-1} + C(p, \theta, T, R) \left[\mathcal{M}_1^p + \mathcal{M}_2^p(N) \exp(apT) \left(2 + \frac{\sigma_1}{L(\sigma_N)} \right) \right] \quad (3.22)$$

where we recall that

$$\mathcal{M}_1 = \sqrt{p}\sigma_1 \quad \text{and} \quad \mathcal{M}_2(N) = \sqrt{pL(\sigma_N)} = \sqrt{p\sigma_2 \ln(2N)^\delta}. \quad (3.23)$$

We now use the above to prove the existence and uniqueness of (1.1) with superlinear σ satisfying (3.19). For any $T > 0$ and $N \geq 2$ with $N \in \mathbb{N}$, we define the following stopping time:

$$\tau_N := \inf \left\{ t > 0 : \sup_{|x| \leq R} |u_N(t, x)| \geq N \right\} \wedge T. \quad (3.24)$$

The uniqueness of the solution and the [local property of the stochastic integral](#) imply that on $\{t < \tau_N\}$, $u_N(t, x) = u_{N+k}(t, x)$ for any $k \in \mathbb{N}$. Hence, $(\tau_N)_{N \geq 2}$ is increasing and bounded above by T .

We now momentarily assume that $\sup_N \tau_N = T$ which would imply that $\{t < \tau_N\} \uparrow \Omega$. On each $\{t < \tau_N\}$, define $(u(t, x) : (t, x) \in [0, T] \times \mathbb{R})$ by $u(t, x) = u_N(t, x)$ and hence $u(t, x) = u_{N+k}(t, x)$ for any $k \in \mathbb{N}$. This implies that on $\{t < \tau_N\}$

$$u(t, x) = 1 + \int_0^t ds \int_{\mathbb{R}} dy Y(t-s, x-y) \sigma_N(u(s, y)) W(ds, dy).$$

However, on $\{t < \tau_N\}$, we have that $\sigma_N(u_N(t, x)) = \sigma(u(t, x))$ and so u satisfies

$$u(t, x) = 1 + \int_0^t ds \int_{\mathbb{R}} dy Y(t-s, x-y) \sigma(u(s, y)) W(ds, dy), \quad \{t < \tau_N\}.$$

Lastly, since $\{t < \tau_N\} \uparrow \Omega$, we conclude that

$$u(t, x) = 1 + \int_0^t ds \int_{\mathbb{R}} dy Y(t-s, x-y) \sigma(u(s, y)) W(ds, dy), \quad (t, x) \in [0, T] \times \mathbb{R}. \quad (3.25)$$

E:SupLinMil

It is mentioned that the stochastic integrals are no longer defined in $L^2(\Omega)$. The extension of the integral is found in Dalang and Sanz-Sole's textbook that is not yet published.

We now focus on proving that $\sup_N \tau_N = T$, or equivalently that $\mathbb{P}(\tau_N < T) \rightarrow 0$ as $N \rightarrow \infty$. To do this, note that by Chebychev's inequality

$$\begin{aligned} \mathbb{P}(\tau_N < T) &\leq \mathbb{P}\left(\sup_{t \in [0, T], x \in [-R, R]} |u_N(t, x)| \geq N\right) \\ &\leq N^{-p} \mathbb{E}\left(\sup_{t \in [0, T], x \in [-R, R]} |u_N(t, x)|^p\right). \end{aligned}$$

An application of (3.22) gives

$$\begin{aligned} N^{-p} \mathbb{E}\left(\sup_{t \in [0, T], x \in [-R, R]} |u_N(t, x)|^p\right) &\leq N^{-p} \left(2^{p-1} + C(p, \theta, T, R) \left[\mathcal{M}_1^p + \mathcal{M}_2(N)^p \exp\{apT\} \left(2 + \frac{\sigma_1}{L(\sigma_N)}\right)^p\right]\right) \\ &= \frac{2^{p-1} + C(p, \theta, T, R) \mathcal{M}_1^p}{N^p} + \frac{C(p, T, R) \mathcal{M}_2(N)^p \exp\{apT\} \left(2 + \frac{\sigma_1}{L(\sigma_N)}\right)^p}{N^p}. \end{aligned}$$

The first term clearly converges to 0 as $N \rightarrow \infty$. As for the second term, we expand using (3.23) to see that

$$\begin{aligned} \frac{\mathcal{M}_2(N)^p \exp\{apT\} \left(2 + \frac{\sigma_1}{L(\sigma_N)}\right)^p}{N^p} &= \frac{\left(\sqrt{p\sigma_2 \ln(2N)^\delta}\right)^p \exp\{apT\} \left(2 + \frac{\sigma_1}{\sqrt{p\sigma_2 \ln(2N)^\delta}}\right)^p}{N^p} \\ &\sim \frac{\ln(2N)^{p\delta/2}}{N^p} \rightarrow \infty, \end{aligned}$$

where above we use the symbol $\sim$ to mean asymptotically equivalent as $N \rightarrow \infty$. This completes the proof of the Theorem. $\square$

Remark 3.12. A potential project would be to show that the solution to (1.1) with superlinear coefficient is Holder continuous. This requires to one to extend the Kolmogorov theorem to the type of stochastic integral used to define the mild solution in (3.25). However these stochastic integrals are no longer defined in $L^2(\Omega)$. Maybe a completely new theory needs to be developed to show continuity. [9, Theorem 5.5.5] may be helpful?

4 Spatially Colored Noise Driven SPDE

In this section we will study (1.1) driven by a Gaussian noise that is white in time and colored in space. In other words, we start with a non-negative and nonnegative definite tempered distribution on $\mathcal{S}'(\mathbb{R}^d)$ which we denote as Λ . Then the Bochner-Schwartz theorem says that Λ is the Fourier transform of a non-negative, tempered and symmetric measure which we denote as μ . μ is referred to as the *spectral measure*. Because of this, the following functional is nonnegative definite and symmetric on $C_0(\mathbb{R}_+ \times \mathbb{R}^d)$:

$$C(\phi, \psi) := \int_0^\infty dt \int_{\mathbb{R}^d} \Lambda(dx) (\phi(t, \cdot) * \tilde{\psi}(t, \cdot))(x), \quad \tilde{\psi}(x) = \psi(-x). \quad (4.1)$$

E:CovNoise

Thus we may identify C as the covariance functional of some Gaussian process. This leads to the following definition.

Definition 4.1. A zero-mean Gaussian processes, $F := \{F(\phi) : \phi \in C_0(\mathbb{R}_+ \times \mathbb{R}^d)\}$ is said to be white in time and homogeneously colored in space if $\mathbb{E}(F(\phi)F(\psi)) = C(\phi, \psi)$.

NoiseAssum1

Assumption 4.2. For the remainder of this paper, we will only consider the Gaussian noises that satisfy the following three conditions:

1. the spectral measure μ is such that $\int_{\mathbb{R}^d} \frac{\mu(d\zeta)}{1+|\zeta|^\alpha} < \infty$,
2. Λ is absolutely continuous with respect to the Lebesgue measure and therefore has a density, $\Lambda(dx) = f(x)dx$. Furthermore, when this is the case, we may often write $\mu(d\zeta) = \varphi(\zeta)d\zeta$ where we have the relation $\mathcal{F}\varphi = f$,
3. the correlation functions satisfies for some $\lambda \in (0, d)$ and for all $c > 0$, $f(cx) = c^{-\lambda}f(x)$, which in return implies that $\varphi(cx) = c^{d-\lambda}\varphi(x)$.

Under this assumption, we may rewrite (4.1) as

$$C(\phi, \psi) = \int_0^\infty dt \int_{\mathbb{R}^{2d}} dx dy \phi(t, x) f(x - y) \psi(t, y)$$

and by applying Placherel's identity, this further implies that

$$\int_0^\infty dt \int_{\mathbb{R}^{2d}} dx dy \phi(t, x) f(x - y) \psi(t, y) = (2\pi)^{-d} \int_0^\infty dt \int_{\mathbb{R}^{2d}} \mu(d\zeta) \mathcal{F}(\phi(t, \cdot))(\zeta) \overline{\mathcal{F}(\psi(t, \cdot))(\zeta)}.$$

We now prove a lemma which will be used to prove the existence and uniqueness of the solution. It is proven by Foondun and Nane [5] in the case that

$$\beta \in (0, 1) \quad \text{and} \quad \gamma = 1 - \beta. \quad (4.2)$$

A:Y=GAssump

L:YfYbdd

Lemma 4.3. Suppose Assumption 4.2 holds. Then there exists a constant $C(\beta, \gamma) =: C > 0$ such that for any $x, y \in \mathbb{R}^d$ we have that

$$\int_{\mathbb{R}^{2d}} Y(t, x - z) Y(t, y - w) f(w - z) dz dw \leq C \left(\frac{\nu}{2}\right)^{-\lambda/\alpha} t^{2(\beta+\gamma-1)-\beta\lambda/\alpha} \int_{\mathbb{R}^d} \frac{\mu(d\zeta)}{(1 + |\xi|^\alpha)^2} < \infty \quad (4.3)$$

E:YYfBdd

Proof. Through a change of variables, we may write the left side of (4.3) as the following:

$$\int_{\mathbb{R}^{2d}} Y(t, x - z) Y(t, y - w) f(w - z) dz dw = \int_{\mathbb{R}^d} Y(t, u - (x - y)) (Y(t, \cdot) * f)(u) du,$$

where $*$ denotes the spatial convolution. Now an application of Parseval's identity gives us that

$$\int_{\mathbb{R}^d} Y(t, u - (x - y)) (Y(t, \cdot) * f)(u) du = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \mathcal{F}Y(t, \cdot - (x - y))(\zeta) \mathcal{F}(Y(t, \cdot) * f)(\zeta) d\zeta.$$

Applying the translation property of the Fourier transform and [4, Theorem 4.1] gives us that

$$\begin{aligned} \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \mathcal{F}Y(t, \cdot - (x - y))(\zeta) \mathcal{F}(Y(t, \cdot) * f)(\zeta) d\zeta \\ = \frac{t^{2(\beta+\gamma-1)}}{(2\pi)^d} \int_{\mathbb{R}^d} \exp(-i(x - y)\zeta) E_{\beta, \beta+\gamma} \left(-\frac{\nu t^\beta |\zeta|^\alpha}{2} \right)^2 \mu(d\zeta). \end{aligned}$$

Hence by taking the absolute value on both sides and recalling the positivity of f , Y and the Mittag-Leffler function we get that

$$\int_{\mathbb{R}^{2d}} Y(t, x - z) Y(t, y - w) f(w - z) dz dw \leq \frac{t^{2(\beta+\gamma-1)}}{(2\pi)^d} \int_{\mathbb{R}^d} E_{\beta, \beta+\gamma} \left(-\frac{\nu t^\beta |\zeta|^\alpha}{2} \right)^2 \mu(d\zeta).$$

Now, by applying [10, Theorem 1.6], we see that there exists a $C > 0$ depending only on β and γ such that

$$\frac{t^{2(\beta+\gamma-1)}}{(2\pi)^d} \int_{\mathbb{R}^d} E_{\beta,\beta+\gamma} \left(-\frac{\nu t^\beta |\xi|^\alpha}{2} \right)^2 \mu(d\xi) \leq \frac{C t^{2(\beta+\gamma-1)}}{(2\pi)^d} \int_{\mathbb{R}^d} \frac{\mu(d\xi)}{(1 + \frac{\nu}{2} t^\beta |\xi|^\alpha)^2}.$$

Another change of variables implies that

$$\frac{C t^{2(\beta+\gamma-1)}}{(2\pi)^d} \int_{\mathbb{R}^d} \frac{\mu(d\xi)}{(1 + \frac{\nu}{2} t^\beta |\xi|^\alpha)^2} \leq \frac{C}{(2\pi)^d} \left(\frac{\nu}{2} \right)^{-\lambda/\alpha} t^{2(\beta+\gamma-1)-\beta\lambda/\alpha} \int_{\mathbb{R}^d} \frac{\mu(d\xi)}{(1 + |\xi|^\alpha)^2}.$$

The finiteness of the above integral follows from Assumption 4.2. $\square$

Nick's Question 4.1 Lemma 4.3 is a problem. Later on we will need to assume that

$$2(\beta + \gamma - 1) - \beta\lambda/\alpha > 0 \quad (4.4)$$

A:Problem

which is stronger than the original assumption that

$$2(\beta + \gamma) - 1 - \beta\lambda/\alpha.$$

(4.4) excludes the SHE and the case when $\gamma = \lceil \beta \rceil - 1$.

RieszKernel

Example 4.4. Consider the function $f(x)$, which is the well-known Riesz kernel with parameter $\lambda \in (0, d)$:

$$f(x) := \frac{1}{|x|^\lambda} \quad \text{and} \quad \varphi(x) := \mathcal{F}f(\xi) = \frac{K_f}{|\xi|^{d-\lambda}},$$

where

$$K_f = \pi^{-d/2} 2^{-\lambda} \frac{\Gamma((d-\lambda)/2)}{\Gamma(\lambda/2)}.$$

Considering Assumption 4.2, if we assume $\lambda < \min\{d, 2\}$, then we may consider our Gaussian processes to have its correlation function given to be the Riesz kernel, $f(x) = |x|^{-\lambda}$.

4.1 Lipschitz Diffusion Coefficient

We first examine (1.1) with Lipschitz diffusion coefficient. As we will see below in Theorem 4.5, to establish the existence and uniqueness of a random field solution to (1.1), we will need to assume Dalang's condition:

$$\int_0^t ds \int_{\mathbb{R}^d} dy_1 dy_2 Y(s, y_1) f(y_1 - y_2) Y(s, y_2) < \infty, \quad \text{for all } t > 0, \quad (4.5)$$

C:DalangC1

which is equivalent to

$$\rho(\lambda) > 0 \quad \text{and} \quad \lambda < 2\alpha, \quad (4.6)$$

E:DalangEqu

where we recall from (3.4) that

$$\rho(x) := 2\beta + 2\gamma - 1 - \beta x/\alpha. \quad (4.7)$$

We assume that there exists some $L_\sigma > 0$ such that

$$|\sigma(x) - \sigma(y)| \leq L_\sigma |x - y|.$$

We can now prove the existence and uniqueness of the solution to (1.1) driven by the noise $\dot{F}$.

Theorem 4.5. Suppose that $\lambda\beta/\alpha - 2(\beta + \gamma - 1) < 1$ and in addition, suppose that $\lambda < 2\alpha$. Then there exists a unique $L^2(\Omega)$ -solution to (1.1) driven by $\dot{F}$ such that

$$u(t, x) = J_0(t, x) + I(t, x), \quad a.s.$$

Proof. The proof follows the strategy laid out in Walsh [12]. We start by defining

$$u_0(t, x) := J_0(t, x) \quad \text{and} \quad u_n(t, x) := J_0(t, x) + \int_0^t \int_{\mathbb{R}^d} Y(t-s, x-y) \sigma(u_n(s, y)) F(ds, dy).$$

In addition, we define

$$D_n(t, x) := \mathbb{E} |u_{n+1}(t, x) - u_n(t, x)|^2 \quad \text{and} \quad H_n(t) := \sup_{x \in \mathbb{R}^d} D_n(t, x).$$

Lastly, we define

$$\Sigma(t, x, n) := |\sigma(u_n(t, x)) - \sigma(u_{n-1}(t, x))|.$$

With this notation we see that

$$D_n(t, x) = \int_0^t ds \int_{\mathbb{R}^d} dz dy Y(t-s, x-y) Y(t-s, x-z) f(y-z) \mathbb{E} (\Sigma(s, y, n) \Sigma(s, z, n)).$$

Now note that by the Lipschitz property of σ , we have that

$$\begin{aligned} \mathbb{E} (\Sigma(s, y, n) \Sigma(s, z, n)) &\leq L_\sigma^2 \mathbb{E} (|u_n(s, y) - u_{n-1}(s, y)| |u_n(s, z) - u_{n-1}(s, z)|) \\ &\leq L_\sigma^2 \|u_n(s, y) - u_{n-1}(s, y)\| \|u_n(s, z) - u_{n-1}(s, z)\| \\ &\leq L_\sigma^2 \sqrt{D_{n-1}(s, y) D_{n-1}(s, z)} \\ &\leq L_\sigma^2 H_{n-1}(s). \end{aligned}$$

Thus,

$$\begin{aligned} D_n(t, x) &\leq L_\sigma^2 \int_0^t H_{n-1}(s) \int_{\mathbb{R}^d} dz dy Y(t-s, x-y) Y(t-s, x-z) f(y-z) \\ &\leq CL_\sigma^2 \left(\frac{\nu}{2}\right)^{-\lambda/\alpha} \int_0^t ds \frac{H_{n-1}(s)}{(t-s)^{\lambda\beta/\alpha - 2(\beta+\gamma-1)}}, \end{aligned}$$

where the last inequality follows from Lemma 4.3. In return, this implies that

$$H_n(t) \leq CL_\sigma^2 \left(\frac{\nu}{2}\right)^{-\lambda/\alpha} \int_0^t ds \frac{H_{n-1}(s)}{(t-s)^{\lambda\beta/\alpha - 2(\beta+\gamma-1)}}.$$

Since we assume that $\lambda\beta/\alpha - 2(\beta + \gamma - 1) < 1$, then the result follows from the discussion before the proof of [12, Lemma 3.3]. $\square$

Remark 4.6. One should note that when considering the SHE ($\beta = 1$, $\alpha = 2$ and $\gamma = 0$), the assumptions $\lambda\beta/\alpha - 2(\beta + \gamma - 1) < 1$ and $\lambda < 2\alpha$ reduce down to $\lambda < 2$ and $\lambda < 3$ which recovers the well-known condition that one must have $\lambda < \min\{2, d\}$.

Nick's Question 4.2 In the existence and uniqueness proof in [5, Theorem 1.10], the following is shown

$$H_n(t) \leq \text{const.} \int_0^t \frac{H_{n-1}(s)}{(t-s)^{\gamma\beta/\alpha}}.$$

Note that in this formula, γ is denoting the Riesz kernel parameter. I believe that we need to assume that

$$\gamma\beta/\alpha < 1$$

however in this proof they say we must assume that $d\beta/\alpha < 1$. I think this is a typo. However, this assumption still is not stated in the statement of the theorem. Another

issue is that they need to assume that

$$\gamma < \alpha$$

in order to apply [5, Lemma 2.7] and this assumption is not stated in the statement of the theorem. This assumption however would imply that $\gamma\beta/\alpha < 1$ since in [5] $\beta \in (0, 1)$.

In addition, I believe that there is a typo in the statement of [12, Lemma 3.3]. I believe that in Lemma 3.3 *ibid.* we should assume that $a > -1$ and not $a > 1$. This would be consistent with [12, (3.14)].

Nick's Question 4.3 The following tries to give an upper bound to $\sup_{x \in \mathbb{R}^d} \mathbb{E}|u(t, x)|^2$ by using techniques from [5]. There is an issue with the upper bound of $J_0(t, x)$ depending on t when the initial condition $v_0 \neq 0$.

Theorem 4.7 ([5, Theorem 1.10]). *Under the assumptions of Theorem 4.5 we have that*

$$\sup_{x \in \mathbb{R}^d} \mathbb{E}|u(t, x)|^2 \leq \dots \quad (4.8)$$

Proof. The proof follows the strategy from the proof of [5, Theorem 1.10]. We start by defining $A(t) = \sup_{x \in \mathbb{R}^d} \mathbb{E}|u(t, x)|^2$. We will prove the following inequality:

$$A(t) \leq C_1 + C_2 t^{2(\lceil \beta \rceil - 1)} + C_3 \int_0^t \frac{A(s)}{(t-s)^{\beta\lambda/\alpha - 2(\beta+\gamma-1)}} ds, \quad (4.9) \quad \boxed{\text{E:Renewal}}$$

To do this, we note that

$$\begin{aligned} \mathbb{E}|u(t, x)|^2 &\leq J_0^2(t, x) + \int_0^t ds \int_{\mathbb{R}^{2d}} dy dz Y(t-s, x-y) Y(t-s, x-z) f(y-z) \mathbb{E}(|\sigma(u(s, y))| |\sigma(u(s, z))|) \\ &:= J_0^2(t, x) + I_2(t, x) \end{aligned} \quad (4.10) \quad \boxed{\text{E:u2normbdd}}$$

Note that for bounded u_0 and v_0 we have that

$$\begin{aligned} |J_0(t, x)| &\leq \begin{cases} |(Z(t, \cdot) * u_0)(x)| & \beta \in (0, 1] \\ |(Z(t, \cdot) * v_0)(x)| + |(Z^*(t, \cdot) * u_0)(x)| & \beta \in (1, 2) \end{cases} \\ &\leq \begin{cases} \|u_0\|_\infty |(Z(t, \cdot) * 1)(x)| & \beta \in (0, 1] \\ \|v_0\|_\infty |(Z(t, \cdot) * 1)(x)| + \|u_0\|_\infty |(Z^*(t, \cdot) * 1)(x)| & \beta \in (1, 2) \end{cases} \\ &= \begin{cases} \|u_0\|_\infty & \beta \in (0, 1] \\ \|v_0\|_\infty t + \|u_0\|_\infty & \beta \in (1, 2). \end{cases} \end{aligned}$$

Next we note that

$$\begin{aligned} \mathbb{E}(|\sigma(u(s, y))| |\sigma(u(s, z))|) &\leq L_\sigma^2 \mathbb{E}(|u(s, y)| |u(s, z)|) \\ &\leq L_\sigma^2 \mathbb{E}(|u(s, z)|)^{1/2} \mathbb{E}(|u(s, y)|)^{1/2} \\ &\leq A(s). \end{aligned}$$

Therefore, we can bound I_2 in (4.10) by the following:

$$I_2(t, x) \leq L_\sigma^2 \int_0^t ds A(s) \int_{\mathbb{R}^{2d}} dy dz Y(t-s, x-y) Y(t-s, x-z) f(y-z).$$

Now applying Lemma ?? we see that

$$I_2(t, x) \leq C_2 \int_0^t \frac{A(s)}{t^{\beta\lambda/\alpha-2(\beta+\gamma-1)}}.$$

Combining the above gives us (4.9). $\square$

Remark 4.8. The bound in (4.9) is different from [5, Proposition 2.12] for the case of $\beta \in (1, 2)$. The $C_2 t$ term changes things. A similar exponential bound may be found in this case by starting from (4.10) and then using the Piccard iteration techniques from [1, Lemma 2.2].

Proposition 4.9 ([5, Proposition 2.12]). *Let $k > 0$ and suppose that f is locally integrable. Suppose there exists a $K > 0$ satisfies*

$$f(t) \leq K(1 + t^2) + K \int_0^t (t - s)^{k-1} f(s) ds. \quad (4.11) \quad \boxed{\text{E:fRenew}}$$

Then

$$f(t) \leq K_1 \exp(K_2 \cdots t). \quad (4.12)$$

Proof. We follow the proof from [6, Proposition 2.5]. For notation, let $g(t) = K(1 + t^2)$ and define

$$(\mathcal{A}\theta)(t) := K \int_0^t (t - s)^{k-1} \theta(s) ds$$

for any locally integrable function θ . Then (4.11) can be written as

$$f(t) \leq g(t) + \mathcal{A}\theta(t).$$

By iterating this inequality, we see that

$$f(t) \leq \sum_{i=1}^{n-1} (\mathcal{A}^i g)(t) + (\mathcal{A}^n f)(t).$$

It is stated in the proof of [6, Proposition 2.5] that

$$(\mathcal{A}^n f)(t) = \frac{(K\Gamma(\rho))^2}{\Gamma(n\rho)} \int_0^t (t - s)^{n\rho-1} f(s) ds.$$

Using this, we can calculate $(\mathcal{A}^i g)(t)$ as follows:

$$\begin{aligned} (\mathcal{A}^i g)(t) &= \frac{(K\Gamma(\rho))^i}{\Gamma(i\rho)} \int_0^t (t - s)^{i\rho-1} g(s) ds \\ &= K \frac{(K\Gamma(\rho))^i}{\Gamma(i\rho)} \int_0^t (t - s)^{i\rho-1} (1 + s^2) ds \\ &= K \frac{(K\Gamma(\rho))^i}{\Gamma(i\rho)} \cdot \frac{t^{\rho i} (1 + \rho i + t)}{\rho i (1 + \rho i)}. \end{aligned}$$

In addition, it is stated in the proof of [6, Proposition 2.5] that $(\mathcal{A}^n f)(t) \rightarrow 0$ as $n \rightarrow \infty$. Thus, we can say

$$\begin{aligned} f(t) &\leq \sum_{i=0}^{\infty} K \frac{(K\Gamma(\rho))^i}{\Gamma(i\rho)} \cdot \frac{t^{\rho i}(1+\rho i+t)}{\rho i(1+\rho i)} \\ &= \sum_{i=0}^{\infty} K \frac{(Kt^{\rho}\Gamma(\rho))^i}{\Gamma(i\rho)} \cdot \frac{1+\rho i+t}{\rho i(1+\rho i)}. \end{aligned}$$

In the proof of [6, Proposition 2.5], it is stated that

$$\sum_{i=0}^{\infty} \frac{(Kt^{\rho}\Gamma(\rho))^i}{\Gamma(i\rho)} = E_{\rho}(K\Gamma(\rho)t^{\rho}),$$

where E_{ρ} is the one-parameter Euler-Mittag function. The $\frac{1+\rho i+t}{\rho i(1+\rho i)}$ term now makes this untrue. $\square$

The following $L^2(\mathbb{R}^d)$ norms may be helpful. I may remove this. Not sure if I actually need it.

2NormZandZ*

Lemma 4.10. *Suppose that $d < 2\alpha$ and that $\beta \in (0, 2)$ and that $\gamma \leq 0$. Then,*

$$\int_{\mathbb{R}^d} Z^2(t, x) dx = C_Z t^{[\beta]-1-d\beta/\alpha} \quad (4.13) \quad \text{E:2NormZ}$$

where $C_Z = (2\pi)^{-d}(\nu/2)^{-d/\alpha} \int_{\mathbb{R}^d} E_{\beta, [\beta]} (-2^{-1}\nu t^{\beta} |\xi|^{\alpha})^2 d\xi$. In addition, we have that

$$\int_{\mathbb{R}^d} Z^{*2}(t, x) dx = C_{Z*} t^{-d\beta/\alpha} \quad (4.14) \quad \text{E:2NormZ*}$$

where $C_{Z*} = (2\pi)^{-d}(\nu/2)^{-d/\alpha} \int_{\mathbb{R}^d} E_{\beta, 1} (-2^{-1}\nu t^{\beta} |\xi|^{\alpha})^2 d\xi$.

Proof. We apply Parseval's identity and [4, Theorem 4.1] to see that

$$\begin{aligned} \int_{\mathbb{R}^d} Z^2(t, x) dx &= \frac{t^{[\beta]-1}}{(2\pi)^d} \int_{\mathbb{R}^d} E_{\beta, [\beta]} (-2^{-1}\nu t^{\beta} |\xi|^{\alpha})^2 d\xi \\ &= \frac{t^{[\beta]-1-d\beta/\alpha}}{(2\pi)^d} \left(\frac{\nu}{2}\right)^{-d/\alpha} \int_{\mathbb{R}^d} E_{\beta, [\beta]} (-|\xi|^{\alpha})^2 d\xi, \end{aligned}$$

where in the last line we applied a change of variables. Note that by [10, Theorem 1.6], the integral above on the right hand side is finite when $2\alpha > d$. This proves (4.13). Similarly we have

$$\begin{aligned} \int_{\mathbb{R}^d} Z^{*2}(t, x) dx &= \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} E_{\beta, 1} (-2^{-1}\nu t^{\beta} |\xi|^{\alpha})^2 d\xi \\ &= \frac{t^{-d\beta/\alpha}}{(2\pi)^d} \left(\frac{\nu}{2}\right)^{-d/\alpha} \int_{\mathbb{R}^d} E_{\beta, 1} (-2^{-1}\nu t^{\beta} |\xi|^{\alpha})^2 d\xi, \end{aligned}$$

where again the integral above on the right hand side is finite when $d < 2\alpha$. $\square$

4.2 Section 4.2 Millet and Sanz-Solé [11]

Proposition 4.11. [11, Proposition 4.4] *Let u_0 and v_0 be Borel functions satisfying $\|u_0\|_{\infty} + \|v_0\|_{\infty} < \infty$. Suppose that $d < 2\alpha$ and for this proof consider $\rho = \rho(\lambda)$ which is defined in (3.4). Then there exists a universal constant $K := K(\alpha, \beta, \gamma, d, \lambda) > 0$ such that for any*

$$a^{\rho} > (8L^2(\sigma)K^2).$$

and

$$p \in \left[2, \frac{a^\rho}{4L^2(\sigma)K^2} \right].$$

we have that

$$N_{a,p}(u) \leq 2\mathcal{T}_0 + \frac{c(\sigma)}{L(\sigma)} \quad (4.15) \quad \boxed{\text{E:ColorNbdd}}$$

where

$$\mathcal{T}_0 = \begin{cases} (\lceil \beta \rceil - 1)^{\lceil \beta \rceil - 1} (ea)^{1 - \lceil \beta \rceil} \|u_0\|_\infty & \beta \in (0, 1] \\ (\lceil \beta \rceil - 1)^{\lceil \beta \rceil - 1} (ea)^{1 - \lceil \beta \rceil} \|v_0\|_\infty + \|u_0\|_\infty & \beta \in (1, 2). \end{cases}$$

Moreover for any $T > 0$,

$$\sup_{x \in \mathbb{R}^d} \mathbb{E}(|u(t, x)|^p) \leq \exp(apt) \left[2\mathcal{T}_0 + \frac{c(\sigma)}{L(\sigma)} \right]^p, \quad t \in [0, T]. \quad (4.16) \quad \boxed{\text{E:ColorPMom}}$$

Proof. We note that $\mathcal{N}_{a,p}(J)$ has the same upper bound as in Proposition 3.4 above. In other words,

$$\mathcal{N}_{a,p}(J) \leq \begin{cases} \|u_0\|_\infty & \beta \in (0, 1] \\ (ea)^{-1} \|v_0\|_\infty + \|u_0\|_\infty & \beta \in (1, 2). \end{cases}$$

We know work to find an upper bound for $\mathcal{N}_{a,p}(I)$. The Burkholder-Davies-Gundy inequality and Hölder's inequality imply that

$$\|I(t, x)\|_p^2 \leq 4p \int_0^t ds \iint_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, x-y_1) f(x-y) Y(t-s, x-y_2) \|\sigma(u(s, y_1))\|_p \|\sigma(u(s, y_2))\|_p.$$

We now apply the Lipschitz property to see that for $i \in \{1, 2\}$,

$$\|\sigma(u(s, y_i))\|_p \leq c(\sigma) + L(\sigma) \|u(s, y_i)\|_p.$$

Therefore, we may say that

$$\begin{aligned} \|I(t, x)\|_p^2 &\leq 4p \int_0^t ds \iint_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, x-y_1) f(x-y) Y(t-s, x-y_2) \\ &\quad \times \prod_{i=1}^2 c(\sigma) + L(\sigma) \|u(s, y_i)\|_p \\ &\leq 4p \int_0^t ds \iint_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, x-y_1) f(x-y) Y(t-s, x-y_2) [c(\sigma) + L(\sigma) \exp(as) N_{a,p}(u)]^2 \\ &\leq 8pc(\sigma)^2 \int_0^t ds \iint_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, x-y_1) f(x-y) Y(t-s, x-y_2) \\ &\quad + 8pN_{a,p}(u)^2 L(\sigma)^2 \int_0^t ds \exp(2as) \iint_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, x-y_1) f(x-y) Y(t-s, x-y_2). \end{aligned}$$

Now by applying Lemma ??, we may bound the $\iint_{\mathbb{R}^{2d}} dy_1 dy_2$ integrals by the quantity

$$K_1(t-s)^{2(\beta+\gamma-1)-\beta\lambda/\alpha},$$

where

$$K_1 := CK_f \left(\frac{\nu}{2} \right)^{-\lambda/\alpha},$$

and C is as in Lemma ?. In return, this gives us that

$$\|I(t, x)\|_p^2 \leq 8pK_1 c(\sigma)^2 \int_0^t (t-s)^{2(\beta+\gamma-1)-\beta\lambda/\alpha} ds$$

$$\begin{aligned}
& + 8pK_1N_{a,p}(u)^2L(\sigma)^2 \int_0^t \exp(2as)(t-s)^{2(\beta+\gamma-1)-\beta\lambda/\alpha} ds \\
& = 8pK_1c(\sigma)^2 \frac{t^\rho}{\rho} + 8pK_1N_{a,p}(u)^2L(\sigma)^2 \frac{\exp(2at) [\Gamma(\rho) - \Gamma(\rho, 2at)]}{(2a)^\rho} \\
& =: \Delta_1(t) + \Delta_2(t),
\end{aligned}$$

where $\Gamma(\cdot)$ denotes the Gamma function and $\Gamma(\cdot, \circ)$ is the incomplete-upper Gamma function. Note that

$$\mathcal{N}_{a,p}(I)^2 \leq \sup_{t \geq 0} \exp(-2at) (\Delta_1(t) + \Delta_2(t)).$$

We start with bounding $\sup_{t \geq 0} \exp(-2at) \Delta_1(t)$. Recall the identities in (3.7). With this we see that

$$\sup_{t \geq 0} \exp(-2at) t^\rho = (\rho)^\rho (2ea)^{-\rho}.$$

Hence,

$$\sup_{t \geq 0} \exp(-2at) \Delta_1(t) \leq \frac{8pK_1c(\sigma)^2}{\rho} \left(\frac{\rho}{2ea} \right)^\rho.$$

Next we bound $\sup_{t \geq 0} \exp(-2at) \Delta_2(t)$. Note that for any $t \geq 0$,

$$\begin{aligned}
\exp(-2at) \Delta_2(t) & = 8pK_1N_{a,p}(u)^2L(\sigma)^2 \frac{[\Gamma(\rho) - \Gamma(\rho, 2at)]}{(2a)^\rho} \\
& \leq 8pK_1N_{a,p}(u)^2L(\sigma)^2 \frac{\Gamma(\rho)}{(2a)^\rho}.
\end{aligned}$$

Putting this all together gives us that

$$\mathcal{N}_{a,p}(I)^2 \leq \frac{8pK_1c(\sigma)^2}{\rho} \left(\frac{\rho}{2ea} \right)^\rho + \frac{8pK_1\Gamma(\rho)L(\sigma)^2}{(2a)^\rho} N_{a,p}(u)^2,$$

and thus,

$$\mathcal{N}_{a,p}(I) \leq 2c(\sigma) \sqrt{\frac{2pK_1}{\rho}} \left(\frac{\rho}{2ea} \right)^{\rho/2} + 2L(\sigma) \frac{\sqrt{2pK_1\Gamma(\rho)}}{(2a)^{\rho/2}} N_{a,p}(u).$$

Similar to the proof of Proposition 3.4 above, we define

$$K = K(\alpha, \beta, \gamma, d, \lambda) := \max \left\{ 2\sqrt{\frac{2K_1}{\rho}} \left(\frac{\rho}{2e} \right)^{\rho/2}, 2\frac{\sqrt{2K_1\Gamma(\rho)}}{(2)^{\rho/2}} \right\}.$$

We now see that

$$\mathcal{N}_{a,p}(I) \leq Kc(\sigma) \frac{\sqrt{p}}{a^{\rho/2}} + KL(\sigma) \frac{\sqrt{p}}{a^{\rho/2}} \mathcal{N}_{a,p}(u) \quad (4.17)$$

This and the bound for $\mathcal{N}_{a,p}(J)$ found above implies that

$$\begin{aligned}
\mathcal{N}_{a,p}(u) & \leq \mathcal{N}_{a,p}(J) + \mathcal{N}_{a,p}(I) \\
& \leq \begin{cases} \|u_0\|_\infty & \beta \in (0, 1] \\ (ea)^{-1} \|v_0\|_\infty + \|u_0\|_\infty & \beta \in (1, 2) \end{cases} \\
& \quad + Kc(\sigma) \frac{\sqrt{p}}{a^{\rho/2}} + KL(\sigma) \frac{\sqrt{p}}{a^{\rho/2}} \mathcal{N}_{a,p}(u).
\end{aligned} \quad (4.18)$$

Now consider

$$a^\rho > 8K^2L(\sigma)^2.$$

This implies that the interval

$$\left[2, \frac{a^\rho}{4K^2L(\sigma)^2}\right]$$

is nonempty. Hence, for any p in this interval we have that

$$\frac{KL(\sigma)\sqrt{p}}{a^{\rho/2}} \leq \frac{1}{2}.$$

This along with (4.18) implies that

$$\mathcal{N}_{a,p}(u) \leq 2 \begin{cases} \|u_0\|_\infty & \beta \in (0, 1] \\ (ea)^{-1} \|v_0\|_\infty + \|u_0\|_\infty & \beta \in (1, 2) \end{cases} + \frac{c(\sigma)}{L(\sigma)}.$$

Its easy to see that (4.16) follows directly from (4.15). $\square$

We now state a lemma which will be used later on.

Parseval

Lemma 4.12. *Let ϕ and ψ be bounded Borel functions. Then for $s, t > 0$ and $z \in \mathbb{R}^d$ we have,*

$$\iint_{\mathbb{R}^{2d}} dx dy \phi(x)Y(t, x)\psi(y)Y(s, y)f(x - y + z) = (2\pi)^{-d} \int_{\mathbb{R}^d} \mathcal{F}(\phi Y(t, \cdot))(\xi) \overline{\mathcal{F}(\psi Y(s, \cdot))(\xi)} e^{-iz \cdot \xi} \mu(d\xi) \quad (4.19)$$

Proof. For notation, define $Y_\phi(t, x) = \phi(x)Y(t, x)$ and $Y_\psi(t, x) = \psi(x)Y(t, x)$. The boundedness and measureability of ϕ and ψ guarantees the existence of $\mathcal{F}Y_\phi$ and $\mathcal{F}Y_\psi$. Thus we have,

$$\iint_{\mathbb{R}^{2d}} dx dy Y_\phi(t, x)Y_\psi(s, y)f(x - y + z) = \int_{\mathbb{R}^d} dx Y_\phi(t, x)(Y_\psi(s, \cdot) * f(\cdot + z))(x).$$

Parseval's identity implies that

$$\begin{aligned} \int_{\mathbb{R}^d} dx Y_\phi(t, x)(Y_\psi(s, \cdot) * f(\cdot + z))(x) &= (2\pi)^{-d} \int_{\mathbb{R}^d} d\zeta \mathcal{F}(Y_\phi(t, \cdot))(\zeta) \overline{\mathcal{F}((Y_\psi(s, \cdot))(\zeta) \mathcal{F}(f(\cdot + z)))} \\ &= (2\pi)^{-d} \int_{\mathbb{R}^d} \mu(d\zeta) \mathcal{F}(Y_\phi(t, \cdot))(\zeta) \overline{\mathcal{F}((Y_\psi(s, \cdot))(\zeta) e^{-iz \cdot \zeta})}, \end{aligned}$$

where $\mu(d\xi) = \varphi(\xi)d\xi$. This completes the proof. $\square$

Proposition 4.13 ([11, Proposition 4.8]). *Statement...*

Proof. By the Burkholder Davies Gundy inequality, we have that

$$\begin{aligned} \|I(t, x) - I(t, z)\|_p^2 &\leq 4p \left\| \int_0^t ds \int_{\mathbb{R}^{2d}} dy_1 dy_2 f(y_1 - y_2) \prod_{i=1}^2 (Y(t - s, x - y_i) - Y(t - s, z - y_i) \sigma(u(s, y_i))) \right\|_{p/2} \\ &=: 4p \|Q\|_{p/2} \end{aligned}$$

We now reorganize the integrand of Q above by following [7, Theorem 3.1]. By expanding the integrand of Q , we get that

$$\begin{aligned} Q &= \int_0^t ds \int_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t - s, x - y_1) Y(t - s, x - y_2) f(y_1 - y_2) \sigma(u(s, y_1)) \sigma(u(s, y_2)) \\ &\quad - \int_0^t ds \int_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t - s, x - y_1) Y(t - s, z - y_2) f(y_1 - y_2) \sigma(u(s, y_1)) \sigma(u(s, y_2)) \end{aligned}$$

$$\begin{aligned}
& - \int_0^t ds \int_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, z-y_1) Y(t-s, x-y_2) f(y_1-y_2) \sigma(u(s, y_1)) \sigma(u(s, y_2)) \\
& + \int_0^t ds \int_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, z-y_1) Y(t-s, z-y_2) f(y_1-y_2) \sigma(u(s, y_1)) \sigma(u(s, y_2)) \\
& =: I_1 + I_2 + I_3 + I_4.
\end{aligned}$$

Now let $\xi = x - z$ and consider the following notation:

$$\begin{aligned}
\Sigma_x(s, y) &= \sigma(u(s, x-y)), \quad \Sigma_{x,z}(s, y) = \sigma(u(s, x-y)) - \sigma(u(s, z-y)), \\
h_1(s, y_1, y_2) &= f(y_1, y_2) \Sigma_{x,z}(s, y_1) \Sigma_{x,z}(s, y_2) \\
h_2(s, y_1, y_2) &= [f(y_1 - y_2 + \xi) - f(y_1 - y_2)] \Sigma_x(s, y_2) \Sigma_{x,z}(s, y_1) \\
h_3(s, y_1, y_2) &= h_2(s, y_2, y_1) \\
h_4(s, y_1, y_2) &= [2f(y_1 - y_2) - f(y_1 - y_2 + \xi) - f(y_1 - y_2 - \xi)] \Sigma_x(s, y) \Sigma_x(s, z).
\end{aligned}$$

With this notation along with a change of variables, we rewrite I_1 through I_4 as the following:

$$\begin{aligned}
I_1 &= \int_0^t ds \int_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, y_1) Y(t-s, y_2) f(y_1-y_2) \Sigma_x(s, y_1) \Sigma_x(s, y_2) \\
I_2 &= - \int_0^t ds \int_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, y_1) Y(t-s, y_2) f(y_1-y_2-\xi) \Sigma_x(s, y_1) \Sigma_z(s, y_2) \\
&= \int_0^t ds \int_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, y_1) Y(t-s, y_2) f(y_1-y_2-\xi) \Sigma_x(s, y_1) \Sigma_{x,z}(s, y_2) \\
&\quad - \int_0^t ds \int_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, y_1) Y(t-s, y_2) f(y_1-y_2-\xi) \Sigma_x(s, y_1) \Sigma_x(s, y_2) \\
I_3 &= \int_0^t ds \int_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, y_1) Y(t-s, y_2) f(y_1-y_2+\xi) \Sigma_x(s, y_2) \Sigma_{x,z}(s, y_1) \\
&\quad - \int_0^t ds \int_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, y_1) Y(t-s, y_2) f(y_1-y_2+\xi) \Sigma_x(s, y_1) \Sigma_x(s, y_2) \\
I_4 &= \int_0^t ds \int_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, y_1) Y(t-s, y_2) f(y_1-y_2) \Sigma_z(s, y_1) \Sigma_z(s, y_2) \\
&= \int_0^t ds \int_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, y_1) Y(t-s, y_2) f(y_1-y_2) (\Sigma_{x,z}(s, y_1) - \Sigma_x(s, y_1) (\Sigma_{x,z}(s, y_2) - \Sigma_x(s, y_2))),
\end{aligned}$$

where above I_3 is derived in the same way I_2 was derived. Now if we expand I_4 and regroup all the terms in $I_1 + \dots + I_4$, we get that

$$Q = \sum_{i=1}^4 Q_i \quad (4.20)$$

where

$$Q_i = \int_0^t \int \int_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, y_1) Y(t-s, y_2) f(y_1-y_2) h_i(s, y_1, y_2). \quad (4.21)$$

Thus

$$\|I(t, x) - I(t, z)\|_p^2 \leq 4p \sum_{i=1}^4 \|Q_i\|_{p/2}.$$

We now estimate each $\|Q_i\|_{p/2}$ separately.

Upper bound for $\|Q_1\|_{p/2}$. Using the Lipschitz property of σ , Minkowski's inequality and the triangle inequality gives is

$$\|Q_1\|_{p/2} \leq L(\sigma)^2 \int_0^t ds \int \int_{\mathbb{R}^{2d}} dy_1 dy_2 Y(t-s, y_1) Y(t-s, y_2) f(y_1-y_2) \left[\sup_{|z_1-z_2|=\xi} \|u(s, z_1) - u(s, z_2)\|_p^2 \right]$$

$$\begin{aligned}
&\leq L(\sigma)^2 CK_f \left(\frac{\nu}{2}\right)^{-\lambda/\alpha} \int_0^t ds (t-s)^{2(\beta+\gamma-1)-\beta\lambda/\alpha} \left[\sup_{|z_1-z_2|=\xi} \|u(s, z_1) - u(s, z_2)\|_p^2 \right] \\
&\leq L(\sigma)^2 CK_f \left(\frac{\nu}{2}\right)^{-\lambda/\alpha} (t)^{2(\beta+\gamma-1)-\beta\lambda/\alpha} \int_0^t ds \left[\sup_{|z_1-z_2|=\xi} \|u(s, z_1) - u(s, z_2)\|_p^2 \right],
\end{aligned}$$

where the second inequality above follows from Lemma ??.

To bound the remaining $Q'_i s$, we will need to consider the following truncations which will then allow us to apply Lemma 4.12. For $k \geq 1$ we define $\Sigma_x^k(s, y) = \Sigma_x(s, y) 1_{\{\Sigma_x(s, y) \leq k\}}$ and $\Sigma_{x,z}^k(s, y) = \Sigma_{x,z}(s, y) 1_{\{\Sigma_{x,z}(s, y) \leq k\}}$ and then we define Q_i^k the same as Q_i but with h_i replaced with h_i^k where h_i^k is defined by replacing $\Sigma_x(s, y)$ and $\Sigma_{x,z}(s, y)$ with $\Sigma_x^k(s, y)$ and $\Sigma_{x,z}^k(s, y)$ respectively in h_i .

Upper Bound for $\|Q_2^k\|_{p/2}$. We apply Lemma 4.12 with $\phi(y_2) = \Sigma_x^k(s, y_2)$ and $\psi(y_1) = \Sigma_{x,z}^k(s, y_1)$. Thus we have that up to a constant of $(2\pi)^{-d}$,

$$Q_2^k = \int_0^t ds \int_{\mathbb{R}^d} \mu(d\zeta) \overline{\mathcal{F}(\Sigma_x^k(s, \cdot) Y(t-s, \cdot))(\xi)} \mathcal{F}(\Sigma_{x,z} Y(t-s, \cdot))(\xi) [e^{-i\xi \cdot \zeta} - 1],$$

where we recall that $\zeta = x - z$. □

References

- [1] CH2017 Chen L. and Huang J. (2017). Comparison Principle for SHE on $\mathbb{R}^d$. *The Annals of Probability* **Vol. 47, No. 2** 989 – 1035.
- [2] ChenHu21 Chen, L and Hu, G. (2021). Hölder regularity of the nonlinear stochastic time-fractional slow and fast diffusion equations on $\mathbb{R}^d$. *arXiv:2105.00891*.
- [3] CHHH16 Chen, L., Hu, G., Hu, Y. and Huang, J. (2016) Space-time fractional diffusions in Gaussian noisy environment *Stochastics*, 89:1, 171–206.
- [4] CHN19 Chen, L., Hu, Y., and Nualart, D. (2019). Nonlinear stochastic time-fractional slow and fast diffusion equations on $\mathbb{R}^d$. *Stochastic Process. Appl.* **129**, 5073–5112
- [5] FN2016 Foondun, M and Nane E. (2016) Asymptotic properties of some space-time fractional stochastic equations. *Mathematische Zeitschrift* 2015, **287**, 493 – 519
- [6] FT2014 Foondun, M. and Tian, K. (2014) On some properties of a class of fractional stochastic heat equations. *arXiv:1404.6791v1*
- [7] HHN13 Hu, Y., Huang, J. and Nualart, D. (2013) On Hölder continuity of the solution of stochastic wave equations *arXiv:1308.6607v1*
- [8] Kilbas [8] H-Transforms: Theory and Applications *An International Series of Monographs in Mathematics*.
- [9] Kuo [9] Kuo, H. (2000) *Introduction to Stochastic Integration* Springer, Universitext
- [10] Podlubny99 Podlubny, I. (1999). *Fractional Differential Equations*. Mathematics in Science and Engineering, Academic Press, Vol198 (1999).
- [11] eMillet2021 Sanz-Sole, M. and Millet, A. (2021) Global solutions to stochastic wave equations with superlinear coefficients. *Stochastic Processes and their Applications*, Elsevier, 2021, 139, pp.175–211.

Walsh89

- [12] Walsh, J. (1989) *An Introduction to Stochastic Partial Differential Equations*. École d'été de Probabilités de Saint-Flour, XIV—1994. Lecture Notes in Math., vol. 1180, pp. 265-439. Springer, Berlin (1986).