

CHEM 1405 Notes

MUFARO MACHAYA

Summer 2024

These are my notes from CHEM 1405 with Professor Priyanka Prashant, which I took at Lone Star College-CyFair (online). I needed the course to enter Science One at The University of British Columbia, haha. Maybe someone else will find them more useful than I did. I made them with **Classnotes** by Neil Rathi. Kudos to him.

Contents

I	Reference Materials	I
I.I	Common Elements	I
I.II	Common Polyatomic Ions	V
1	Quantitative Skills	2
1.1	Lecture 1: Introduction	2
1.1.1	<i>The Scientific Method</i>	2
1.1.2	<i>Qualitative vs. Quantitative Data</i>	3
1.1.3	<i>Scientific Facts, Laws, Hypotheses, and Theories</i>	3
1.2	Lecture 2: Measurement	4
1.2.1	<i>Exact vs. Inexact and Accuracy vs. Precision</i>	4
1.2.2	<i>Significant Figures and Rounding-Off Rules</i>	4
	<i>Rounding Rules</i>	5
1.2.3	<i>Significant Figures and Mathematical Operations</i>	5
1.2.4	<i>Scientific Notation</i>	6
1.3	Lecture 3: Unit System and Dimensional Analysis	6
1.3.1	<i>The Metric System</i>	6
1.3.2	<i>SI Units</i>	7
1.3.3	<i>Metric to Metric Conversion</i>	8
1.3.4	<i>Common Equation Reference</i>	8
1.3.5	<i>Densities of Common Substances at 25 degrees Celsius</i>	9
1.3.6	<i>Temperature Conversions</i>	9
2	Matter	10
2.1	Lecture 1: Basic Concepts About Matter	10
2.1.1	<i>Matter</i>	10
2.1.2	<i>The Three States of Matter</i>	10
2.1.3	<i>The Properties of Matter</i>	11
2.1.4	<i>Changes in Matter</i>	12
	<i>Naming Changes Between Matter States</i>	12
2.1.5	<i>Pure Substances and Mixtures</i>	13

CONTENTS

2.1.6	<i>Elements and Compounds</i>	14
3	Elements	15
3.1	Lecture 1: Subatomic Particles and Isotopes	15
3.1.1	<i>The Atomic Theory of Matter</i>	15
3.1.2	<i>Molecules</i>	15
	Molecule Types	16
3.1.3	<i>Atoms</i>	16
3.1.4	<i>Subatomic Particles</i>	17
	Nuclear Notation	18
3.1.5	<i>Isotopes</i>	19
	Isobars	20
3.1.6	<i>Atomic Mass</i>	20
3.2	Lecture 2: Electronic Structure and Chemical Periodicity	20
3.2.1	<i>The Periodic Table</i>	20
	Representative Elements	21
	Alkali Metals (Group IA)	22
	Alkaline Earth Metals (Group IIA)	22
	Halogens (Group VIIA)	22
	Noble Gases	22
	Transition Element	23
	Inner Transition Element	23
3.2.2	<i>Physical Properties of Table Elements</i>	23
	Metals	23
	Non-Metals	23
3.2.3	<i>Periodic Table Trends</i>	24
	Metallic Character	24
	Non-Metallic Character	24
	Metalloids and Semiconductors	24
	Atomic Size	25
	Electronegativity	25
3.2.4	<i>Electron Shells, Subshells, and Orbitals</i>	25
	Base Review of Electrons	25
	Electron Shells	26
	Electron Subshells	26
	Electron Orbitals	27
	The Aufbau Principle	27
3.2.5	<i>Electron Configuration</i>	27
	Condensed Electron Configuration	28
3.2.6	<i>Paramagnetic and Diamagnetic Substances</i>	28

CONTENTS

Visualizing Electron Orbitals	29
4 Compounds	30
4.1 Lecture 1: Ionic and Covalent Compounds	30
4.1.1 <i>Types of Chemical Bonds</i>	30
The Octet Rule	31
Lewis Dot Symbols	31
4.1.2 <i>Ionic Bonds</i>	31
Ionic Compound Formulas	32
Ionic Compounds	33
4.1.3 <i>Covalent Bonds</i>	33
4.1.4 <i>Molecular Geometry and VSEPR</i>	34
The Core of Molecular Geometry based on VSEPR . . .	34
Types of VSEPR Configurations	35
4.1.5 <i>Bond Polarity</i>	36
Calculating Bond Polarity	36
4.1.6 <i>Molecular Polarity</i>	37
4.2 Lecture 2: Naming of Compounds	37
4.2.1 <i>Ionic Compounds</i>	38
Metal Ions	38
4.2.2 <i>Naming Binary Ionic Compounds</i>	38
4.2.3 <i>Naming Compounds with Polyatomic Ions</i>	39
4.2.4 <i>Binary Molecular Compounds</i>	40
Common Names to Know	40
4.2.5 <i>Introduction to Acids</i>	41
Acid Nomenclature	41
4.2.6 <i>Common Student Mistakes</i>	41
5 Reactions	43
5.1 Lecture 1: Reactions	43
5.1.1 <i>Introduction to Reactions</i>	43
The Law of Conservation of Mass	43
5.1.2 <i>Chemical Equations</i>	43
Balancing Equations	44
5.1.3 <i>Types of Chemical Reactions</i>	46
6 Moles	47
6.1 Lecture 1: Counting Atoms in Moles	47
6.1.1 <i>The Law of Definite Proportions/Constant Composition</i> . .	47
6.1.2 <i>Percent Composition</i>	47

CONTENTS

	Percent Composition from Known Formulas	47
	Percent Composition from Experimental Data	48
6.1.3	<i>Counting Atoms</i>	48
6.1.4	<i>Empirical Formula vs. Molecular Formula</i>	48
	Calculating Empirical Formulas	49
	Calculating Molecular Formulas	50
6.2	Lecture 2: Stoichiometry	51
6.2.1	<i>Calculating Theoretical Yield</i>	51
6.2.2	<i>Percent Yield</i>	52
7	Intermolecular Forces	53
8	Solutions	54

PERIODIC TABLE OF ELEMENTS

1 H 1.008																	2 He 4.003
3 Li 6.941	4 Be 9.012											5 B 10.81	6 C 12.011	7 N 14.007	8 O 15.999	9 F 18.998	10 Ne 20.180
11 Na 22.990	12 Mg 24.305											13 Al 26.981	14 Si 28.085	15 P 30.974	16 S 32.066	17 Cl 35.453	18 Ar 39.948
19 K 39.098	20 Ca 40.078	21 Sc 44.956	22 Ti 47.88	23 V 50.941	24 Cr 51.996	25 Mn 54.938	26 Fe 55.847	27 Co 58.933	28 Ni 58.69	29 Cu 63.546	30 Zn 65.39	31 Ga 69.72	32 Ge 72.61	33 As 74.922	34 Se 78.96	35 Br 79.904	36 Kr 83.80
37 Rb 85.468	38 Sr 87.62	39 Y 88.906	40 Zr 91.224	41 Nb 92.906	42 Mo 95.94	43 Tc (98)	44 Ru 101.07	45 Rh 102.905	46 Pd 106.42	47 Ag 107.868	48 Cd 112.41	49 In 114.82	50 Sn 118.710	51 Sb 121.757	52 Te 127.60	53 I 126.905	54 Xe 131.29
55 Cs 132.905	56 Ba 137.33	57 *La 138.905	72 Hf 178.49	73 Ta 180.948	74 W 183.85	75 Re 186.207	76 Os 190.2	77 Ir 192.22	78 Pt 195.08	79 Au 196.967	80 Hg 200.59	81 Tl 204.383	82 Pb 207.2	83 Bi 208.980	84 Po (209)	85 At (210)	86 Rn (222)
87 Fr (223)	88 Ra 226.025	89 †Ac 227.028	104 Rf (261)	105 Db (262)	106 Sg (263)	107 Bh (262)	108 Hs (265)	109 Mt (266)	110 Ds (271)	111 Rg (272)	112 Cn	113 Nh	114 Fl	115 Mc	116 Lv	117 Ts	118 Og

*LANTHANIDES	58 Ce 140.12	59 Pr 140.908	60 Nd 144.24	61 Pm (145)	62 Sm 150.36	63 Eu 151.96	64 Gd 157.25	65 Tb 158.925	66 Dy 162.50	67 Ho 164.930	68 Er 167.26	69 Tm 168.934	70 Yb 173.04	71 Lu 174.967
†ACTINIDES	90 Th 232.038	91 Pa 231.036	92 U 238.029	93 Np 237.048	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (257)	101 Md (258)	102 No (259)	103 Lr (260)

Module I

Reference Materials

I.I Common Elements

Symbol	Name	Group/Type	Charge	Charge Type
H	Hydrogen	Reactive Non-Metal	+1	Non-Metal
He	Helium	Noble Gas	-1	Non-Metal
Li	Lithium	Alkali Metal	+1	Fixed
Be	Beryllium	Alkaline Earth Metal	+2	Fixed
B	Boron	Metalloid	+3	Non-Metal
C	Carbon	Reactive Non-Metal	± 4	Non-Metal
N	Nitrogen	Pnictogen/Reactive Non-Metal	-3	Non-Metal
O	Oxygen	Chalcogen/Reactive Non-Metal	-2	Non-Metal
F	Flourine	Halogen/Reactive Non-Metal	-1	Non-Metal
Ne	Neon	Noble Gas	None	Noble Gas
Na	Sodim	Alkali Metal	+1	Fixed
Mg	Magnesium	Alkaline Earth Metal	+2	Fixed
Al	Aluminum	Post-Transition Metal	+3	Fixed
Si	Sillicon	Metalloid	± 4	Non-Metal
P	Phosphorus	Pnictogen/Reactive Non-Metal	-3	Non-Metal
S	Sulfur	Chalcogen/Reactive Non-Metal	-2	Non-Metal
Cl	Chlorine	Halogen/Reactive Non-Metal	-1	Non-Metal
Ar	Argon	Noble Gas	None	Noble Gas
K	Potassium	Alkali Metal	+1	Fixed
Ca	Calcium	Alkaline Earth Metal	+2	Fixed
Sc	Scandium	Transition Metal	N/A	Variable
Ti	Titanium	Transition Metal	N/A	Variable
Continued on next page				

MODULE I. REFERENCE MATERIALS

Table I.1 – continued from previous page

Symbol	Name	Group/Type	Charge	Charge Type
V	Vanadium	Transition Metal	N/A	Variable
Cr	Chromium	Transition Metal	N/A	Variable
Mn	Manganese	Transition Metal	N/A	Variable
Fe	Iron	Transition Metal	N/A	Variable
Co	Cobalt	Transition Metal	N/A	Variable
Ni	Nickel	Transition Metal	+2	Fixed
Cu	Copper	Transition Metal	N/A	Variable
Zn	Zinc	Transition Metal	+2	Fixed
Ga	Gallium	Post-Transition Metal	+3	Fixed
Ge	Germanium	Metalloid	N/A	Variable
As	Arsenic	Pnictogen/Metalloid	N/A	Variable
Se	Selenium	Chalcogen/Reactive Non-Metal	N/A	Variable
Br	Bromine	Halogen/Reactive Non-Metal		
Kr	Krypton	Noble Gas		
Rb	Rubidium	Alkali Metal		
Sr	Strontium	Alkaline Earth Metal		
Y	Yttrium	Transition Metal		
Zr	Zirconium	Transition Metal		
Nb	Niobium	Transition Metal		
Mo	Molybdenum	Transition Metal		
Tc	Technetium	Transition Metal		
Ru	Ruthenium	Transition Metal		
Rh	Rhodium	Transition Metal		
Pd	Palladium	Transition Metal		
Ag	Silver	Transition Metal		
Cd	Cadmium	Transition Metal		
In	Indium	Post-Transition Metal		
Sn	Tin	Post-Transition Metal		
Sb	Antimony	Pnictogen/Metalloid		
Te	Tellurium	Chalcogen/Metalloid		
I	Iodine	Halogen/Reactive Non-Metal		
Xe	Xenon	Noble Gas		
Cs	Cesium	Alkali Metal		
Ba	Barium	Alkaline Earth Metal		
La	Lanthanum	Lanthanoid		
Ce	Cerium	Lanthanoid		
Pr	Praseodymium	Lanthanoid		
Continued on next page				

MODULE I. REFERENCE MATERIALS

Table I.1 – continued from previous page

Symbol	Name	Group/Type	Charge	Charge Type
Nd	Neodymium	Lanthanoid		
Pm	Promethium	Lanthanoid		
Sm	Samarium	Lanthanoid		
Eu	Europium	Lanthanoid		
Gd	Gadolinium	Lanthanoid		
Tb	Terbium	Lanthanoid		
Dy	Dysprosium	Lanthanoid		
Ho	Holmium	Lanthanoid		
Er	Erbium	Lanthanoid		
Tm	Thulium	Lanthanoid		
Yb	Ytterbium	Lanthanoid		
Lu	Lutetium	Lanthanoid		
Hf	Hafnium	Transition Metal		
Ta	Tantalum	Transition Metal		
W	Tungsten	Transition Metal		
Re	Rhenium	Transition Metal		
Os	Osmium	Transition Metal		
Ir	Iridium	Transition Metal		
Pt	Platinum	Transition Metal		
Au	Gold	Transition Metal		
Hg	Mercury	Transition Metal		
Tl	Thallium	Post-Transition Metal		
Pb	Lead	Post-Transition Metal		
Bi	Bismuth	Pnictogen/Post-Transition Metal		
Po	Polonium	Chalcogen/Post-Transition Metal		
At	Astatine	Halogen/Metalloid		
Rn	Radon	Noble Gas		
Fr	Francium	Alkali Metal		
Ra	Radium	Alkaline Earth Metal		
Ac	Actinium	Actinoid		
Th	Thorium	Actinoid		
Pa	Protactinium	Actinoid		
U	Uranium	Actinoid		
Np	Neptunium	Actinoid		
Pu	Plutonium	Actinoid		
Am	Americium	Actinoid		
Cm	Curium	Actinoid		
Continued on next page				

MODULE I. REFERENCE MATERIALS

Table I.1 – continued from previous page

Symbol	Name	Group/Type	Charge	Charge Type
Bk	Berkelium	Actinoid		
Cf	Californium	Actinoid		
Es	Einsteinium	Actinoid		
Fm	Fermium	Actinoid		
Md	Mendelevium	Actinoid		
No	Nobelium	Actinoid		
Lr	Lawrencium	Actinoid		
Rf	Rutherfordium	Transition Metal		
Db	Dubnium	Transition Metal		
Sg	Seaborgium	Transition Metal		
Bg	Bohrium	Transition Metal		
Hs	Hassium	Transition Metal		
Mt	Meitnerium	Synthetic/Unknown		
Ds	Damstadtium	Synthetic/Unknown		
Rg	Roentgenium	Synthetic/Unknown		
Cn	Copernicium	Synthetic/Unknown		
Nh	Nihonium	Synthetic/Unknown		
Fl	Flerovium	Synthetic/Unknown		
Mc	Moscovium	Synthetic/Unknown		
Lv	Livemonium	Synthetic/Unknown		
Ts	Tennessine	Synthetic/Unknown		
Og	Oganesson	Synthetic/Unknown		

Note: <https://sciencenotes.org/element-charges-chart-how-to-know-the-charge-of-an/>

I.II Common Polyatomic Ions

Formula	Sign	Name
$\text{C}_2\text{H}_3\text{O}_2^-$	Anion	Acetate
NH_3	Not Ionic!	Ammonia
NH_4^+	Cation	Ammonium
AsO_4^{3-}	Anion	Arsenate
AsO_3^{3-}	Anion	Arsenite
N_3^-	Anion	Azide
$\text{C}_7\text{H}_5\text{O}_2^-$	Anion	Benzoate
BO_3^{3-}	Anion	Borate
BrO_3^-	Anion	Bromate
CO_3^{2-}	Anion	Carbonate
ClO_3^-	Anion	Chlorate
ClO_2^-	Anion	Chlorite
CrO_4^{2-}	Anion	Chromate
CN^-	Anion	Cyanide
$\text{Cr}_2\text{O}_7^{2-}$	Anion	Dichromate
H_2PO_4^-	Anion	Dihydrogen phosphate
H_2PO_3^-	Anion	Dihydrogen phosphite
HCO_3^-	Anion	Hydrogen carbonate
HPO_4^{2-}	Anion	Hydrogen phosphate
HPO_3^{2-}	Anion	Hydrogen phosphite
HSO_4^-	Anion	Hydrogen sulfate
HSO_3^-	Anion	Hydrogen sulfite
HS^-	Anion	Hydrogen sulfide
OH^-	Anion	Hydroxide
ClO^-	Anion	Hypochlorite
H_3O^+	Cation	Hydronium Ion
IO_3^-	Anion	Iodate
MnO_4^{2-}	Anion	Manganate
NO_3^-	Anion	Nitrate
NO_2^-	Anion	Nitrite
$\text{C}_2\text{O}_4^{2-}$	Anion	Oxalate
ClO_4^-	Anion	Perchlorate
MnO_4^-	Anion	Permanganate
O_2^{2-}	Anion	Peroxide
Continued on next page		

MODULE I. REFERENCE MATERIALS

Table I.2 – continued from previous page

Formula	Sign	Name
PO_4^{3-}	Anion	Phosphate Ion
PO_3^{3-}	Anion	Phosphite Ion
SiO_3^{2-}	Anion	Silicate
SO_4^{2-}	Anion	Sulfate Ion
SO_3^{2-}	Anion	Sulfite Ion
$\text{C}_4\text{H}_4\text{O}_6^{2-}$	Anion	Tartrate
SCN^-	Anion	Thiocyanate
$\text{S}_2\text{O}_3^{2-}$	Anion	Thiosulfate

Module 1

Quantitative Skills

1.1 Lecture 1: Introduction

References chapter 1, sections 1.2 and 1.3 of the textbook.

1.1.1 The Scientific Method

Definition 1.1.1. The Scientific Method is the set of general procedures for acquiring scientific knowledge.

1. Identify the problem
2. Collect past data/perform background research
3. Analyze and organize the data into general summaries of previous observations
4. Suggest probable explanations/hypotheses for the generalizations
5. Perform further experiments to prove or disprove these explanations/theorems

1.1.2 Qualitative vs. Quantitative Data

Definition 1.1.2. Qualitative vs. Quantitative Data

- **Quantitative Data** - Numerical data yielded from *measurements*.
- **Qualitative Data** - Any data yielded from other observation, *including rough estimates*.

- The patient's fever has reached 105.3°F - *Quantitative*
- The packet of Candy contains *about* 100 gummy bears - *Qualitative*

1.1.3 Scientific Facts, Laws, Hypotheses, and Theories

- **Scientific Fact** - Reproducible data/observations obtained from an experiment.
- **Scientific Law** - Summarizes scientific facts without interpretation. Accepted as fact but can be changed/contested.
- **Scientific Hypothesis** - Explains or contests scientific laws. Tested by experiments.
- **Scientific Theory** - Scientific hypotheses that are tested and validated as correct/true over a long period of time. *Scientific Theories can never be proven true.*

Example 1.1.1. Facts, Laws, and Theories

- The burning candle generated both heat and light - *Scientific Fact*
- As a candle burns, its wax gradually disappears - *Scientific Fact*
- All burning candles generate heat and light - *Scientific Law*
- Burning candles generate heat as the result of the decomposition of melted wax. - *Scientific Hypothesis*

1.2 Lecture 2: Measurement

1.2.1 Exact vs. Inexact and Accuracy vs. Precision

Definition 1.2.1. Exact vs. Inexact and Accuracy vs. Precision

- **Exact numbers** - Numbers with no uncertainty and are known exactly either in definition, counting, or as simple fractions (cannot be an irrational number).
- **Inexact numbers** - Numbers with some degree of uncertainty.
- **Accuracy** - How close the measured values are to the intended/-goal values
- **Precision** - How close the measured values are to each other

- Measurements require units.
- It is impossible to make exact measurements as all measurements have some degree of uncertainty due to limits of the measuring instrument (systematic error) and random error.

When measuring, one has to keep the certain digits and uncertain digits in mind. The magnitude of uncertainty is indicated with a plus-minus notation alongside the acceptable range.

1.2.2 Significant Figures and Rounding-Off Rules

Significant figures are all the certain digits of measurement plus one digit of uncertainty. When determining significant figures for unknown measurements, the following rules are used:

- Nonzero digits are always significant.
- Leading zeroes are never significant.
- Confined zeroes are always significant.
- Trailing zeroes are only significant if there is a decimal point or the zeroes carry overbars (i.e. are repeating).

Note: Remember that the last significant figure is always uncertain.

MODULE 1. QUANTITATIVE SKILLS

Example 1.2.1. Significant Figures

- $14.232 \rightarrow 5 \text{ sig. figs, } \pm 0.001$
- $0.0045 \rightarrow 2 \text{ sig. figs, } \pm 0.0001$
- $2.075 \rightarrow 4 \text{ sig. figs, } \pm 0.001$
- $4300.00 \rightarrow 6 \text{ sig. figs, } \pm 0.01$
- $36,003 \rightarrow 5 \text{ sig. figs, } \pm 1$
- $6310 \rightarrow 3 \text{ sig. figs, } \pm 10$

Rounding Rules

1. If the first digit to be dropped is less than five, then all proceeding digits are just dropped. (ex. $98.623 \rightarrow 98.6$)
2. If the first digit to be dropped is greater than five or is a five followed by all zeroes, the previous digit is increased by one. (ex. $25.0566 \rightarrow 25.06$)
3. If the first digit to be dropped is a five not followed by any other digit or a five followed by only zeroes, then the previous digit is increased by one only if it is odd. If it is even, it will not be incremented. (ex. $63.3500 \rightarrow 63.4$, but $62.65 \rightarrow 62.6$)

1.2.3 Significant Figures and Mathematical Operations

- For multiplication and division, always use the *fewest significant figures*.
- For addition and subtraction, always use the number with the most uncertainty (or the number with the uncertainty digit in the highest place).

Example 1.2.2. Significant Figures with Operations

- $6.038 \times 2.57 \approx 15.5$
- $677 + 39.2 + 6.23 \approx 722$

1.2.4 Scientific Notation

Definition 1.2.2. Scientific notation is a numerical system in which an ordinary decimal number is expressed as a product of a number between 1 and 10 and 10 raised to a power. (as in: $A \times 10^N$) The value N refers to how many places the decimal of a number is moved, where $+N$ is right and $-N$ is left.

Example 1.2.3. Scientific Notation

- $905000 = 9.05 \times 10^5$
- $0.0006030 = 6.030 \times 10^{-4}$
Note: The trailing zero is kept as it is significant.
- $1.894 \times 10^{-7} = 0.0000001894$
- $1.327 \times 10^5 = 132700$

1.3 Lecture 3: Unit System and Dimensional Analysis

1.3.1 The Metric System

Quantity	Metric System	Customary System	Tool for Measure
Mass	Grams (g)	Pounds (lbs)	Scale
Length	Meters (m)	Inches (in.)/Various	Ruler
Volume	Liters (L)	Fluid Ounces (fl. oz.)/Various	Graduated Cylinder
Temperature	Celsius (°C)	Fahrenheit (°F)	Thermometer

Note: The metric system is based on base 10, meaning that it follows prefixes that get multiplied to each number by a series of ten.

MODULE 1. QUANTITATIVE SKILLS

Prefix	Symbol	Value
Tera	T	10^{12}
Giga	G	10^9
Mega	M	10^6
Kilo	k	10^3
Hecto	h	10^2
Deca	da	10^1
Deci	d	10^{-1}
Centi	c	10^{-2}
Milli	m	10^{-3}
Micro	μ	10^{-6}
Nano	n	10^{-9}
Pico	p	10^{-12}

Note: This can be remembered with: *The Great Mega King Henry Doesn't [usually] Drink Chocolate Milk Mixed [with] Nana's Peanuts*. **It is incredibly important that these values and their prefixes are remembered!**

1.3.2 SI Units

SI is the international system of units, and denotes the standard units for each measurement.

- SI Length is in Meters (m) ("Metres" internationally but "Meters" domestically).
- SI Mass is in Kilograms (kg). **Note:** Mass is the total number of matter, weight is mass multiplied by the force of gravity.
- SI Volume is in Liters (L). ("Litres" internationally but "Liters" domestically), where $1 \text{ L} = 1000 \text{ cm}^3$.
- SI time is in seconds (s).
- SI temperature is in Kelvin (K).
- SI Current is in Amperes (amp).
- SI Amount/Count is in Moles (mol).

MODULE 1. QUANTITATIVE SKILLS

Note: When measuring the volume of a solid, use $V = LWH$. For liquids in a graduated cylinder, read from the bottom of the meniscus.

1.3.3 Metric to Metric Conversion

1. Determine what the end goal unit is.
2. Determine what the given unit is.
3. Multiply the given unit by the ratio fraction of the two units, where the initial unit's value is the denominator and the ending unit's value is the numerator.
4. Round the final product to the original number of significant figures.

For English-to-Metric conversions, you will be given the conversion rates between English and Metric conversions. That is all that you need to know for that.

1.3.4 Common Equation Reference

- Density: $\rho = \frac{\text{Mass}}{\text{Volume}}$
- Percent Error: $\frac{\text{Measured Value} - \text{Accepted Value}}{\text{Accepted Value}} \times 100$ (**Note:** Percent error can be negative, as the sign indicates the relation between the measured and the intended value.)
- Rate or Velocity: $v = \frac{\text{Distance}}{\text{Time}}$
- Concentration: $M = \frac{\text{Moles}}{\text{Volume}}$

MODULE 1. QUANTITATIVE SKILLS

1.3.5 Densities of Common Substances at 25 degrees Celsius

Substance	Density (g/mL)	Substance	Density (g/cm ³)
Ethanol	0.789	Balsa Wood	0.160
Acetone	0.791	Cork	0.210
Rubbing Alcohol	0.786	Magnesium	1.74
Motor Oil	0.899	Aluminum	2.70
Olive Oil	0.920	Iron	7.86
Water	1.00	Copper	8.92
Seawater	1.03	Silver	10.5
Honey	1.45	Lead	11.34
Bromine	3.12	Gold	19.32
Mercury	13.6	Platinum	21.45

Note: Cubic centimeters equal milliliters.

1.3.6 Temperature Conversions

- $T_K = T_C + 273.15^\circ$
- $T_C = \frac{5}{9}(T_F - 32)$
- $T_F = \frac{9}{5}T_C + 32$
- $273^\circ K = 0^\circ C$
- $373^\circ K = 100^\circ C$
- $32^\circ F = 0^\circ C$
- $212^\circ F = 100^\circ C$

Module 2

Matter

2.1 Lecture 1: Basic Concepts About Matter

2.1.1 Matter

Definition 2.1.1. *Matter* is anything that has mass and occupies space. The *mass* of an object is the measure of the amount of matter in an object, and the *volume* of an object is the amount of space the object takes up in three-dimensional space.

Note: Matter includes living, nonliving, viewable, and non-viewable things, such as wood, rocks, gasoline, air, bacteria, etc.

2.1.2 The Three States of Matter

- **Solids** have *definite shape and definite volume*, and have *the least energy* out of the three states.
- **Liquids** have *indefinite shape but definite volume* by taking the shape of its container, and thus, they have *more energy than solids but less energy than gases*.
- **Gases** have *indefinite shape and indefinite volume* as they expand to take the shape and volume of their container, and thus, they have *the most energy* of the three types.

Note: As the temperature of matter increases, particles move faster as they have more energy. Particles in solids have a lower/near-zero velocity, but

MODULE 2. MATTER


particles in gases or liquids will have higher velocities and thus higher energies.

2.1.3 The Properties of Matter

The properties of matter are the distinguishing characteristics of a substance, and they are useful for the identification/classification of unknown substances, characterization of a newly-discovered substance, distinguishing between different substances, and predicting the usefulness of a substance for a specific purpose.

Note: Properties of Matter are split into two types: **Physical** and **Chemical**.

Definition 2.1.2. A **physical property** is a characteristic of a substance that can be observed without changing the substance into another substance *chemically*. A **chemical property** describes the way a substance undergoes a change into another substance.



© Encyclopædia Britannica, Inc.

Figure 2.1: Properties of Matter Flowchart. **Note:** Remember the difference between Intensive and Extensive physical properties

2.1.4 Changes in Matter

Physical changes to matter change a substance's physical appearance but not its chemical composition, and chemical changes affect its chemical composition (and usually additionally change its physical qualities in the process) **to form a completely new substance**. Physical changes include melting ice, forming clouds, cutting, breaking, etc. Chemical changes include heating/cooking, reacting a substance with another substance, healing a wound, rusting, fermentation, etc.

Note: Remember that the key difference between physical and chemical changes is whether or not a new substance is formed by the change.

Note: Dissolving salt into water is actually a chemical change because when salt dissolves, it dissociates into ions, thus changing its chemical composition, but dissolving sugar into water is physical because the sugar simply changes from a solid state to an aqueous state.

Naming Changes Between Matter States

The movement from each state of matter to another has a completely different term based on the nature of the change.




Figure 2.2: States of Matter Flowchart

2.1.5 Pure Substances and Mixtures

Matter can be classified into one of two types: mixtures or pure substances. Pure substances are composed of a singular substance and cannot be physically separated (think like Pure Gold). Whereas mixtures are comprised of more than one substance and can be physically separated into its component substances.




Figure 2.3: Pure Substance vs. Mixtures Flowchart

- *Pure Substances* only ever have one substance present, and thus, have a definite and constant composition and consistent chemical and physical properties. **Note:** Pure substances are always chemically homogeneous, but they can be physically homogeneous or heterogeneous.
- *Mixtures* are combinations of more than one type of substance, and thus, have variable composition and properties. **Note:** Mixtures are always chemically heterogeneous but, like pure substances, can be phys-

MODULE 2. MATTER

ically homogeneous or heterogeneous.

- *Elements* are molecules with a singular *type* of atom.
- *Compounds* are molecules with two or more types of atoms joined together.
- *Homogeneous Mixtures* are mixtures where a substance (known as the solute, such as salt or sugar) is dissolved into another substance (known as the solvent, such as water) at a *uniform distribution* (think saltwater). These mixtures are also known as **solutions** and have one distinct physical appearance (or phase).
- *Heterogeneous Mixtures* are non-solution mixtures with a non-uniform distribution of its comprising particles and thus, two or more distinct physical appearances/phases (think Chex Mix).

2.1.6 Elements and Compounds

Definition 2.1.3. Elements are the building blocks of all higher types of matter and are made from atoms. They are the smallest classification of substances, and thus, cannot be broken down into simpler substances through chemical or physical means. **Note:** There are currently 118 known atom types, and 90 of those types occur naturally in nature.

Module 3

Elements

3.1 Lecture 1: Subatomic Particles and Isotopes

Matter consists of various mixtures of substances comprised of molecules of compounds comprised of elements comprised of atoms, which are in turn comprised of protons, neutrons, and electrons.

3.1.1 The Atomic Theory of Matter

1. All matter is made of atoms, and each distinct type of atom is known as an element. (118 different types of which are currently known.)
2. All atoms of a given type will be similar to one another and different from all others.
3. Relative number and arrangement of different types of atoms in a pure substance determine its identity.
4. Chemical change is a *union, separation, or rearrangement* of atoms to produce new substances.
5. Only whole atoms can participate in or result from any chemical change, and **for the purposes of this course, atoms are considered indestructible during chemical changes.**

3.1.2 Molecules

MODULE 3. ELEMENTS

Definition 3.1.1. Molecules are made from two or more atoms bonded **covalently** to function as a singular entity, known as a *molecular compound*. In a molecule, *the relative number and arrangement of its atoms determine its identity*.

Note: Whereas the atom is the limit of chemical subdivision, the molecule is the limit of physical subdivision.

- There is only ever one type of molecule for any given molecular substance (otherwise it is a mixture).
- Molecules have their own properties which will be different from their individual atoms.

Molecule Types

- Diatomic Molecule: Two atoms
- Polyatomic Molecule: More than one atoms
- Homoatomic Molecule: The atoms in the molecules are the same type
- Heteroatomic Molecule: The atoms in the molecules are different types

Note: There are seven elements that naturally exist as diatomic molecules: Hydrogen (H_2), Nitrogen (N_2), Oxygen (O_2), Fluorine (F_2), Chlorine (Cl_2), Bromine (Br_2), and Iodine (I_2).

- Molecules containing Carbon are called **organic**.
- Molecules containing **only** Carbon, Hydrogen, and Oxygen are called **carbohydrates**.
- Molecules that only contain Carbon and Hydrogen are called **Hydrocarbons**.

3.1.3 Atoms

- Atoms are always neutral, as they always have the same number of protons and electrons. This balance is broken in **ions**.
- Atoms make up elements and compounds.

MODULE 3. ELEMENTS

- Individual atoms are smaller than the wavelength of light, so they cannot be seen with an optical microscope.
- The mass of the atom lies solely in the nucleus, but the vast majority of the atom's size is in the electron cloud. (The ratio of the size of the electron cloud to the nucleus is 1 to 100,000.)




Figure 3.1: Chadwick's Model of an Atom

3.1.4 Subatomic Particles

MODULE 3. ELEMENTS

Definition 3.1.2. Atoms are comprised of smaller fundamental particles known as subatomic particles. The three core subatomic particles are **Protons, Neutrons, and Electrons**.

- Protons have a positive charge and weigh slightly less (but effectively equal to) a Neutron.
- Neutrons have no/neutruual charge and weigh the most out of the previous two particles.
- Electrons have a negative charge and weigh roughly 1800 times less than Protons and Neutrons, with effectively no mass.

The atom is split into two key portions, the nucleus and the electron cloud. The nucleus is the small, incredibly dense center of the atom that contains its protons and neutrons (which *always results in a positive charge*), and these protons and neutrons within the nucleus are referred to as *nucleons*. The electron cloud is the large outer region of the atom where the electrons move freely and rapidly throughout the nucleus, *thus always resulting in a negative charge*. It has no firmly defined area for any atom and contains various shell and subshell levels which contain orbitals for the electrons.

Nuclear Notation

Nuclear Notation is used to indicate the atomic number and mass number of an atom of isotope, and it is formed by writing an elemental symbol preceded by a subscript indicating its atomic number (Z) and a superscript indicating its mass number (A).

Note: Together, an atom with a specific atomic number and a specific mass number form a **nuclide**.

Nuclear Notation




Figure 3.2: Nuclear Notation Model

Definition 3.1.3. The Atomic Number, denoted by Z , is the number of protons in the nucleus of the atom, and it acts as the primary definition of each type of atom (or Element). Atomic Numbers 1 through 92 (excluding Technetium and Promethium) are naturally occurring and the remaining 93 through 118 are synthetic. **Note:** As Atoms always hold a neutral charge, the number of electrons will always equal the number of protons and the atomic number.

The Mass Number is the sum of the number of protons and neutrons within the nucleus of an atom, and it is a count, **not** an atomic mass.

3.1.5 Isotopes

MODULE 3. ELEMENTS

Definition 3.1.4. Isotopes are atoms of an element that have the same number of protons and electrons but a different number of neutrons, and they are named by denoting the name of the element followed by its newfound mass number (such as Cobalt-60 or Carbon-14).

As an element is determined by the number of protons in the nucleus, the nucleus can have a variable number of neutrons without changing the charge but resulting in changes to the atom's mass number, and this change results in slightly different physical properties but generally similar chemical properties.

Note: The mass number minus the atomic number is equal to the number of neutrons present in an isotope.

Isobars

Definition 3.1.5. Isobars are atoms/nucleides of different chemical elements that have the same number of nucleons. Examples include Iron-58 ($^{58}_{26}\text{Fe}$) and Nickel-58 ($^{58}_{27}\text{Ni}$), or Calcium-40 ($^{40}_{20}\text{Ca}$) and Argon-40 ($^{40}_{18}\text{Ar}$).

3.1.6 Atomic Mass

Definition 3.1.6. The atomic mass is a weighted average that accounts for the number of isotopes for the element, the percent abundance of each isotope, and the relative mass of each isotope (with Carbon-12 as the reference point).

3.2 Lecture 2: Electronic Structure and Chemical Periodicity

References chapter 4, sections 4.4 to 4.7 of the textbook.

3.2.1 The Periodic Table

MODULE 3. ELEMENTS

Definition 3.2.1. The Periodic Law

When elements are arranged in order of increasing atomic number, elements with similar chemical behavior occur at regularly recurring (or *periodic*) intervals.

- Period: The horizontal row of elements in the periodic table.
- Group: The vertical column of elements in the periodic table.
 - There are three ways to name the columns: US, EU, and IUPAC
 - There are four groups with non-numerical names: Alkali Metals, Alkali Earth Metals, Halogens, and Noble Gases
- Chemical Periodicity: The variation in properties of elements as a function of their position on the periodic table (or how atoms change as they travel up/down the periodic table).
 - There are three properties of elements that exhibit chemical periodicity: metallic vs. non-metallic character, atomic size, and electronegativity.

Note: Increasing atomic number **does not** always equal increasing atomic mass.

Elements in the periodic table can be classified in one of two ways: either by their electron configurations (noble gases, representative elements, transition metals, or inner transition/rare earth metals) or their physical properties (metals/non-metals/metalloids)

Representative Elements

Definition 3.2.2. Representative elements are all of the elements located in the first two columns (denoted as the $\langle s \rangle$ area) or columns 13-17 (denoted as the $\langle p \rangle$ area).

Most elements are representative elements, and they generally have a partially or completely full $\langle s \rangle$ electron level and either an empty or partially filled $\langle p \rangle$ electron level.

MODULE 3. ELEMENTS

Alkali Metals (Group IA)

- Members: Lithium (Li), Sodium (Na), Potassium (K), Rubidium (Rb), and Cesium (or Caesium) (Cs)
- They only have one valence electron.
- All soft, shiny metals.
- All very reactive with water.

Alkaline Earth Metals (Group IIA)

- Members: Beryllium (Be), Magnesium (Mg), Calcium (Ca), Strontium (St), and Barium (Ba)
- They all have two valence electrons.
- All soft, shiny metals.
- Only moderately reactive with water.

Halogens (Group VIIA)

- Members: Fluorine (F), Chlorine (Cl), Bromine (Br), Iodine (I), and Astatine (At)
- They all have seven valence electrons.
- They all exist as diatomic molecules.
- They are all very reactive colored substances.
- Almost all are all gases at room temperature (or slightly above room temperature), except for Bromine, which is a liquid at room temperature.

Noble Gases

- Members: Helium, Neon, Argon, Krypton, Xenon, and Radon
- They all have a full valence shell of electrons.
- They are all completely unreactive (hence the name "Noble," as they are too noble to react with other substances).
- They are all gases at room temperature.

MODULE 3. ELEMENTS

Transition Element

Definition 3.2.3. Transition elements are located in the $\langle d \rangle$ area of the periodic table, and their electrons fill the $\langle D \rangle$ energy level.

Inner Transition Element

Definition 3.2.4. Inner Transition Elements are located in the $\langle f \rangle$ area of the periodic table, and their electrons fill the $\langle f \rangle$ energy level. This series are referred to as **Lanthanides and Actinides** and are generally known as "Rare Earth Elements."

3.2.2 Physical Properties of Table Elements

Metals

- Luster (Shine)
- Thermal Conductivity
- Electrical Conductivity
- Malleability (can be rolled into sheets)
- Ductility (can be drawn into wires)
- All solids at room temperature (except for Mercury)
- High Density
- High Melting Point

Note: There are 94 actively known metals.

Non-Metals

Note: Non-metals are generally characterized by their lack of the properties of *luster, thermal conductivity, electrical conductivity, and malleability* in reference to metals.

- Can be solids, liquids, or gases at room temperature.

MODULE 3. ELEMENTS

- Nitrogen (N), Oxygen (O), Hydrogen (H), Chlorine (Cl), Fluorine (F), Helium (He), Neon (Ne), Argon (Ar), Krypton (Kr), Xenon (Xe), and Radon (Rn) are all **gases**.
- Carbon (C), Phosphorus (P), Sulfur (S), Selenium (Se), Iodine (I), Astatine (At), and Tellurium (Te) are all **solids**.
- Bromine (Br) is a **liquid**.
- Generally have lower density and lower melting points than metals.

3.2.3 Periodic Table Trends

Metallic Character

Definition 3.2.5. Metallic character is a general chemical property that describes how 'metallic' an element is, and for any given period/row in the periodic table, it increases from right to left, and for any given group, it increases going top to bottom. For example, Sodium (Na) is more Metallic than Magnesium (Mg) but less metallic than Potassium (K).

Non-Metallic Character

Definition 3.2.6. Non-metallic character is another general chemical property that describes how 'non-metallic' an element is (and subsequently, it works in the exact opposite manner to metallic character). As such, for any given period, it increases from left to right, and for any given group, it increases going bottom to top. For example, Chlorine (Cl) is more non-metallic than Phosphorus (P) but less non-metallic than Fluorine (F).

Metalloids and Semiconductors

Definition 3.2.7. Metalloids are an element whose properties intermediate between those of metals and non-metals, and there are eight known metalloids: Boron (B), Silicon (Si), Germanium (Ge), Arsenic (As), Antimony (Sb), Tellurium (Te), Polonium (Po), and Astatine (At).

MODULE 3. ELEMENTS

Note: Metalloids reside along the stepped line which divides the metals and non-metals.

Definition 3.2.8. Semiconductors are metalloids that do not conduct electrical currents at room temperatures but are able to conduct electrical currents at higher temperatures, and there are three main semiconductors: Silicon (Si), Germanium (Ge), and Antimony (Sb).

Note: Semiconductors are vital in the electronics industry.

Atomic Size

Atomic Radii tend to increase from right to left within any given period and increase from top to bottom for any given group. For example, Helium is roughly the smallest whereas Francium is roughly the largest.

Electronegativity

Definition 3.2.9. Electronegativity is a measure of the relative attraction that an atom has for the shared electrons within its bonds.

- The higher the electronegativity of an element, the greater the electron-attracting ability of the atoms of that element.
- Electronegativity generally increases from left to right within a given period and from bottom to top within a given group.
- Electronegativity depends on numerous factors: atomic size, nuclear charge, and the number of non-valence electrons.

3.2.4 Electron Shells, Subshells, and Orbitals

Base Review of Electrons

- Electrons are the smallest of the three main subatomic particles.
- Electrons have very small mass.
- They reside in the electron cloud surrounding the nucleus.
- Their rapid movement around the nucleus defines the size of the atom.

MODULE 3. ELEMENTS

- The information regarding the behavior and the arrangements of electrons outside of the nucleus is defined from quantum mechanics.
- Electrons have spin.
- Energy in an electron is defined and restricted, so it cannot just go 'anywhere.'
- An energy on electrons is quantized, meaning that it can only have certain values.

Electron Shells

Definition 3.2.10. An **electron shell** is a defined region of space about a nucleus that only contains electrons with approximately the same energy. The **shell number**, $\langle n \rangle$, is used to identify the electron shell. Electron shells are numbered 1 to 7, and electrons in higher shell numbers have more energy.

- Not all electron shells have an equal number of electrons.
- The number of electrons in a shell generally follows the formula $N = 2n^2$, where $\langle n \rangle$ is the electron shell level. (I.e., level 3 would have $2(3)^2$ or 18 electrons.)
- Lower level shells fill before higher ones.

Electron Subshells

Within an electron shell, there are subshells and orbitals. A subshell is a defined region of space within an electron shell that contains electrons of the same energy level, such as $\langle s \rangle$, $\langle p \rangle$, $\langle d \rangle$, or $\langle f \rangle$.

- The number of subshells in an electron shell is equal to the shell number.
- The number of electrons in each subshell is defined and independent of the shell number.
 - $\langle s \rangle$ has 2 electrons.
 - $\langle p \rangle$ has 6 electrons.
 - $\langle d \rangle$ has 10 electrons.
 - $\langle f \rangle$ has 14 electrons.

MODULE 3. ELEMENTS

- Electron subshells are denoted by the electron shell number followed by the letter of the electron subshell with a superscript denoting the number of the number of electrons in the given subshell, such as $1s^2$.

Electron Orbitals

Definition 3.2.11. Electron orbitals are regions of space within an electron subshell where with a specific energy is most likely to be found.

Note: The number of orbitals in each subshell is one-half of the number of electrons within the subshell, as each orbital can hold two electrons.

- $\langle s \rangle$ has 1 orbital.
- $\langle p \rangle$ has 3 orbitals.
- $\langle d \rangle$ has 10 orbitals.
- $\langle f \rangle$ has 14 orbitals.

The Aufbau Principle

Electron shells fill according to the *Aufbau Principle*, wherein electrons normally occupy electron subshells in an atom in order of increasing subshell energy, and the energy of subshells can overlap.

3.2.5 Electron Configuration

Definition 3.2.12. Electron configuration serves as a shorthand notation describing the subshells in an atom that are occupied by electrons.

Electron configuration is written by starting at the beginning (no electrons) and then continuously filling electron subshells until the correct number of electrons are represented. (Example table below)

Element	Number of Protons (Z)	Electron Configuration
Hydrogen	1	$1s^1$
Helium	2	$1s^2$
Carbon	6	$1s^2 2s^2 2p^2$
Neon	10	$1s^2 2s^2 2p^6$
Sodium	11	$1s^2 2s^2 2p^6 3s^1$
Potassium	19	$1s^2 2s^2 2p^6 3s^2 3p^6 4s^1$
Rubidium	37	$1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^{10} 4p^6 5s^1$
Cesium	55	$1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^{10} 4p^6 5s^2 4d^{10} 5p^6 6s^1$

MODULE 3. ELEMENTS




Figure 3.3: Ground State Electron Configurations of the Elements

Condensed Electron Configuration

Electron configuration can also be condensed by utilizing a starting point element (commonly a noble gas) surrounded by brackets then continuing the electron configuration from there.

For example, as Sodium's electron configuration is $1s^2 2s^2 2p^6 3s^1$ and Neon's electron configuration is $1s^2 2s^2 2p^6$, one could condense the electron configuration for Sodium down to $[\text{Ne}]3s^1$.

Note: For (d), use one less than the period, and for (f), use two less than the period.

3.2.6 Paramagnetic and Diamagnetic Substances

- Atoms where all electron orbitals are occupied by pairs of electrons are called *diamagnetic* atoms.

MODULE 3. ELEMENTS

- Atoms where all electron orbits are not occupied by pairs of electrons are called *paramagnetic* atoms.

Visualizing Electron Orbitals

We can visualize electron orbitals using arrows. Levels that are completely filled are diamagnetic and those that are not are paramagnetic.




Figure 3.4: Electron orbitals visualized with arrows. In this diagram, Scandium (Sc) is paramagnetic and Zinc (Zn) is diamagnetic.

Module 4

Compounds

4.1 Lecture 1: Ionic and Covalent Compounds

References chapter 5 of the textbook.

4.1.1 Types of Chemical Bonds

There are two main types of chemical compounds: ionic compounds and molecular compounds.

Note: The attractive forces that hold atoms together are called chemical bonds.

Definition 4.1.1. An ionic bond is a chemical bond formed through the transfer of electrons from one atom (or group of atoms) to another to form ions. A covalent bond is a chemical bond formed through the sharing of electron **pairs** between atoms.

MODULE 4. COMPOUNDS

Property	Ionic Bond	Covalent Bond
Chemical Formula	Formula Unit	Molecular Formula
Molecule Discreteness	No Discrete Molecule	Forms Discrete Molecules
Melting Point	High Melting Point	Lower Melting Points
Electrical Conductivity	Good Conductors when Molten	Cannot Conduct
Solubility	Soluble in Polar Solvents and Not Soluble in Non-Polar Solvents	Soluble in Non-Polar Solvents and Some Polar Solvents
State of Matter	N/A	Gases, Liquids, and Low-Melting Solids

Note: Chemical bonds can be strongly ionic or strongly covalent, but they will usually be a mix of both.

Note: Only valence electrons are able to be utilized for bonding.

The Octet Rule

Electron bonds occur to further stabilize a chemical or group of chemicals by reaching a state of completely full valence electrons. Thus, Noble Gases are entirely unreactive, as they already have completely full valence electrons. Through these bonds, electrons can either **lose, gain, or share** electrons to reach a full valence state (which would be the same configuration as any one of the noble gases). This process is often referred to as **the octet rule**.

Lewis Dot Symbols

Definition 4.1.2. A Lewis Dot Symbol is a chemical symbol of an element surrounded by dots equal to the number of valence electrons.

Note: The choice of where the dots are placed is not important.

4.1.2 Ionic Bonds

MODULE 4. COMPOUNDS

Definition 4.1.3. Ions are atoms (or groups of atoms) that are electrically charged as the result of losing or gaining valence electrons. Anions are negatively charged (usually nonmetals), and cations are positively charged (usually metals).

Whether or not an element loses or gains electrons is dependent on how many valence electrons the element has. If the number of valence electrons is closer to 0, it is more likely to lose electrons, and if the number of valence electrons is closer to 8, it is more likely to gain electrons.

This property is why metals, which are generally closer to the left of the periodic table and thus have less valence electrons, are more likely to lose electrons, and non-metals, which are further to the right and thus have more valence electrons, and more likely to gain electrons.

Note: By this rule, common ionic behavior is best observed in the representative elements (but not in the transition elements, as they have a more balanced number of valence electrons). Group 1 will normally have a 1+ charge, group 2 will normally have a 2+ charge, group 13 will have a 3+ charge, group 14 will have a 4+ or 4- charge, group 15 will have a 3- charge, group 16 will have a 2- charge, and group 17 will have a 1- charge.

Ions with the same electron configuration as a noble gas are considered **Isoelectronic** with the noble gas, and **isoelectronic species** are the set of ions (and possibly an element as well) that all have the same electron configuration.

Ionic Compound Formulas

The net charge of every compound must be equal to zero, with positive ions written first and negative ions written after. Charges on each individual ion are not reflected in the formula, and subscript numbers in the formula give the combining ratio of the ions rather than the exact count.

For example, in the ion NaCl, Na carries a +1 charge and Cl carries a -1 charge. Thus, NaCl is neutral.

MODULE 4. COMPOUNDS

Ionic Compounds

Ionic compounds are usually comprised of a metal and a nonmetal (also known as a **polyatomic ion**), and the ratio in which these ions combine is always the lowest possible ratio that garners a net neutral charge between the two. These ratios are known as **Formula Units**.

Note: Metals are groups IA (1), IIA (2), and IIIA (13), and Non-Metals are groups VA (15), VIA (16), and VIIA (17).

Metallic atoms usually lose electrons and become positively charged, and non-metallic atoms usually gain electrons and become negatively charged.

4.1.3 Covalent Bonds

Definition 4.1.4. Covalent bonds are bonds between two non-metal atoms as a result of two atoms **sharing** the same valence electrons.




Figure 4.1: Covalent Bond

Atoms in covalent bonds can have both bonding electrons and non-bonding

MODULE 4. COMPOUNDS

electrons, where the shared electrons are considered 'bonding.'

Note: Covalent compounds can form single, double, or triple bonds by sharing one, two, or three pairs of electrons, with each bond represented with a straight line.

4.1.4 Molecular Geometry and VSEPR

Definition 4.1.5. Molecular Geometry is a description of the three dimensional arrangement of atoms within a molecule, and it plays a critical role in determining the physical and chemical properties of substances.

Definition 4.1.6. Valence Shell Electron Pair Repulsion Theory, or VSEPR Theory, is a set of procedures for predicting the geometric structure of a molecule from its lewis dot structure.

In VSEPR, the non-binding electron pairs and the number of atoms bonded to an atom determine the geometric structure of an atom as **electrons naturally repel each other**. (**Note:** this is why remembering the number of electron groups is key.)

A VSEPR electron group is a group of valence electrons around an atom or molecule. (**Note:** It is slightly different from an electron pair.)

The Core of Molecular Geometry based on VSEPR

1. Atoms arrange themselves to be as far from other electrons as possible.
2. There is no distinction made between bonding/shared electrons and non-bonding/lone electrons.
3. Single, double, and triple bonds are all counted equally as one pair attached to the central atom.

MODULE 4. COMPOUNDS

Types of VSEPR Configurations





















Number of Electron Dense Areas	Electron-Pair Geometry	Molecular Geometry				
		No Lone Pairs	1 lone Pair	2 lone Pairs	3 lone Pairs	4 lone Pairs
2 	Linear	 Linear				
3 	Trigonal planar	 Trigonal planar	 Bent			
4 	Tetrahedral	 Tetrahedral	 Trigonal pyramidal	 Bent		
5 	Trigonal bipyramidal	 Trigonal bipyramidal	 Sawhorse	 T-shaped	 Linear	
6 	Octahedral	 Octahedral	 Square pyramidal	 Square planar	 T-shaped	 Linear

Figure 4.2: Types of VSEPR Configurations

- Molecules with 2 or 3 VSEPR pairs are **planar**.
- Molecules with VSEPR pairs that are strictly 180° apart are **linear**.
- Molecules with VSEPR pairs that are strictly 120° will be **Triagonal Planar** or **Angular/Bent**.
- Molecules with VSEPR pairs that are strictly 109.5° will be either **Tetrahedral**, **Trigonal Pyramidal**, or **Angular/Bent**

MODULE 4. COMPOUNDS

Formula	VSEPR Groups	Shape
XB_2	2	Linear
XB_3	3	Trigonal Planar
XB_2N_1	3	Bent
XB_4	4	Tetrahedral
XB_3N_1	4	Trigonal Pyramidal
XB_2N_2	4	Bent

Table 4.1: X is the Central Atom, B is the Bonding Electron Groups, and N is the Non-Bonding Electron Groups.

4.1.5 Bond Polarity

The greater the electronegativity of an atom, the greater its electron attraction. By this rule, atoms with different electronegativities will have bonds of different strengths.

Definition 4.1.7. Bond Polarity is the measure of the magnitude of difference in electron bonds between atoms.

- Bonds with equal sharing of electrons are called non-polar covalent bonds.
- Bonds that do not have an equal sharing of electrons are known as polar covalent bonds and are represented with a "D-" or "D+" for negative and positive respectively.
- Polar covalent bonds create a polarity in the resulting molecule, which influences its resulting properties.

Calculating Bond Polarity

Electronegativity differences can be used as a rough measure of calculating bond polarity given the following rules:

MODULE 4. COMPOUNDS

Electronegativity Range	Bond Polarity Type
$\Delta\chi < 0.4$	Non-Polar Covalent
$0.4 \leq \Delta\chi < 2$	Polar Covalent
$2 \geq \Delta\chi$	Ionic

Table 4.2: $\Delta\chi$ is the magnitude of difference between the electronegativities between the two elements.

For example, Hydrogen has an electronegativity of 2.1 and Chlorine has an electronegativity of 3.0, causing a difference of $\Delta\chi = 0.9$, and thus, a polar covalent bond. *The greater this difference, the higher the polarity.*

Note: Although the bonds can be polarized, sometimes the polarity can cancel out, causing the molecule to be non-polar.

4.1.6 Molecular Polarity

Definition 4.1.8. Molecular Polarity is a measure of degree inequality in the attraction of bonding electrons to various locations within a molecule, where polar molecules have unsymmetrical charge distribution and non-polar molecules have symmetrical charge distribution.

Note: Molecular geometry and bond polarity combined determine molecular polarity.

4.2 Lecture 2: Naming of Compounds

The International Union of Pure and Applied Chemistry, or IUPAC, sets the following standard rules for naming compounds in chemistry.

- **Binary Ionic Compounds** contain a metal and a non-metal.
- **Binary Molecular Compounds** consist of only non-metals. (**Note:** For naming purposes, metalloids are treated as non-metals.)
- **Binary Acids** are compounds that release hydrogen ions when dissolved in water.

MODULE 4. COMPOUNDS

4.2.1 Ionic Compounds

- Ionic Compounds are the attraction between oppositely-charged ions, not atoms.
- Ionic compounds form crystal lattices, not molecules.
- The charge on an ionic compounds is always neutral.
- Ionic compounds may contain one or more *polyatomic ions*.

Metal Ions

For naming purposes, there are two types of metal ions: fixed and variable charge.

Definition 4.2.1. **Fixed charge** ions are metal ions that form one type of positive ion that always have the same charge. **Variable charge** ions are metal ions that can form more than one type of positive ion.

The only fixed-charge ions are the metal elements in the IA, IIA, IB, IIB, and IIIA groups, or Lithium, Sodium, Potassium, Rubidium, Cesium, Beryllium, Magnesium, Calcium, Strontium, Barium, Aluminum, Gallium, Silver, Zinc, and Cadmium. *All other metals have variable charge*, such as Chromium, Cobalt, or Gold.

4.2.2 Naming Binary Ionic Compounds

- When naming ionic compounds, positively-charged ions (cations) are named first, and negatively-charged ions (anions) are named after.
- Metal ions (fixed or variable) take the name of the metal they come from.
- Non-metal ions are named by taking the stem of the non-metal and adding the suffix "-ide."

Example 4.2.1. Li^+ is a Lithium Ion, Mg^{2+} is a Magnesium Ion, but F^- is a Fluoride ion and O^{2-} is an Oxide ion.

Note: Polyatomic ions always bring their own name to the compound.

MODULE 4. COMPOUNDS

Element	Stem	Name of Ion	Formula
Bromine	brom-	bromide ion	Br^-
Carbon	carb-	carbide ion	C^{4-}
Chlorine	chlor-	chloride ion	Cl^-
Fluorine	fluor-	fluoride ion	F^-
Hydrogen	hydr-	hydride ion	H^-
Iodine	iod-	iodide ion	I^-
Nitrogen	nitr-	nitride ion	N^{3-}
Oxygen	ox-	oxide ion	O^{2-}
Phosphorus	phosph-	phosphide ion	P^{3-}
Sulfur	sulf-	sulfide ion	S^{2-}

Table 4.3: Names for Common Non-Metal Ions

Example 4.2.2. CaF_2 would be Calcium fluoride, and no importance is placed on the amount for each element.

When naming binary ionic compounds with variable-charge metals, there has to be a differentiation between the different charges of the metal for situations such as differentiating between FeS and Fe_2S_3 given the ions Fe^{2+} and Fe^{3+} .

The solution to this is referring to each ion of the metal by its cation charge in roman numerals within parentheses, so Fe^{2+} becomes Iron (II) and Fe^{3+} becomes Iron (III).

Note: If there is a metal in the compound, the metal is ionic.

4.2.3 Naming Compounds with Polyatomic Ions

1. Most polyatomic ions are negatively charged, and the only two main positive ones are ammonium and hydronium.
2. Four polyatomic ions have names that end in "-ide" already: hydroxide, cyanide, azide, and peroxide.
3. Pay attention to the "-ate" and "-ite" pairs, as they differ by the number of oxygen atoms, where "-ate" always has a greater number of oxygen atoms. For example, Sulfate (SO_4^{2-}) and Sulfite (SO_3^{2-}).
4. Some pairs differ by the addition of a hydrogen ion, such as Carbonate (CO_3^{2-}) and Hydrogen Carbonate (HCO_3^{2-}).

MODULE 4. COMPOUNDS

5. When Sulfur replaces an Oxygen atom, the prefix "Thio-" is added to the name, such as with Cyanate (OCN) and Thiocyanate (SCN) or Sulfate (SO_4^{2-}) and Thiosulfate ($\text{S}_2\text{O}_3^{2-}$).

4.2.4 Binary Molecular Compounds

Binary molecular compounds consist of two non-metal elements, and these compounds are named as the formula is written.

- The least electronegative element is named first.
- The stem of the second non-metal follows with the suffix "-ide."
- Numerical prefixes are used to indicate the number of both atoms present.

Note: The number of atoms present is not clarified for polyatomic ions, but it is for binary molecular compounds.

Prefix	Value
mono	1
di	2
tri	3
tetra	4
penta	5
hexa	6
hepta	7
octa	8
nona	9
deca	10

Note: The exception to this rule is that binary compounds with hydrogen listed as the first element are named without using the prefix for how many atoms there are, and the prefix "mono" is always dropped at the beginning of a name.

Common Names to Know

Note: The above examples are all binary compounds, but other molecular compounds with more than 2 elements don't follow these rules.

MODULE 4. COMPOUNDS

Formula Unit	Name
H ₂ O	Water
H ₂ O ₂	Hydrogen Peroxide
NH ₃	Ammonia
N ₂ H ₄	Hydrazine
CH ₄	Methane
C ₂ H ₆	Ethane
PH ₃	Phosphine
AsH ₃	Arsine

4.2.5 Introduction to Acids

Definition 4.2.2. An acid is a hydrogen-containing molecular compound whose molecules yield Hydrogen ions (H⁺) when dissolved in water (H₂O).

These acids can be identified by having Hydrogen as the first element shown in its formula, such as HCl, H₂S, H₂SO₄, or HNO₃ (which are all acids), whereas NH₃, CH₄, PH₃, or SiH₄ are all non-acidic.

Note: Acids produce H⁺ in water, but an anion is also produced, depending on the structure of the molecular compound. For example, HCl produces H⁺ and Cl⁻.

Acid Nomenclature

Anion Ending	Acid Name
"-ide"	hydro-(<i>stem</i>)-ic acid
"-ate"	(<i>stem</i>)-ic acid
"-ite"	(<i>stem</i>)-ous acid

4.2.6 Common Student Mistakes

- For **ionic compounds ONLY**, formulas **must be reduced to its simplest ratios**. For example, Mn₂O₄ must be reduced to MnO₂.
- Do not confuse Ammonia (NH₃) and an Ammonium Ion (NH₄⁺).
- Chemical formulas do not show ionic charges. For example, write table salt as NaCl, not Na⁺Cl⁻.

MODULE 4. COMPOUNDS

- Don't put parentheses unless they are needed to show more than one of the same ion. For example, write NaNO_3 , not $\text{Na}(\text{NO}_3)$

Module 5

Reactions

5.1 Lecture 1: Reactions

5.1.1 Introduction to Reactions

References Chapter 7, Sections 7.6 and 7.11 of the textbook.

Definition 5.1.1. A chemical reaction is the process in which at least one or more new substances are produced as a result of a chemical change.

Evidence of chemical changes include a change in color, temperature, or odor, or the formation of a gas or insoluble solid (or precipitate).

Definition 5.1.2. The input/starting substances of a reaction are known as *reactants*, and the output/ending substances of a reaction are known as *products*.

The Law of Conservation of Mass


Definition 5.1.3. Mass is neither created nor destroyed (in any ordinary chemical reaction). Therefore, the total mass of reactants is always equal to the total mass of products.

5.1.2 Chemical Equations

MODULE 5. REACTIONS

Definition 5.1.4. A chemical equation is a representation for a chemical reaction that uses chemical symbols and chemical formulas to describe changes that occur in a chemical reaction.


In a chemical equation, the reactants are written on the left side (before the arrow), and products are written on the right side (after the arrow), with each substance separated by a plus sign on each side.



In order for a chemical equation to be valid, it must satisfy two conditions:

1. Only the reactants and products involved in the reaction are represented in the equation.
2. Accurate or correct formulas must be used, not simply empirical formulas. For example, diatomic molecules must always be presented as diatomic rather than as single atoms.
3. The same number of atoms of each kind must be present on each side of the equation.

In a chemical formula, the physical state of a substance may be indicated using symbols in parentheses.



Note: *aq* means aqueous.

Balancing Equations

To balance an equation, adjust the number of atoms of each element so that they are the same on each side of the equation.

Note: When balancing an equation, only change the coefficients on each side. **NEVER** alter a substance's formula (i.e. subscripts) to balance the equation, as it creates a completely new substance.

MODULE 5. REACTIONS

Note: Remember that the coefficients in a balance equation are the lowest whole numbers that balance the equation. As such, it's incredibly useful to consider polyatomic ions as single entities, provided that they maintain their identities during the reaction.

When balancing equations, follow the order:


1. Balance the metals
2. Balance the polyatomic ions if they stay together
3. Balance the non-metals
4. Balance hydrogen
5. Balance oxygen

Note: Additionally, it also helps to create a table when balancing an equation.

Example 5.1.3. $\text{C}_4\text{H}_{10(g)} + \text{O}_{2(g)} \longrightarrow \text{CO}_{2(g)} + \text{H}_2\text{O}_{(l)}$

Elements	Reactant Count	Product Count
Carbon	4	1
Hydrogen	10	2
Oxygen	2	3

Naturally, this process can be followed to reach a balanced state.



Elements	Reactant Count	Product Count
Carbon	8	8
Hydrogen	20	20
Oxygen	26	26

5.1.3 Types of Chemical Reactions

Type of Reaction	General Equation
Combination (Synthesis)	$A + B \longrightarrow AB$
Decomposition	$AB \longrightarrow A + B$
Single Replacement	$A + BC \longrightarrow AC + B$
Double Replacement	$AB + CD \longrightarrow AD + BC$
Combustion	$C_x + H_y + O_2 \longrightarrow CO_2 + H_2O$

- Synthesis: Two or more reactants combining to form only one product.
- Decomposition: One reactant breaks down to produce two or more different substances.
- Single Replacement: One element reacts with a compound to replace one of the elements of that compound.
- Double Replacement: Two elements react with a compound to replace two of the elements of that compound.
- Combustion: Substance reacts with oxygen and produces heat and/or a flame.

Note: A double replacement reaction will usually result in one or more of these three: the formation of a precipitate, the formation of a gas/bubbles, the release or absorption of heat, or the formation of a molecular compound (such as H_2O).

Module 6

Moles

6.1 Lecture 1: Counting Atoms in Moles

6.1.1 The Law of Definite Proportions/Constant Composition

Definition 6.1.1. The law of definite proportions (also known as the Law of Constant Composition) states that in a pure compound, the elements described by its formula are present in the same definite proportion by mass, and thus, it will always retain the same physical and chemical properties for all objects of this substance type.

For example, water always contains only two elements, Hydrogen and Oxygen, and it will always be about 11.2% Hydrogen and 88.8% Oxygen by mass. In the event that a compound differs from these qualities whatsoever, it is no longer water.

6.1.2 Percent Composition

Additionally, this knowledge can be used to calculate percent composition of substances from its molecular formulas.

Percent Composition from Known Formulas

MODULE 6. MOLES

Example 6.1.1. For example, given the formula MgCl_2 and the atomic masses of Mg and Cl at 24.30 amu and 35.45 amu respectively, one can compute the total molecular mass of MgCl_2 as $1 \times 24.30 + 2 \times 35.45 = 95.20 \text{ amu}$, and from there, the percent composition of each element as $\frac{24.30}{95.20} \times 100 = 25.53\%$ for Mg and $\frac{2 \times 35.45}{95.20} \times 100 = 74.47\%$ for Cl.

Percent Composition from Experimental Data

Additionally, when calculating percent compositions from unknown compositions with experimental data, simply manually calculate the total mass from the data then divide by the mass of each element.

6.1.3 Counting Atoms

Definition 6.1.2. Large numbers, such as the number of atoms in a substance, are counted in units known as moles, where **one mole is equal to 6.022×10^{23}** , a value known as **Avogadro's number**. Moles are abbreviated as *mol*.

The mass of each mole of an element (or every 6.022×10^{23} atoms of the element combined) is known as the **molar mass**, and it is different for each element. For all elements, the molar mass is equal to the atomic mass when the element is present in an atomic form. As such, atomic mass is the same as atomic molar mass.

Molar mass of a compound is measured in grams and is numerically equal to the formula or molecular mass of the compound (which is measured in *amu*). For example, the formula/molecular mass of water is 18.02 amu and the molar mass is 18.02 grams.

6.1.4 Empirical Formula vs. Molecular Formula

MODULE 6. MOLES

Definition 6.1.3. The empirical formula of a compound gives the simplest whole number ratio of the atoms present for each formula unit in a compound, whereas the molecular formula represents the total number of atoms of each element present in one molecule of a compound (which is **not simplified**).

Note: The molecular formula of a substance is the true formula of the compound, and it is calculated by using the empirical formula and the molar mass together.

Example 6.1.2. The empirical formula for Ethylene is CH_2 , but given its molecular mass, the actual molecular formula is C_2H_4 .

Calculating Empirical Formulas

1. Assume some definite starting quantity (like 100.0g) of the compound if it isn't given and then calculate the mass of each element component in grams.
2. Following this conversion formula (where m_s is the empirical mass of the element within the substance, m_a is the atomic molecular mass of the element, and x is the number of moles of atoms of this element within the substance,

$$x = \frac{m_s}{m_a} \quad (6.1.1)$$

3. Divide the moles of atoms of each element by the moles of atoms of the element that had the smallest value. (If they are whole numbers, use them as subscripts and write the empirical formula).
4. Otherwise, multiply the values obtained previously by the smallest numbers that will convert them to whole numbers.

MODULE 6. MOLES

Example 6.1.3. *An analysis of salt shows that it contains 56.58% Potassium (K), 8.68% Carbon (C), and 34.73% Oxygen (O). Calculate the empirical formula for this substance.*

Step 1: Express each element in grams, assuming we have 100 grams of the compound.

$$K = 56.58\text{g}$$

$$C = 8.68\text{g}$$

$$O = 34.73\text{g}$$

Step 2: Convert the grams of each element into moles.

$$56.58\text{gK} \left(\frac{1\text{molK}}{39.10\text{gK}} \right) = 1.447\text{molK}$$

$$8.68\text{gC} \left(\frac{1\text{molC}}{12.01\text{gC}} \right) = 0.723\text{molC}$$

$$34.73\text{gO} \left(\frac{1\text{molO}}{16.00\text{gO}} \right) = 2.171\text{molO}$$

Step 3: Divide each number of moles by the smallest molar value calculated previously (0.723 mol).

$$K = \frac{1.447\text{mol}}{0.723\text{mol}} = 2.00$$

$$C = \frac{0.723\text{mol}}{0.723\text{mol}} = 1.00$$

$$O = \frac{2.171\text{mol}}{0.723\text{mol}} = 3.00$$

Therefore, the simplest ratio of K:C:O is 2:1:3, resulting in an empirical formula of K_2CO_3 .

Calculating Molecular Formulas

The molecular formula can be calculated from the empirical formula if the molar mass is known, given that the mass of the molecular formula will be

MODULE 6. MOLES

equal to some multiple of the empirical mass.

The simplest way to calculate the molecular formula is to divide the molar mass of the substance by the empirical mass to determine the scaling value, then multiply all of the coefficients of the formula by this number.

$$n = \frac{\text{Molar Mass}}{\text{Empirical Mass}} \quad (6.1.2)$$

Note: Where n is the number of empirical formula units.

Example 6.1.4. *What is the molecular formula of a compound which has an empirical formula of CH_2 and a molar mass of 126.2 grams?*

As we know the empirical mass of CH_2 to be 14.03 grams per formula unit, we can calculate

$$n = \frac{126.2\text{g}}{14.03\text{g}} = 9$$

therefore, the molecular formula is scaled from CH_2 to $(\text{CH}_2)_9$ or C_9H_{18} .

6.2 Lecture 2: Stoichiometry

Definition 6.2.1. Chemical Stoichiometry is the study of quantitative and molecular relationships between reactants and products within chemical reactions.


6.2.1 Calculating Theoretical Yield

Definition 6.2.2. The **theoretical yield** is the maximum amount of product you will get if the reaction occurs with complete efficiency.

The theoretical yield of a reaction is calculated using the coefficients of a chemical equation, which not only describe the number of atoms/molecules being reacted/produced but also describe the number of moles of atoms/-molecules being reacted/produced.

MODULE 6. MOLES

Example 6.2.1. *Bismuth (III) chloride will react with hydrogen sulfide to form Bismuth (III) sulfide and Hydrochloric acid. Write the balanced equation for this reaction then calculate how many moles of acid would be formed if 15.0 mol of Hydrogen sulfide react.*



$$15.0 \text{ mol H}_2\text{S} \times \frac{6 \text{ mol HCl}}{3 \text{ mol H}_2\text{S}} = 30.0 \text{ mol HCl}$$

6.2.2 Percent Yield

Definition 6.2.3. The **actual yield** is the actual amount of product obtained from a chemical reaction and the **percent yield** is the ratio of the actual yield to the theoretical yield (expressed as a percentage).

$$\text{Percent Yield} = \frac{\text{Actual Yield}}{\text{Theoretical Yield}} \times 100 \quad (6.2.1)$$

Module 7

Intermolecular Forces

Module 8

Solutions