



Estimating survival using telemetry or age distribution data



Estimating survival using telemetry ~~or age distribution data~~



Options

(1) Telemetry

- ▶ Binomial model
- ▶ Kaplan-Meier model

Options

(1) Telemetry

- ▶ Binomial model
- ▶ Kaplan-Meier model

(2) Age distribution data

- ▶ Often used but rarely valid

Options

(1) Telemetry

- ▶ Binomial model
- ▶ Kaplan-Meier model

(2) Age distribution data

- ▶ Often used but rarely valid

(3) Capture-mark-recapture

- ▶ Covered in the previous lecture

Pros

- Fate is often known
- Analysis can be straight-forward
- Much additional information can be obtained



Pros

- Fate is often known
- Analysis can be straight-forward
- Much additional information can be obtained

Cons

- Batteries don't last as long as bands
- When fate is unknown, analysis can be hard
- Transmitters may influence vital rates and behavior



Design

- n animals are randomly sampled

Design

- n animals are randomly sampled
- The fates of all n are known at end of the study period

Design

- n animals are randomly sampled
- The fates of all n are known at end of the study period
- x of the n animals survive

Design

- n animals are randomly sampled
- The fates of all n are known at end of the study period
- x of the n animals survive
- The estimate of survival over the time interval is:

$$\hat{S} = \frac{x}{n}$$

Assumptions

- (1) Fates are independent
- (2) Fates are known
- (3) All individuals are exposed to mortality during the same time interval

Assumptions

- (1) Fates are independent
- (2) Fates are known
- (3) All individuals are exposed to mortality during the same time interval

When **censoring** occurs, assumptions (2) and (3) are violated

Assumptions

- (1) Fates are independent
- (2) Fates are known
- (3) All individuals are exposed to mortality during the same time interval

When **censoring** occurs, assumptions (2) and (3) are violated

Censoring occurs when the time of mortality is not directly observed

Right censoring

- Occurs when animals leave the sample before they die.
- Examples
 - ▶ Battery failure
 - ▶ Emigration

Right censoring

- Occurs when animals leave the sample before they die.
- Examples
 - ▶ Battery failure
 - ▶ Emigration

Left censoring and interval censoring are rarely a concern


Right censoring

- Occurs when animals leave the sample before they die.
- Examples
 - ▶ Battery failure
 - ▶ Emigration

Left censoring and interval censoring are rarely a concern

If censoring occurs, the Kaplan-Meier estimator is more appropriate than the binomial model.

CENSORING




Diamonds indicate mortality. Circles indicate censoring.
Black lines show the observation period.

$$\hat{S}_t = \prod_{j:t_i \leq t} \left(\frac{r_j - d_j}{r_j} \right)$$

S_t – probability of surviving to time t

r_j – number of individuals “at risk” prior to time t

d_j – number of mortalities prior to time t



<https://youtu.be/Mgp46izlfeo>

Critical Assumption:

Censoring must be independent of survival

- It's fine if transmitters fail randomly
- It's a problem if they fail because of mortality

Critical Assumption:

Censoring must be independent of survival

- It's fine if transmitters fail randomly
- It's a problem if they fail because of mortality


Consequence of violating this assumption:

If an animal dies, but you mistakenly right censor it because you think its battery died, survival will be over-estimated.

AGE DISTRIBUTION DATA

Standard practice


- We have a pile of wings that can be aged
- We assume the age ratios tell us something about survival



AGE DISTRIBUTION DATA

Standard practice

- We have a pile of wings that can be aged
- We assume the age ratios tell us something about survival



There are many problems with this

Suppose a cohort of individuals is monitored over time, so that we can obtain information about survival.

$$n_{i+1,t+1} = n_{i,t}S_{i,t}$$

Suppose a cohort of individuals is monitored over time, so that we can obtain information about survival.

$$n_{i+1,t+1} = n_{i,t} S_{i,t}$$

$$S_{i,t} = \frac{n_{i+1,t+1}}{n_{i,t}}$$

Suppose a cohort of individuals is monitored over time, so that we can obtain information about survival.

$$n_{i+1,t+1} = n_{i,t}S_{i,t}$$

$$S_{i,t} = \frac{n_{i+1,t+1}}{n_{i,t}}$$

All is well. This is a valid approach based on the binomial model.

The problem is that this...

$$S_{i,t} = \frac{n_{i+1,t+1}}{n_{i,t}}$$

The problem is that this...

$$S_{i,t} = \frac{n_{i+1,t+1}}{n_{i,t}}$$

...is not the same as data from “standing age distribution”

$$S_{i,t} = \frac{n_{i+1,t}}{n_{i,t}}$$

Standing age distribution data can't be used to estimate survival, unless. . .

- Population is at the stable age distribution
- $\lambda = 1$

Population reconstruction refers to the calculation of the abundance and age distribution of a cohort at some initial time, usually based on “ages at death”.

Population reconstruction refers to the calculation of the abundance and age distribution of a cohort at some initial time, usually based on “ages at death”.

This is only possible if we know the ages of death for **all** individuals in the cohort.

Population reconstruction refers to the calculation of the abundance and age distribution of a cohort at some initial time, usually based on “ages at death”.

This is only possible if we know the ages of death for **all** individuals in the cohort.

Shouldn't use this type of “reconstruction data” as real data in a subsequent analysis.

The age data are biased

- Age data not representative of actual age distribution
- Hunting, trapping, and capture-recapture methods do not evenly sample the population

When estimating survival, it is always good to...

- Collect more than 1 year of data
- Follow multiple cohorts of individuals over long periods of time
- Estimate survival using telemetry or mark-recapture methods