


Occupancy Estimation and Modeling



Occupancy models were developed to estimate metapopulation parameters when **detection probability** (p) is < 1 .

Occupancy models were developed to estimate metapopulation parameters when **detection probability** (p) is < 1 .

Parameters of interest:

- ψ – Probability that a site is occupied

Occupancy models were developed to estimate metapopulation parameters when **detection probability** (p) is < 1 .

Parameters of interest:

- ψ – Probability that a site is occupied
- γ – Probability that an unoccupied site becomes colonized

Occupancy models were developed to estimate metapopulation parameters when **detection probability** (p) is < 1 .

Parameters of interest:

- ψ – Probability that a site is occupied
- γ – Probability that an unoccupied site becomes colonized
- ε – Probability that an occupied site goes extinct

Occupancy models were developed to estimate metapopulation parameters when **detection probability** (p) is < 1 .

Parameters of interest:

- ψ – Probability that a site is occupied
- γ – Probability that an unoccupied site becomes colonized
- ε – Probability that an occupied site goes extinct
- p – Probability of detecting at least one individual at a site that is occupied (on a single sampling occasion)

If $p < 1$, we might incorrectly conclude that a site is unoccupied if we don't detect any individuals

If $p < 1$, we might incorrectly conclude that a site is unoccupied if we don't detect any individuals

Consequences

- We will underestimate the state variable: The proportion of sites occupied

If $p < 1$, we might incorrectly conclude that a site is unoccupied if we don't detect any individuals

Consequences

- We will underestimate the state variable: The proportion of sites occupied
- We might make incorrect conclusions about habitat relationships

Scenario

- No interest in colonization or extinction events

Scenario

- No interest in colonization or extinction events
- Instead, we directly estimate ψ during one season

Scenario

- No interest in colonization or extinction events
- Instead, we directly estimate ψ during one season
- Useful for assessing snapshot of habitat relationships or for modeling a species' distribution


Scenario

- No interest in colonization or extinction events
- Instead, we directly estimate ψ during one season
- Useful for assessing snapshot of habitat relationships or for modeling a species' distribution
- We assume **population closure**: A site's occupancy state does not change during the survey period (season)

Scenario

- No interest in colonization or extinction events
- Instead, we directly estimate ψ during one season
- Useful for assessing snapshot of habitat relationships or for modeling a species' distribution
- We assume **population closure**: A site's occupancy state does not change during the survey period (season)
- Definitions of site and season are very important

HOW MANY POTHOLES ARE OCCUPIED BY MALLARDS?



Model for the occurrence state

$$O_i \sim \text{Bernoulli}(\psi)$$

Model for the occurrence state

$$O_i \sim \text{Bernoulli}(\psi)$$

Model for the data

$$y_{ij} \sim \text{Bernoulli}(O_i \times p)$$

Model for the occurrence state

$$O_i \sim \text{Bernoulli}(\psi)$$

Model for the data

$$y_{ij} \sim \text{Bernoulli}(O_i \times p)$$

We need to estimate ψ (occurrence probability) and p (detection probability)

To estimate ψ and p , we should use a good study design:

- We need multiple sites

To estimate ψ and p , we should use a good study design:

- We need multiple sites
- Sites should be randomly selected

To estimate ψ and p , we should use a good study design:

- We need multiple sites
- Sites should be randomly selected
- We must have multiple sampling occasions at (a subset of) the sites

10 sites and 3 sampling occasions

	Occasion 1	Occasion 2	Occasion 3
Site 1	0	0	1
Site 2	0	0	0
Site 3	0	0	0
Site 4	1	0	0
Site 5	0	0	0
Site 6	1	0	1
Site 7	1	0	0
Site 8	0	0	0
Site 9	0	1	0
Site 10	0	0	0

10 sites and 3 sampling occasions

	Occasion 1	Occasion 2	Occasion 3
Site 1	1	1	1
Site 2	0	0	0
Site 3	0	0	0
Site 4	1	1	1
Site 5	0	0	0
Site 6	1	0	1
Site 7	1	1	1
Site 8	0	0	0
Site 9	0	1	1
Site 10	0	0	0

HOW DO WE ESTIMATE ψ AND p ?

HOW DO WE ESTIMATE ψ AND p ?


Maximum likelihood

- This method tries to find the most “likely” values of ψ and p , given the data
- It can also be used to estimate standard errors (SEs) and confidence intervals (CIs)

HOW DO WE ESTIMATE ψ AND p ?

Maximum likelihood

- This method tries to find the most “likely” values of ψ and p , given the data
- It can also be used to estimate standard errors (SEs) and confidence intervals (CIs)




HOW DO WE ESTIMATE ψ AND p ?

Maximum likelihood

- This method tries to find the most “likely” values of ψ and p , given the data
- It can also be used to estimate standard errors (SEs) and confidence intervals (CIs)

Software

Program PRESENCE, Program MARK, R package unmarked



10 sites and 3 sampling occasions

	Occasion 1	Occasion 2	Occasion 3
Site 1	0	0	1
Site 2	0	0	0
Site 3	0	0	0
Site 4	1	1	0
Site 5	0	0	0
Site 6	1	1	1
Site 7	1	0	0
Site 8	0	0	0
Site 9	0	1	0
Site 10	0	0	0

Naive occupancy = $5/10 = 0.50$

Estimated occupancy = $\hat{\psi} = 0.61 \pm 0.22$

Estimated detection = $\hat{p} = 0.44 \pm 0.16$

If you sample a site K times, the overall detection probability (\bar{p}) is:

$$\bar{p} = 1 - (1 - p)^K$$

\bar{p} is detection probability after K sampling occasions

p is detection probability on a single occasion

K is the number of sampling occasions (e.g., visits to a site)

- (1) Population closure: Occurrence state does not change during sampling

- (1) Population closure: Occurrence state does not change during sampling
- (2) Occurrence probability is the same for all sites, unless we account for **covariates**

- (1) Population closure: Occurrence state does not change during sampling
- (2) Occurrence probability is the same for all sites, unless we account for **covariates**
- (3) Detection probability is the same for all sites and sampling occasions, unless we account for covariates

- (1) Population closure: Occurrence state does not change during sampling
- (2) Occurrence probability is the same for all sites, unless we account for **covariates**
- (3) Detection probability is the same for all sites and sampling occasions, unless we account for covariates
- (4) Statistical independence

The value of ψ or p may depend on other variables, e.g. habitat, weather, observer abilities

The value of ψ or p may depend on other variables, e.g. habitat, weather, observer abilities



How do we accommodate covariates?

The value of ψ or p may depend on other variables, e.g. habitat, weather, observer abilities

How do we accommodate covariates?

The key is to think of detection probability as a function. A common choice is the logit-linear model:

$$\text{logit}(\psi_i) = \beta_0 + \beta_1 \text{ELEV}_i$$



With more than 1 season, we can estimate all the parameters of our metapopulation model, plus detection probability p

With more than 1 season, we can estimate all the parameters of our metapopulation model, plus detection probability p

$$\psi_{i,t+1} = O_{i,t}(1 - \varepsilon) + (1 - O_{i,t})\gamma$$

$$O_{i,t+1} \sim \text{Bernoulli}(\psi_{i,t+1})$$

$$y_{i,j,t} \sim \text{Bernoulli}(O_{i,t} \times p)$$

5 sites, 2 seasons, and 3 sampling occasions

	Season 1			Season 2		
	Occasion 1	Occasion 2	Occasion 3	Occasion 1	Occasion 2	Occasion 3
Site 1	0	0	1	1	0	0
Site 2	0	0	0	0	0	0
Site 3	0	0	0	1	0	0
Site 4	1	1	1	0	0	0
Site 5	0	0	0	1	1	0

Occupancy models let us estimate metapopulation parameters when detection is imperfect.

Occupancy models let us estimate metapopulation parameters when detection is imperfect.

There are no individual-level data or parameters.

Occupancy models let us estimate metapopulation parameters when detection is imperfect.

There are no individual-level data or parameters.

These methods are often easy to implement over large areas and so are used in monitoring programs.

Occupancy models let us estimate metapopulation parameters when detection is imperfect.

There are no individual-level data or parameters.

These methods are often easy to implement over large areas and so are used in monitoring programs.

Definition of site and season are very important considerations.

Occupancy models let us estimate metapopulation parameters when detection is imperfect.

There are no individual-level data or parameters.

These methods are often easy to implement over large areas and so are used in monitoring programs.

Definition of site and season are very important considerations.

Models can be used in other contexts, such as when a site is a human, and we are interested in proportion of people with some disease.