

# Sampling and Estimation





Fig 5.5



## Key concepts

- All models include parameters (e.g.,  $N$ ,  $r$ ,  $K$ , etc. . . )

## Key concepts

- All models include parameters (e.g.,  $N$ ,  $r$ ,  $K$ , etc. . . )
- Parameters are almost never known. Why?

## Key concepts

- All models include parameters (e.g.,  $N$ ,  $r$ ,  $K$ , etc. . . )
- Parameters are almost never known. Why?
  - ▶ We usually have to sample
  - ▶ Animals are hard to detect

## Key concepts

- All models include parameters (e.g.,  $N$ ,  $r$ ,  $K$ , etc. . . )
- Parameters are almost never known. Why?
  - ▶ We usually have to sample
  - ▶ Animals are hard to detect
- Good sampling designs yield accurate estimates of unknown parameters

## **Parameter**

A characteristic of a population

## **Parameter**

A characteristic of a population

## **Statistic**

A characteristic of a dataset (often used to estimate a parameter)

## Parameter

A characteristic of a population

## Statistic

A characteristic of a dataset (often used to estimate a parameter)

	Population parameter	Parameter estimate
Population size	$N_t$	$\hat{N}_t$
Growth rate	$r$	$\hat{r}$
Occurrence probability	$\psi$	$\hat{\psi}$



## **Properties of a good design:**

## Properties of a good design:

- (1) Clearly defined objective, in terms of:
  - ▶ Parameter(s) that will be estimated
  - ▶ Population of interest
  - ▶ Criteria for reliability (e.g., precision)
  - ▶ Practical constraints (e.g., costs)

## Properties of a good design:

- (1) Clearly defined objective, in terms of:
  - ▶ Parameter(s) that will be estimated
  - ▶ Population of interest
  - ▶ Criteria for reliability (e.g., precision)
  - ▶ Practical constraints (e.g., costs)

**Bad objective:** “Quantify the abundance of tigers in India”

# HOW DO YOU GET ACCURATE ESTIMATES?

## Properties of a good design:

- (1) Clearly defined objective, in terms of:
  - ▶ Parameter(s) that will be estimated
  - ▶ Population of interest
  - ▶ Criteria for reliability (e.g., precision)
  - ▶ Practical constraints (e.g., costs)

**Bad objective:** “Quantify the abundance of tigers in India”

**Better objective:** “Obtain an unbiased estimate of tiger density on the Corbett Tiger Reserve during July 2009, with a coefficient of variation  $< 25\%$  for under \$5,000.”

# HOW DO YOU GET ACCURATE ESTIMATES?

## Properties of a good design:

- (1) Clearly defined objective, in terms of:
  - ▶ Parameter(s) that will be estimated
  - ▶ Population of interest
  - ▶ Criteria for reliability (e.g., precision)
  - ▶ Practical constraints (e.g., costs)
- (2) Replication

**Bad objective:** “Quantify the abundance of tigers in India”

**Better objective:** “Obtain an unbiased estimate of tiger density on the Corbett Tiger Reserve during July 2009, with a coefficient of variation  $< 25\%$  for under \$5,000.”

# HOW DO YOU GET ACCURATE ESTIMATES?

## Properties of a good design:

- (1) Clearly defined objective, in terms of:
  - ▶ Parameter(s) that will be estimated
  - ▶ Population of interest
  - ▶ Criteria for reliability (e.g., precision)
  - ▶ Practical constraints (e.g., costs)
- (2) Replication
- (3) Randomization

**Bad objective:** “Quantify the abundance of tigers in India”

**Better objective:** “Obtain an unbiased estimate of tiger density on the Corbett Tiger Reserve during July 2009, with a coefficient of variation  $< 25\%$  for under \$5,000.”

# HOW DO YOU GET ACCURATE ESTIMATES?

## Properties of a good design:

- (1) Clearly defined objective, in terms of:
  - ▶ Parameter(s) that will be estimated
  - ▶ Population of interest
  - ▶ Criteria for reliability (e.g., precision)
  - ▶ Practical constraints (e.g., costs)
- (2) Replication
- (3) Randomization
- (4) Controls (when conducting an experiment)

**Bad objective:** “Quantify the abundance of tigers in India”

**Better objective:** “Obtain an unbiased estimate of tiger density on the Corbett Tiger Reserve during July 2009, with a coefficient of variation  $< 25\%$  for under \$5,000.”

## **Target population**

The population of interest. Should be defined in terms of time and space.



## **Target population**

The population of interest. Should be defined in terms of time and space.

## **Sampled population**

The sampled portion of the population of interest, usually defined in terms of the sample units (such as plots, quadrats, etc.).


## Accuracy has two components:

- (1) **Bias:** The difference between the average estimate and the true parameter
- (2) **Variance:** The variability of the estimates.

## Accuracy has two components:

- (1) **Bias:** The difference between the average estimate and the true parameter
- (2) **Variance:** The variability of the estimates.

Fig 5.5



# HOW DO WE REDUCE VARIANCE?


Huge sample size

# EFFECT OF SAMPLE SIZE


Suppose we want to estimate the height of students on campus, and we have enough resources to repeat a survey many times.

Each point below is an estimate.

**Sample size of 10**



**Sample size of 100**




# EFFECT OF SAMPLE SIZE


Suppose we want to estimate the height of students on campus, and we have enough resources to repeat a survey many times.

Each point below is an estimate.

**Sample size of 10**



**Sample size of 100**




# EFFECT OF SAMPLE SIZE


Suppose we want to estimate the height of students on campus, and we have enough resources to repeat a survey many times.

Each point below is an estimate.

**Sample size of 10**



**Sample size of 100**






# EFFECT OF SAMPLE SIZE


Suppose we want to estimate the height of students on campus, and we have enough resources to repeat a survey many times.

Each point below is an estimate.

**Sample size of 10**



**Sample size of 100**




# EFFECT OF SAMPLE SIZE


Suppose we want to estimate the height of students on campus, and we have enough resources to repeat a survey many times.

Each point below is an estimate.

**Sample size of 10**



**Sample size of 100**




# EFFECT OF SAMPLE SIZE


Suppose we want to estimate the height of students on campus, and we have enough resources to repeat a survey many times.

Each point below is an estimate.

**Sample size of 10**



**Sample size of 100**




# EFFECT OF SAMPLE SIZE


Suppose we want to estimate the height of students on campus, and we have enough resources to repeat a survey many times.

Each point below is an estimate.

**Sample size of 10**



**Sample size of 100**




# EFFECT OF SAMPLE SIZE


Suppose we want to estimate the height of students on campus, and we have enough resources to repeat a survey many times.

Each point below is an estimate.

**Sample size of 10**




**Sample size of 100**



## EFFECT OF SAMPLE SIZE

Suppose we want to estimate the height of students on campus, and we have enough resources to repeat a survey many times.

Each point below is an estimate.




# EFFECT OF SAMPLE SIZE


Suppose we want to estimate the height of students on campus, and we have enough resources to repeat a survey many times.

Each point below is an estimate.

**Sample size of 10**



**Sample size of 100**




# EFFECT OF SAMPLE SIZE


Suppose we want to estimate the height of students on campus, and we have enough resources to repeat a survey many times.

Each point below is an estimate.

**Sample size of 10**



**Sample size of 100**



The standard deviation of the sampling distribution is called the standard error (SE)




# HOW DO WE REDUCE BIAS?

## Randomization


## Simple random sampling

- All sample units have the same inclusion probability




## Simple random sampling

- All sample units have the same inclusion probability
- Easiest and most reliable method




## Simple random sampling

- All sample units have the same inclusion probability
- Easiest and most reliable method
- But not always cost effective




## Stratified random sampling

- Useful when study area is characterized by several homogeneous regions




## Stratified random sampling

- Useful when study area is characterized by several homogeneous regions
- Regions with higher variability should be sampled more intensively than regions with low variability



## Stratified random sampling

- Useful when study area is characterized by several homogeneous regions
- Regions with higher variability should be sampled more intensively than regions with low variability
- Often more cost effective than simple random sampling






## Systematic sampling

- Sample units are selected according to a regular, ordered scheme with the first unit being sampled randomly.


(E)



## Systematic sampling


- Sample units are selected according to a regular, ordered scheme with the first unit being sampled randomly.
- Easy to implement in the field

(E)



## Systematic sampling

- Sample units are selected according to a regular, ordered scheme with the first unit being sampled randomly.
- Easy to implement in the field
- Potentially dangerous because sample unit spacing could coincide with natural spacing of environmental features



## Main points

- We have to estimate model parameters
- Reliable estimates require good sampling design
- Replication reduces variance
- Randomization reduces bias