

# A Differential Geometry Basics Cheat Sheet

By Nick Konz

*This is designed to be a quick, yet rigorous introduction/reference to the basic principles found within differential geometry, covering only the bare minimum of topics needed. Largely adapted from Spivak's A Comprehensive Introduction to Differential Geometry, 3rd Edition and Mendelson's Introduction to Topology, 3rd Edition*

## 1 Prerequisite Knowledge

We only assume basic knowledge of continuity and differentiability in the context of functions and topology. Here,  $\forall$  means “for all” or “for each”, and  $\exists$  means “there exists”.

### 1.1 Some Basic Topology

**Definition 1.1. Neighborhood.** Let  $a \in \mathbb{R}^n, \delta > 0$ . The  $\delta$ -**neighborhood** of  $a$  is the set

$$U(a, \delta) = \{x \in \mathbb{R}^n : \|x - a\| < \delta\}. \quad (1)$$

**Definition 1.2. Metric Space.** Let  $X$  be some non-empty set and  $d$  be the mapping/function  $d : X \times X \rightarrow \mathbb{R}$ . The pair  $(X, d)$  is a **metric space** if,  $\forall x, y, z \in X$ ,

1.  $d(x, y) \geq 0$
2.  $d(x, y) = 0$  if and only if (iff)  $x = y$
3.  $d(x, y) = d(y, x)$
4.  $d(x, z) \leq d(x, y) + d(y, z)$  (*triangle inequality*).

**Definition 1.3. Topological Space.** Let  $X$  be a non-empty set and  $\mathcal{J}$  be a collection of subsets of  $X$ .  $(X, \mathcal{J})$  is a **topological space** if

1.  $X \in \mathcal{J}$
2.  $\emptyset \in \mathcal{J}$ , where  $\emptyset$  is the empty set
3. If  $O_1, O_2, \dots, O_n \in \mathcal{J}$ , then  $\bigcap_{i=1}^n O_i \in \mathcal{J}$
4. If  $\forall \alpha \in I, O_\alpha \in \mathcal{J}$ , then  $\bigcup_{\alpha \in I} O_\alpha \in \mathcal{J}$   
( $I$  is some *indexing set*).

We label  $X$  as the **underlying set**,  $\mathcal{J}$  as the **topology** on  $X$ , and members of  $\mathcal{J}$  as **open sets**.

*Remark.* Topological and metric spaces  $(X, \mathcal{J})$  and  $(X, d)$ , respectively, are sometimes notated simply as  $X$ .

**Definition 1.4. Homeomorphism.** Let  $(X, \mathcal{J})$  and  $(Y, \mathcal{K})$  be topological spaces.  $(X, \mathcal{J})$  and  $(Y, \mathcal{K})$  are **homeomorphic** if  $\exists$  inverse functions, or **homeomorphisms**,  $f : X \rightarrow Y$  and  $g : Y \rightarrow X$  such that  $f, g$  are continuous.

## 2 Manifolds

**Definition 2.1. Manifold.** A **manifold** is a metric space  $M$  such that if  $x \in M$ ,  $\exists$  some neighborhood  $U$  of  $x$  and some  $n \in \{0, 1, 2, \dots\}$  such that  $U$  is homeomorphic to  $\mathbb{R}^n$ . If  $\exists$  such an  $n$  that is the same  $\forall x \in M$ , we say that  $M$  is  **$n$ -dimensional**, which can be notated as  $M^n$ .

*Remark.* Think of a manifold as being a surface that is locally Euclidean.

**Definition 2.2.  $C^\infty$ -related Homeomorphisms.** Let  $M$  be some manifold, and let  $U, V$  be open subsets of  $M$ . Two homeomorphisms  $x : U \rightarrow x(U) \subset \mathbb{R}^n$  and  $y : V \rightarrow y(V) \subset \mathbb{R}^n$  (for some  $n$ ) are  **$C^\infty$ -related** if the maps

$$y \circ x^{-1} : x(U \cap V) \rightarrow y(U \cap V) \quad (2)$$

$$x \circ y^{-1} : y(U \cap V) \rightarrow x(U \cap V) \quad (3)$$

are infinitely differentiable, or  **$C^\infty$** .

**Definition 2.3. Atlas.** A family of *mutually*  $C^\infty$ -related homeomorphisms whose domains cover  $M$  (i.e. their union equals  $M$ ) is an **atlas** of  $M$ .

**Definition 2.4. Chart/Coordinate System.** A **chart** or **coordinate system** for some manifold  $M$  is a homeomorphism  $x$  from some open  $U \in M$  to an open subset of  $\mathbb{R}^n$ , denoted  $(x, U)$ . A chart of  $M$  is a member of some atlas of  $M$ .

*Remark.* Charts/coordinate systems  $(x, U)$  create a way of assigning coordinates to points on  $U$ , and are sometimes notated simply with  $x$ .

**Definition 2.5. Differentiable Manifold.** A **differentiable, smooth** or  **$C^\infty$  manifold** is a pair  $(M, \mathcal{A})$ , where  $M$  is some manifold, and  $\mathcal{A}$  is some *maximal atlas* for  $M$ , i.e. the union of all possible atlases of  $M$ .